

# Molecular characterization of adenovirus circulating in Central and South America during the 2006–2008 period

Josefina García<sup>a</sup>, Merly Sovero<sup>a</sup>, Víctor Alberto Laguna-Torres<sup>a</sup>, Jorge Gomez<sup>b</sup>, Wilson Chicaiza<sup>c</sup>, Melvin Barrantes<sup>d</sup>, Felix Sanchez<sup>e</sup>, Mirna Jimenez<sup>f</sup>, Guillermo Comach<sup>g</sup>, Ivette L. de Rivera<sup>h</sup>, Roberto Agudo<sup>i</sup>, Ana E. Arango<sup>j</sup>, Alma Barboza<sup>k</sup>, Nicolas Aguayo<sup>k</sup>, Tadeusz J. Kochel<sup>a</sup>

<sup>a</sup>US Naval Medical Research Center Detachment, Lima, Peru. <sup>b</sup>Dirección General de Epidemiología, Ministerio de Salud, Lima, Perú. <sup>c</sup>Hospital Vozandes, Quito, Ecuador. <sup>d</sup>Hospital Solano, Buenos Aires, Argentina. <sup>e</sup>Hospital Infantil Manuel de Jesus Rivera, Managua, Nicaragua. <sup>f</sup>Hospital Nacional de Metapan, Metapan, El Salvador. <sup>g</sup>LARDIDEV-Biomed-UC, Maracay, Venezuela. <sup>h</sup>Universidad Nacional Autónoma de Honduras, Tegucigalpa, Honduras. <sup>i</sup>Dirección General de Epidemiología, Ministerio de Salud, Cochabamba, Bolivia. <sup>j</sup>Universidad de Antioquia, Medellín, Colombia. <sup>k</sup>ONG Rayos de Sol, Asuncion, Paraguay

Correspondence: Josefina García, US Naval Medical Research Center Detachment, Lima, Peru. E-mail: josefina.garcia@med.navy.mil

Accepted 18 August 2009. Published Online 5 October 2009.

**Background** Human Adenoviruses are recognized pathogens, causing a broad spectrum of diseases. Serotype identification is critical for epidemiological surveillance, detection of new strains and understanding of HAdvs pathogenesis. Little data is available about HAdvs subtypes in Latin America.

**Methods** In this study, we have molecularly characterized 213 adenoviruses collected from ILI presenting patients, during 2006–08, in Central and South America.

**Results** Our results indicate that 161(76%) adenoviruses belong to subgroup C, 45 (21%) to subgroup B and 7 (3%) to subtype E4.

**Keywords** Adenovirus, Latin America, viruses.

Please cite this paper as: García *et al.* (2009) Molecular characterization of adenovirus circulating in Central and South America during the 2006–2008 period. *Influenza and Other Respiratory Viruses* 3(6), 327–330.

## Introduction

Human adenoviruses (HAdv) are well-recognized pathogens, causing a broad spectrum of diseases, including upper and lower respiratory tract infections, gastroenteritis, conjunctivitis, and keratoconjunctivitis, mostly in young children.

Adenoviruses are non-enveloped, double-stranded DNA viruses that cause an estimated 8% of clinically relevant viral disease globally.<sup>1</sup> To date, 51 human adenoviral serotypes have been described on the basis of type-specific anti-serum-mediated neutralization of infectivity and grouped in six species or subgroups (A, B, C, D, E, and F), on the basis of hemagglutination inhibition and biochemical criteria.<sup>2</sup> Each species has been associated with different spectra of disease: respiratory infections with HAdvs are usually associated with species B, C, and E whereas species F viruses are restricted to the enteric tract infections.<sup>3,4</sup>

Serotype identification is critical for epidemiological surveillance, detection of new strains and understanding of HAdv pathogenesis. The use of molecular methods for adenovirus diagnosis has significantly increased in the last years, although they have some disadvantages as the

homology within species ranges from 50% to 100%, but between subgroups it can go as low as 4%.<sup>5</sup>

Little data are available about HAdvs subtypes circulating in Latin America. There are few studies showing the different proportions of each species and/or subtypes in this region.<sup>6–9</sup> In this study, we have analyzed samples from patients presenting with influenza-like illness (ILI) in Central and South America during the 3-year period 2006–2008 by amplification of the last 300 nucleotides of the hexon gene and comparison to published sequences, to classify adenovirus into species in outpatients with ILI symptoms in this region.

## Material and methods

### Specimen collection and identification of adenovirus positive samples


In collaboration with 10 Central and South American countries ILI-surveillance networks, 14 106 nasopharyngeal and throat swab were collected from ILI presenting subjects, regardless of age, with an onset of fever ( $\geq 38^{\circ}\text{C}$ ) and cough or sore throat during 2006–2008 under the Surveillance for

Respiratory Pathogens Protocol, NMRC.D.2002.0019.<sup>10</sup> Adenoviruses were isolated from ILI specimens from: Nicaragua ( $n = 21$ ), Honduras ( $n = 19$ ), El Salvador ( $n = 1$ ), Venezuela ( $n = 7$ ), Colombia ( $n = 27$ ), Bolivia ( $n = 3$ ), Ecuador ( $n = 17$ ), Peru ( $n = 267$ ), Paraguay ( $n = 1$ ), and Argentina ( $n = 39$ ). Most of the samples came from Peru, thus the number of adenovirus is greater in this country, but the proportion of adenovirus by ILI cases remains low (2.8%). Once collected, swabs were placed in viral transport media and stored at  $-80^{\circ}\text{C}$  until transported on dry ice to the NMRC

in Lima, Peru. Virus isolation was carried out by inoculation in Vero cell line for 10 days. Then, viral identification was performed by indirect immunofluorescence using specific anti-sera (D3 Ultra DFA Respiratory Virus Screening & ID Kit; Diagnostic Hybrids, Athens, OH, USA).

#### DNA extraction and polymerase-chain reaction (PCR)

Viral DNA was extracted from immunofluorescence positive samples using the AmpliTaq Gold<sup>®</sup> kit (Applied



**Figure 1.** Adenovirus-phylogenetic tree. The last 300 nucleotides of the adenovirus *hexon* gene were amplified, sequenced, and compared with published sequences from GenBank. We have labeled the samples according to the following format: 'Country of collection/Month-Year of collection.' The comparison sequences are complete genome sequences from GenBank, these are presented in the following format: 'Accession Number/Serotype in Bold' (Number of isolated virus sharing the same sequence). Nucleotide sequences were aligned by using clustal x. Phylogenetic analyses were performed using the Kimura two-parameter model as a model of nucleotide substitution and using the neighbor-joining method to reconstruct phylogenetic trees (mega version 2.1). The samples are grouped into species. Canine adenoviruses were used as out-group sequences for comparison.

**Table 1.** Adenoviral subgroups in circulation

Country	HAdv B	HAdv C	HAdv E	Total	% of adenovirus isolates/ILI cases
El Salvador	0	1	0	1	0.38
Honduras	1	11	0	12	5.28
Nicaragua	2	9	0	11	2.23
Venezuela	3	3	0	6	1.72
Colombia	3	6	0	9	4.29
Ecuador	2	7	0	9	1.81
Peru	32	114	7	153	2.84
Paraguay	0	1	0	1	0.43
Argentina	2	9	0	11	6.43
Total	45	161	7	213	

Number of adenovirus from each subgroup (B, C, and E) determined by molecular analysis and sequencing by country of collection. The last column presents the percentage of adenovirus isolations among the total ILI cases by country.

ILI, influenza-like illness; HAdv, human adenoviruses.

Biosystems, Foster City, CA, USA) and tested by PCR. An amplicon of a moderately conserved region of the adenovirus Hexon gene (amino acid 900–971) was amplified with the following specific primers: Adeno 3 (5'-CCTTTGG CGCATCCCATCT-3') and Adeno 4 (5'-GCGCTTGTCATAGGTGCCCA-3').<sup>11</sup>

### Sequencing and phylogenetic analysis

For direct sequencing of viral nucleic acids from clinical specimens, genes fragments were amplified and sequenced with the use of Big Dye terminator cycle sequencing kit (version 3.1; Applied Biosystems, Ann Arbor, MI, USA) on a Genetic Analyser system (version 3130xL; Applied Biosystems).

Gene sequences were assembled aligned and edited using sequencer and bioedit (version 7.0.0; Isis Pharmaceuticals, Inc., Dublin, Ireland) software. Phylogenetic trees were generated with clustal x version 2.0.1 and mega version 3.1 software.<sup>12</sup>

## Results and discussion

Influenza-like illnesses surveillance programs help to monitor the progression and distributions of many viruses and may be used to alert early changes in strains causing differences in infectivity or severity of infections and possible outbreaks.

Molecular typing provides a rapid identification of almost any serotype belonging to the Adv species A, B, C, D, E, and F. Many studies about the phylogenetic evolution of adenovirus exist focusing mainly on the *hexon* gene, with different results depending on the region chosen. We chose the last 300 nucleotides of this gene, which is a moderately conserved region common to all HAdv subgroups.

A total of 402 HAdv were isolated from the ILI-surveillance-network. 213 adenoviral samples were selected trying to include all listed countries; amplified and sequenced, subgroups and subtypes were inferred by comparison of sequenced samples with published ones from GenBank. Table 1 shows the analysis of subgroups B, C, and E by country (no other subgroups were detected), and we can see that during the 2006–2008 period, there were mostly two adenoviral species in circulation: C (76%) and B (21%). Subgroup E was only seen in Peru after August 2008 and represents 3% of the total HAdv found. This distribution agrees with recently published studies<sup>4,13,14</sup> but differs from previous studies showing that the predominant species circulating in Latin America was B.<sup>7,13,15,16</sup>

Some studies have reported that adenoviral outbreaks are seasonal, during a 3-year period we did not observe any seasonality pattern on adenoviral infectivity by subgroups or subtypes.

Finally, Figure 1 shows the phylogenetic tree of all sequenced HAdv, where the most represented subtype is HAdv C2, followed by HAdv C1 and then by B species, subtypes 7 and 3.

As this is passive surveillance study, the collected data does not allow the analysis of incidence rates or disease burden, it is a descriptive view of respiratory specimens from symptomatic cases. We are currently planning additional studies to address these issues.

## Acknowledgements

We are very grateful to Ms. María Ester Gamero for cell culture and viral isolation support. We also thank Dr. Patricia Aguilar and Dr. Brett Forshey for critical reading of this manuscript.

## Funding statement

This study was funded by the United States Department of Defense Global Emerging Infections Systems Research Program, Work Unit Number: 847705.82000.25GB.B0016.

## Conflict of interest statement

We declare that we have no conflict of interest.

## Disclaimers

The views expressed in this article are those of the authors and do not necessarily reflect the official policy or position of the Department of the Navy, Department of Defense, nor the U.S. Government.

The study protocol was approved by the Naval Medical Research Center Institutional Review Board (Protocols NMRC.D.2002.0019) in compliance with all applicable Federal regulations governing the protection of human subjects.

## Disclosure

None of the authors has a financial or personal conflict of interest related to this study. The corresponding author had full access to all data in the study and final responsibility for the decision to submit this publication.

## Copyright statement

Authors Tadeusz J. Kochel is a military service member, Alberto Laguna-Torres and Merly Sovero are employees of the U.S. Government. This work was prepared as part of their official duties. Title 17 U.S.C. § 105 provides that 'Copyright protection under this title is not available for any work of the United States Government'. Title 17 U.S.C. § 101 defines a U.S. Government work as a work prepared by a military service members or employees of the U.S. Government as part of those person's official duties.

## References

- 1 Rux JJ, Burnett RM. Adenovirus structure. *Hum Gene Ther* 2004; 15:1167–1176.
- 2 Russell WC. Adenoviruses: update on structure and function. *J Gen Virol* 2009; 90:1–20.
- 3 Echavarría M. Adenoviruses in immunocompromised hosts. *Clin Microbiol Rev* 2008; 21:704–715.
- 4 Echavarría M, Maldonado D, Elbert G, Videla C, Rappaport R, Carballal G. Use of PCR to demonstrate presence of adenovirus species B, C, or F as well as coinfection with two adenovirus species in children with flu-like symptoms. *J Clin Microbiol* 2006; 44:625–627.
- 5 Schilham MW, Claas EC, van Zaane W *et al.* High levels of adenovirus DNA in serum correlate with fatal outcome of adenovirus infection in children after allogeneic stem-cell transplantation. *Clin Infect Dis* 2002; 35:526–532.
- 6 de Azevedo JP, Nascimento LR, Cortinovis MC, Oliveira SS, da Costa EV, da Silva EE. Characterization of species B adenoviruses isolated from fecal specimens taken from poliomyelitis-suspected cases. *J Clin Virol* 2004; 31:248–252.
- 7 Kajon AE, Mistchenko AS, Videla C, Hortal M, Wadell G, Avendano LF. Molecular epidemiology of adenovirus acute lower respiratory infections of children in the south cone of South America (1991–1994). *J Med Virol* 1996; 48:151–156.
- 8 Moura PO, Roberto AF, Hein N *et al.* Molecular epidemiology of human adenovirus isolated from children hospitalized with acute respiratory infection in Sao Paulo, Brazil. *J Med Virol* 2007; 79:174–181.
- 9 Palomino MA, Larranaga C, Villagra E, Camacho J, Avendano LF. Adenovirus and respiratory syncytial virus-adenovirus mixed acute lower respiratory infections in Chilean infants. *Pediatr Infect Dis J* 2004; 23:337–341.
- 10 Laguna-Torres VA, Gomez J, Ocana V *et al.* Influenza-like illness sentinel surveillance in Peru. *PLoS ONE* 2009; 4:e6118.
- 11 Hierholzer JC, Halonen PE, Dahlen PO, Bingham PG, McDonough MM. Detection of adenovirus in clinical specimens by polymerase chain reaction and liquid-phase hybridization quantitated by time-resolved fluorometry. *J Clin Microbiol* 1993; 31:1886–1891.
- 12 Kumar S, Tamura K, Nei M. MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform* 2004; 5:150–163.
- 13 Frabasile S, Vituriera N, Perez G, Mateos S, Arbiza J. Genotyping of Uruguayan human adenovirus isolates collected between 1994 and 1998. *Acta Virol* 2005; 49:129–132.
- 14 Moura FE, Mesquita JR, Portes SA, Ramos EA, Siqueira MM. Antigenic and genomic characterization of adenovirus associated to respiratory infections in children living in Northeast Brazil. *Mem Inst Oswaldo Cruz* 2007; 102:937–941.
- 15 Carballal G, Videla C, Misirlian A, Requeijo PV, Aguilar Mdel C. Adenovirus type 7 associated with severe and fatal acute lower respiratory infections in Argentine children. *BMC Pediatr* 2002; 2:6.
- 16 Larranaga C, Kajon A, Villagra E, Avendano LF. Adenovirus surveillance on children hospitalized for acute lower respiratory infections in Chile (1988–1996). *J Med Virol* 2000; 60:342–346.