

On the behavior of the $n_1(K, T)_{n_2}^A$ series of doubly excited states in He-like atoms immersed in weakly coupled plasmas

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2014 J. Phys.: Conf. Ser. 488 062010

(<http://iopscience.iop.org/1742-6596/488/6/062010>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 200.24.16.228

This content was downloaded on 24/01/2017 at 19:04

Please note that [terms and conditions apply](#).

You may also be interested in:

[Resonances of a hydrogen atom in strong parallel electric and magnetic fields using B-spline basis sets](#)

Zhang Yue-Xia, Meng Hui-Yan and Shi Ting-Yun

[Determination of Nonlinear Resonance Parameters in Electron Storage Rings](#)

Tsukasa Miyajima and Yukinori Kobayashi

[Photodetachment cross sections of the positronium negative ion](#)

Akinori Igarashi, Isao Shimamura and Nobuyuki Toshima

[Factorized cumulant expansion for homogeneous turbulence](#)

Tomomasa Tatsumi, Michio Yamada and Toshifumi Takei

[A new moving coil loud speaker](#)

[Feshbach-like study of He resonant states in Debye plasmas using explicitly correlated wave functions](#)

J C Cardona, A F Ordoñez and J L Sanz-Vicario

[Computation of resonant states using explicitly correlated coordinates in Be-like B, C and N](#)

J C Cardona

On the behavior of the $n_1(K, T)_{n_2}^A$ series of doubly excited states in He-like atoms immersed in weakly coupled plasmas

A. F. Ordóñez*, J. C. Cardona† and J. L. Sanz-Vicario*¹

* Grupo de Física Atómica y Molecular. Instituto de Física, Universidad de Antioquia, Medellín, Colombia.

† GEOEL, Universidad del Atlántico, Barranquilla, Colombia.

Synopsis The stability of resonant states in Helium-like atoms while under the influence of a plasma environment is studied. We analyze the variation of resonance parameters (energies and lifetimes) against the strength of the screening parameter within the Debye-Hückel model. We use a Feshbach-like projection formalism with explicitly correlated CI-wave functions to uncover the evolution of resonance parameters until they cross the upper A^{9+} ($N=2$) threshold, and a complex scaling method to analyze the survival of these *Feshbach* resonances across the threshold, which eventually makes them behave as *shape* resonances.

The interest in the fundamental problem of atoms immersed in plasmas stems from its potential application in atomic physics, plasma physics, solid-state physics, and astrophysics. Plasma embedded He-like atoms represent here also a benchmark to understand the physics involved. The Hamiltonian for a two-electron atom immersed into a weakly coupled plasma reads

$$\mathcal{H} = -\frac{\nabla_1^2}{2} - \frac{\nabla_2^2}{2} - \frac{Z \exp(-r_1/D)}{r_1} - \frac{Z \exp(-r_2/D)}{r_2} + \frac{\exp(-r_{12}/D)}{r_{12}}. \quad (1)$$

Here, the Debye length $D=(k_B T/4\pi e^2 \rho)^{1/2}$ (a function of the plasma electron temperature T and density ρ) characterizes the interaction strength of the atomic system with the surrounding plasma. We use a Hylleraas configuration interaction (HyCI) method to compute accurate energies and wave functions, the latter consisting of antisymmetrized correlated configurations $\psi_i(\mathbf{x}_1, \mathbf{x}_2)=\mathcal{A}\{\phi(\mathbf{r}_1, \mathbf{r}_2)\chi(s_1, s_2)\}$ where the spatial part of the wave function is built in terms of correlated Slater type orbitals $\phi = r_1^{n_1} r_2^{n_2} r_{12}^{n_{12}} e^{-\alpha r_1 - \beta r_2} \mathcal{Y}_{L,M}(\Omega_1, \Omega_2)$. Their advantage is that all integrals required for the solution of the plasma free ($D=\infty$) as well as for finite values of D may be calculated in closed form [1, 2].

We have also implemented a Feshbach-like projection method (at variance with previous extensive work by Ho *et al* [3] that uses the stabilization method) by solving separately the resonant $QH\mathcal{Q}$ (resonance energies), the non-resonant $\mathcal{P}H\mathcal{P}$ (continuum discretized energies) eigenvalue problems and the $\mathcal{Q}H\mathcal{P}$ couplings (from which the Auger widths are extracted). For instance, we perform a systematic study on the behavior of resonance parameters (energies, life-

times and interelectronic angle) of the lowest series of plasma-embedded He $1,3S^e$, $1,3P^o$ and $1,3D^e$ doubly excited states located below the $\text{He}^+(N=2)$ ionization threshold, as a function of the Debye characteristic length D . At variance with one-electron atoms (where *shape* resonance widths vary monotonically with the screening strength) the evolution of the Auger width with respect to screening is found to be different for each series of $(K, T)^A$ Herrick-Sinanoğlu-Lin pseudo-quantum numbers, until resonances cross the upper $\text{He}^+(2s, 2p)$ thresholds, then merging into new open electronic continuum channels. The geometrical distribution of the electron pair as a function of the screening is also analyzed through the interelectronic angle for each resonance state. In addition, to shed light on the survival of Feshbach resonances (below the threshold) transforming into shape resonances (above the threshold) we also implement the complex rotation method for the problem at hand. This transformation of the resonant character across the threshold [4] may involve dramatic changes in the wave function structure and particularly in the Auger widths.

References

- [1] R.J. Tweed (1972) *J. Phys. B: At. Mol. Phys.*, **5**, 810; J.C. Cardona and J.L. Sanz-Vicario (2008) *J. Phys. B: At. Mol. Opt. Phys.*, **41**, 055003
- [2] J.L. Calais and P.-O. Löwdin (1962) *J. Mol. Spectrosc.*, **8**, 203
- [3] S. Kar and Y.K. Ho (2005) *Chem. Phys. Lett.*, **402**, 544; (2009) *J. Phys. B: At., Mol. Opt. Phys.*, **42**, 044007
- [4] S.B. Zhang, J.G. Wang and R.K. Janev (2010) *Phys. Rev. Lett.*, **104**, 023203; (2010) *Phys. Rev. A*, **81**, 032707

¹E-mail: sanjose@fisica.udea.edu.co

