

Original Article

Obesity and Metabolic Syndrome

Diabetes & Metabolism Journal 2017;41(4):303-315.

Published online: May 24, 2017

DOI: <https://doi.org/10.4093/dmj.2017.41.4.303>


In Vitro Effect of Fatty Acids Identified in the Plasma of Obese Adolescents on the Function of Pancreatic β -Cells

[Claudia Velasquez](#)¹, [Juan Sebastian Vasquez](#)², [Norman Balcazar](#)^{2,3}  

¹Research Group in Food and Human Nutrition, School of Dietetics and Human Nutrition, University of Antioquia, Medellin, Colombia.

²Genetics Molecular Group, University of Antioquia, Medellin, Colombia.

³Department of Physiology and Biochemistry, School of Medicine, University of Antioquia, Medellin, Colombia.

 Corresponding author: Norman Balcazar. Genetics Molecular Group and Department of Physiology and Biochemistry, School of Medicine, University of Antioquia, Calle 62 #52-59, Medellin 1226, Colombia. norman.balcazar@udea.edu.co

Received September 29, 2016 Accepted January 31, 2017

Copyright © 2017 Korean Diabetes Association

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background

The increase in circulating free fatty acid (FFA) levels is a major factor that induces malfunction in pancreatic β -cells. We evaluated the effect of FFAs reconstituted according to the profile of circulating fatty acids found in obese adolescents on the viability and function of the murine insulinoma cell line (mouse insulinoma [MIN6]).

Methods

From fatty acids obtained commercially, plasma-FFA profiles of three different youth populations were reconstituted: obese with metabolic syndrome; obese without metabolic syndrome; and normal weight without metabolic syndrome. MIN6 cells were treated for 24 or 48 hours with the three FFA profiles, and glucose-stimulated insulin secretion, cell viability, mitochondrial function and antioxidant activity were evaluated.

Results

The high FFA content and high polyunsaturated ω 6/ ω 3 ratio, present in plasma of obese adolescents with metabolic syndrome had a toxic effect on MIN6 cell viability and function, increasing oxidative stress and decreasing glucose-dependent insulin secretion.

Conclusion

These results could help to guide nutritional management of obese young individuals, encouraging the increase of ω -3-rich food consumption in order to reduce the likelihood of deterioration of β -cells and the possible development of type 2 diabetes mellitus.

Keywords: [Fatty acids, nonesterified](#); [Insulin-secreting cells](#); [Lipotoxicity](#); [Metabolic syndrome](#); [Obesity](#)

INTRODUCTION

Overweight and the associated metabolic disorders affect more people in the world than any other condition [1]. Metabolic syndrome (MetS) describes the principal clinical manifestations generated by obesity, and it can be more precisely referred to as a chronic immune-MetS, that includes changes such as hyperglycaemia, insulin resistance (IR), atherogenic dyslipidaemia, fatty liver, hypercoagulability, arterial hypertension, neurodegeneration, and autoimmunity [2].

Close to 90% of individuals with type 2 diabetes mellitus (T2DM) are overweight and present MetS [3]. Abdominal (central) obesity principally occurs because of hypertrophy of the visceral adipocytes and is strongly associated with MetS [4]. The dysfunction of adipocyte populations that are in some way “sensitive” to cellular lesions due to over-eating is recognized as one of the initial factors in the development of MetS.

One of the connections between abdominal obesity and the metabolic changes of MetS could be circulating free fatty acids (FFAs) [5]. Adipocytes in adipose tissues are capable of activating proinflammatory intracellular pathways during their development towards hypertrophy in the context of over-eating. With time, a state of IR and local inflammation is established, which favours the release of excess FFAs that progress to different tissues that are not metabolically designed to store FFAs. Some fatty acids released by inflamed visceral adipose tissue can promote an atherogenic state, altering the secretion of insulin or concomitantly generating IR in multiple tissues, especially muscle, liver, and subcutaneous adipose tissues [6].

This inflammatory state and cytotoxic serum lipid profiles can induce chronic damage in the β -cells of the pancreas and loss of the insulin-secreting function. The effect of FFAs on β -cell function is complex and pleiotropic; it depends on the chemical nature, concentration, exposure time, and interaction with other nutrients. For example, culturing cells or islets for 24 or 48 hours with palmitic or stearic acid reduces glucose-stimulated insulin secretion (GSIS), induces apoptosis and reduces cell viability. However, coincubation with oleic or linoleic acid does not induce cell damage, and this treatment combined with palmitic or stearic acid also restores GSIS. These differential effects of FFAs, some with protective potential and others with marked cytotoxicity, show that serum lipid levels are key factors in the progression from MetS to T2DM [7,8,9].

In adults, the data available on the FFA profile of obese persons with MetS are affected by lifestyle variables, such as alcohol consumption, smoking, or comorbidities. Children and adolescents present a disturbing increase in the incidence of obesity; this age range also offers advantages for the study of serum lipids due to the shorter time required to evaluate the effects of

confounding lifestyle factors. For this reason, evaluating the effect of FFAs in adolescents with immune-metabolic changes on the function and viability of β -cells is highly relevant both for clinical applications and at the basic research level. We specifically study adolescents who have a very high risk of developing T2DM. It is important to determine the overall effect of each FFA mixture on the function of the pancreatic β component; this detailed information will allow us to identify early diagnostic biomarkers of progression and biomarkers for prognosis and to alter serum lipid profiles using individual or public nutritional strategies.

METHODS

Profiles of FFAs

Based on the previous study by Bermudez and Velasquez [10], the serum FFA profile of three groups of adolescents between the ages of 10 and 18 years was evaluated; these were (1) obese adolescents with MetS (OWMS), (2) obese adolescents without MetS (ONMS), and (3) normal weight (NW) adolescents. In the initial phase, the profile of each group of adolescents was evaluated using the eight fatty acids found with the highest frequencies in the three groups (Table 1). Each profile was reconstructed with the same quantity and proportion of each fatty acid found in each group. The second phase of the experiments was performed to determine whether the experimentally observed effects were due to the quantity or the ratio of FFA in each profile; the total quantity of FFAs in the mixture corresponding to the OWMS profile was decreased to be the same level as the total quantity corresponding to the NW profile without any change in the proportion of FFAs (Table 2). With these new profiles, that were equalized with respect to the total amount of FFAs but had the same proportions of FFAs, functional and viability tests in MIN6 (mouse insulinoma) cells were performed.

Table 1

Principal fatty acids in the FFAs of the adolescents in each study group

Data obtained from Bermudez et al. [10].

FFA, free fatty acid; NW, normal weight; ONMS, obese adolescents without metabolic syndrome (MetS); OWMS, obese adolescents with MetS; NS, not significant.

^a $P < 0.05$, ^bKruskal-Wallis, ^cNW vs. OWMS, ^dONMS vs. OWMS, ^eAnalysis of variance followed by Tukey *post hoc* test for multiple comparisons, ^fNW vs. ONMS.

Fatty acid	NW		ONMS		OWMS		P value ^a
	Concentration, mg/dL	No. (%)	Concentration, mg/dL	No. (%)	Concentration, mg/dL	No. (%)	
Total FFA	103.25	31 (100)	106.16	32 (100)	121.90	32 (100)	b,c,d
C16:0 (palmitic)	30.01	31 (28.71)	31.85	32 (30.00)	36.75	32 (30.15)	b,c,d

Fatty acid	NW		ONMS		OWMS		P value ^a
	Concentration, mg/dL	No. (%)	Concentration, mg/dL	No. (%)	Concentration, mg/dL	No. (%)	
C16:1 (palmitoleic)	1.74	8 (1.72)	1.84	19 (1.74)	2.28	27 (1.87)	NS
C18:0 (stearic)	15.60	31 (15.16)	15.30	32 (14.41)	17.30	32 (14.20)	b,c
C18:1 <i>n</i> -9c (oleic)	20.14	31 (19.01)	19.64	32 (18.50)	23.95	32 (19.65)	NS
C18:2 <i>n</i> -6c (linoleic, LA)	21.80	31 (21.49)	22.35	32 (21.05)	25.07	32 (20.57)	c,e
C20:3 <i>n</i> -6 (cis-8,11,14-dihomo- γ -linolenic, DGLA)	3.41	30 (3.36)	4.13	31 (3.89)	5.19	31 (4.26)	b,c,d,f
C20:4 <i>n</i> (arachidonic, AA)	6.97	31 (7.01)	7.48	31 (7.05)	8.13	32 (6.67)	b,c
C22:6 <i>n</i> -3 (cis-4,7,10,13,16,19-docosahexaenoic, DHA)	3.59	25 (3.54)	3.56	26 (3.36)	3.21	31 (2.63)	NS
(LA+DGLA+AA)/DHA	8.96	25	9.00	26	11.96	31	NS

[Open in a separate window](#)

 Download Table

Table 2

Concentration of FFA in OWMS adolescents, adjusted to the concentration found for NW adolescents

FFA, free fatty acid; OWMS, obese adolescents with metabolic syndrome (MetS); NW, normal weight.

Fatty acid	NW		OWMS	
	Concentration, mg/dL	Percentage	Concentration, mg/dL	Percentage
Total FFA	103.25	100.00	103.25	100.00
C16:0 (palmitic)	30.01	28.71	31.13	30.15
C16:1 (palmitoleic)	1.74	1.72	1.93	1.87
C18:0 (stearic)	15.60	15.16	14.66	14.20
C18:1 <i>n</i> -9c (oleic)	20.14	19.01	20.29	19.65
C18:2 <i>n</i> -6c (linoleic, LA)	21.80	21.49	21.24	20.57
C20:3 <i>n</i> -6 (cis-8,11,14-dihomo- γ -linolenic, DGLA)	3.41	3.36	4.40	4.26
C20:4 <i>n</i> (arachidonic, AA)	6.97	7.01	6.89	6.67
C22:6 <i>n</i> -3 (cis-4,7,10,13,16,19-docosahexaenoic, DHA)	3.59	3.54	2.72	2.63
(LA+DGLA+AA)/DHA	8.96	-	11.96	-

Preparation of the FFA profiles

The FFAs and bovine serum albumin (BSA) were obtained from Sigma (St. Louis, MO, USA). Saturated and monounsaturated fatty acids (MUFAs) were solubilized in distilled water. The polyunsaturated fatty acids (PUFAs) were prepared in methanol under nitrogen atmosphere. The mixtures of eight different fatty acids in the quantity and proportion identified in the sera of the three groups were complexed to BSA with shaking for 3 hours at 37°C. A fixed FFA:BSA ratio of 6:1 was maintained. This mixture was dissolved in Dulbecco's modified Eagle medium (DMEM), with a high glucose level and no fetal bovine serum (FBS), and was added to the MIN6 cells for the various evaluations.

Cellular culture

MIN6 cells were obtained from AddexBio Technologies (San Diego, CA, USA). The cells were maintained in DMEM, containing 25 mmol/L glucose, 2 mmol/L glutamine, 500 μM pyruvate, and 15% FBS at 37°C, with 5% CO₂ and 95% humidity. The cells were seeded for the experiments in 48-well plates at 100,000 cells/well in medium without FBS for 24 hours. They were then washed in phosphate-buffered saline (PBS), and the FFA:BSA mixture with the appropriate profile was added to each group, whereas the control group received medium that was free of BSA and FFA. The cells were incubated for 24 or 48 hours, and the medium was changed after 24 hours for the 48-hour experiments. For the experiments on the adenosine triphosphate/adenosine diphosphate (ATP/ADP) relationship, the cells were cultured in 96-well plates at a density of 10,000 cells/well under the conditions described above.

Glucose-stimulated insulin secretion

GSIS was evaluated after incubating the cells with FFA profiles or without FFA (control). The cells were washed in PBS with 2 mmol/L glucose and maintained for 2 hours prior to collecting the supernatant (baseline secretion). Immediately thereafter, DMEM with 25 mmol/L glucose without FBS was added, the cells were incubated for 30 minutes, and the supernatant was collected (GSIS). The insulin secreted in the baseline condition and under glucose stimulation was quantified using an ultrasensitive enzyme immunoassay (Mercodia, Uppsala, Sweden). The results are expressed as a percentage of the change in GSIS [(stimulated concentration-baseline concentration)/baseline concentration]×100.

Cellular viability assay

The effect of the different FFA profiles on cellular viability was evaluated using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) colorimetric assay [11]. The absorbance of each well was measured at 490 nm in a spectrophotometer (Varioskan Lux; Thermo Fisher, Waltham, MA, USA) and cell viability (percentage) was determined by normalizing each group to the control.

Evaluation of mitochondrial function

To evaluate the mitochondrial function in the cells treated with the FFA mixtures corresponding to the profiles of each group, GSIS was induced, and the cells were then immediately washed in PBS. After treatment with trypsin, 5 μ L of the MitoTracker fluorophore (50 nM) and propidium iodide (250 nM) were added, both in PBS. Using flow cytometry (BD FACSCanto II Flow Cytometer; BD Biosciences, San Jose, CA, USA), morphological patterns were determined, and the mean intensity of MitoTracker fluorescence was quantified. In addition, the ATP/ADP ratio was quantified using a commercial kit (Abcam, Cambridge, UK).

Lipid peroxidation and antioxidant capacity

The content of intracellular lipid peroxidation products was determined using the thiobarbituric acid reactive substances (TBARS) assay [12]. Briefly, 1 mL cell extract was combined with 1 mL 0.02 M thiobarbituric acid and heated at 90°C for 45 minutes; the mixture was then cooled to room temperature, and the absorbance was determined at 532 nm (Varioskan Lux). The results are expressed as malondialdehyde (MDA) equivalents/mg of total protein. The antioxidant capacity of the cell lysates after incubation with the FFAs was evaluated using the peroxynitrite anion (ONOO⁻) trapping capacity, following the protocol described by Whiteman et al. [13]. The capacity of the lysates to inhibit the peroxynitrite oxidation of the reagent was quantified relatively to ascorbic acid (AA) and normalized to the quantity of total protein. The results are shown in milligram of AA/mg of total protein.

Statistical analysis


The difference between groups was evaluated using the analysis of variance with *post hoc* Tukey test for multiple comparisons. The FFA concentrations were expressed as the mean \pm standard error or the median \pm interquartile range. Differences were considered statistically significant at $P < 0.05$. The statistical analysis was performed using the program GraphPad Prism version 7.00 for Mac OS X (GraphPad Software, La Jolla, CA, USA).

Go to :

RESULTS

The serum FFA mixture of OWMS adolescents reduces the secretory function and the viability of MIN6 cells

To determine the overall effect of the FFAs of each profile (NW, ONMS, and OWMS) on the function of β -cells, GSIS was quantified in MIN6 cells treated for 24 or 48 hours. After stimulation with the culture medium containing 25 mmol/L glucose without FBS, GSIS in the cells treated with the NW and ONMS FFA mixtures did not significantly differ from that of the control cells. However, GSIS at 24 hours was significantly lower in the cells treated with the OWMS FFA mixture than in the cells treated with the FFA mixtures corresponding to the NW and ONMS profiles. The reduction in GSIS was maintained after treatment for 48 hours with the FFA mixture corresponding to the OWMS profile ([Fig. 1A and B](#)).










[Open in a separate window](#)

To determine whether the FFA mixtures impact the viability of MIN6 cells, MTT assays were carried out for 24 and 48 hours (Fig. 1C and D). In the cells treated with the FFA mixture corresponding to the OWMS profile, cell viability decreased after 24 and 48 hours of treatment to 40% compared to that of the cells treated with the FFA mixture corresponding to the NW profile. The other groups did not show significant changes.

The FFA mixture corresponding to the OWMS profile reduces the mitochondrial function of MIN6 cells

Mitochondrial function was evaluated by quantifying the potential of the internal mitochondrial membrane (IMM) using the MitoTracker fluorophore. Through flow cytometry, the mean intensity of fluorescence (MIF) was quantified in each cell group treated with the FFA mixtures and in the untreated control ([Fig. 2](#)). The IMM potential remained uniform in the control and in the NW and ONMS groups after both 24 and 48 hours of incubation, but the group of cells treated with the FFA mixture corresponding to the OWMS profile had a significantly lower potential at 24 and 48 hours (50%) compared to that of the control cells.



 [Open in a separate window](#)

The ATP/ADP results demonstrated that the cells treated with the FFA mixture corresponding to the OWMS profile also reduced the ATP/ADP ratio compared to that of the control after both 24 and 48 hours of coincubation, with a cumulative effect over time ([Fig. 2C and D](#)).


The FFA mixture corresponding to the OWMS profile increases lipid peroxidation and reduces the antioxidant capacity

The results of a TBARS assay showed that treatment with the FFA mixture corresponding to the OWMS profile increased the concentration of MDA, which is the principal product of lipid peroxidation, at both 24 and 48 hours of coincubation. Although the cells treated with the FFA mixtures corresponding to the NW and ONMS profiles tended to present increased lipid peroxidation at 24 hours, this increase was markedly lower than observed in the OWMS group, and it remained the same in cells treated for 48 hours ([Fig. 2E and F](#)). β -Cells are particularly vulnerable to oxidative stress given their reduced antioxidant defence. Treatment with the FFA mixture corresponding to the OWMS profile reduced the capacity to capture peroxynitrite compared to that of the control or for the treatment with the FFA mixture corresponding to the NW profile; in the latter two groups, the antioxidant capacity was maintained without changes after both 24 and 48 hours of treatment ([Fig. 2G and H](#)).

Effect of the proportion of FFAs on the function and viability of MIN6 cells

To determine whether the negative effects of the OWMS mixture of FFAs on the function and viability of the MIN6 cells observed in this study were due to the total quantity of FFAs or to their proportion in the total profile, the total quantity of FFAs in the OWMS mixture was equalized so as to be the same as that in the NW mixture while maintaining the relative proportion of each fatty acid ([Table 2](#)). The effect of the ONMS FFA mixture was not further studied because in all tests, it showed a pattern very similar to that of the NW profile. At 24 hours, the treatment with the NW FFA mixture significantly increased GSIS compared to that of the control ([Fig. 3A and B](#)). In contrast, treatment with the OWMS FFA mixture decreased insulin secretion by approximately 50%. At 48 hours, the cells treated with the NW FFA mixture maintained GSIS without changes compared to 24 hours, whereas the cells treated with the OWMS FFA mixture recovered GSIS, with an 86% increase in secretion compared to baseline secretion ([Fig 3A and B](#)). Interestingly, cell viability determined by the MTT assay did not show changes in the cells treated with

the OWMS FFA mixture at 24 hours but did show a reduction of approximately 20% after 48 hours of treatment ([Fig. 3C and D](#)).











[Open in a separate window](#)

The IMM potential remained uniform in the control and NW groups at both 24 and 48 hours of incubation, but the treatment with the OWMS FFA mixture caused a significant reduction in the potential at 24 and 48 hours (40%) compared to that of the control group ([Fig. 4A and B](#)). Quantifying the

ATP/ADP ratio indicated no significant change between the groups, although the ratio tended to be lower in the OWMS group ([Fig. 4C and D](#)).

24 Hours



 [Open in a separate window](#)

Compared to control cells and to cells treated with the NW FFA mixture, the cells treated with the OWMS FFA mixture presented a significantly higher MDA concentration after 48 hours of coincubation. However, cells treated with the NW FFA mixture showed a significant increase in the production of MDA compared to that produced in control cells ([Fig. 4E and F](#)). A similar effect was observed when evaluating the antioxidant capacity. Compared with the control cells and cells treated with the NW FFA mixture, cells treated with the OWMS FFA mixture had a lower capacity to capture peroxynitrite after 48 hours of treatment. The antioxidant capacity of cells treated with the NW FFA mixture was also lower than that of the control cells ([Fig. 4G and H](#)). The significant decrease in the antioxidant capacity and the increase in the reactive oxygen species (ROS) observed ([Fig. 4F and H](#)) in the cells treated with the NW FFA mixture compared to control cells may be explained by the high passage number in which the experiments were performed. In phase 2, cells with a much higher passage number were used than those used in phase 1, and the cells became sensitive to oxidative stress due to a decrease in antioxidant capacity.

Go to : 

DISCUSSION

IR in peripheral tissues and the dysfunction of pancreatic β -cells are the principal pathological characteristics of T2DM and the latter abnormality results from prolonged exposure to high concentrations of glucose and FFAs; that is, frequently associated with obesity and MetS.

The increase in circulating FFA levels, particularly in saturated FFAs, is the principal factor inducing apoptosis and poor function of pancreatic β -cells. However, the effects vary with the length of the carbon chain and degree of saturation of circulating FFAs, via different mechanisms. There is significant controversy regarding the effect of fatty acids on β -cells [[7,8,9](#)]. For example, the presence of unsaturated FFAs rescues cells from the toxic effects of saturated FFAs [[14,15](#)]. There are no published studies that evaluate the effect

of the FFA mixture in the serum of OWMS or ONMS on the function and viability of pancreatic β -cells.

The FFA composition of serum can predict the risk of diabetes and cardiovascular disease, and it is closely related to components of MetS. Diverse studies have reported higher concentrations of myristic and stearic saturated fatty acids, palmitoleic (16:1 *n*-7) MUFA and dihomo- γ -linolenic (20:3 *n*-6) PUFA, and lower concentrations of linoleic acid (18:2 *n*-6) and docosahexaenoic acid (DHA; 22:6 *n*-3) in obese children than in non-obese children. In obese children with MetS, the concentrations of oleic (18:1 *n*-9) and palmitoleic (16:1 *n*-7) MUFAs and γ linoleic (18:3 *n*-6) PUFAs were higher, and the concentrations of linoleic (18:2 *n*-6) PUFAs also remained low [16,17]. The cellular defects induced by high levels of circulating FFAs and glucose (glucolipototoxicity), that may be responsible for cell death or deterioration in the secretory function of β -cells, include epigenetic changes [18], inflammation [19], changes in cellular signalling [20], oxidative stress [21], endoplasmic reticulum (ER) stress [22], and mitochondrial dysfunction [23]. The main aim of present study was to evaluate the effect the total fatty acid mixture had on adolescent's plasma on the function of MIN6 cells. We considered that a more real approach was to analyze the global effect of this FFA mixture.

In OWMS adolescents, the total amount of FFAs is higher than that of ONMS or NW adolescents (Table 1) and probably reflects the inflammatory status, IR and adrenergic hypersensitivity manifested by hypertrophied adipocytes. In these circumstances, there is substantial mobilisation of the FFAs that reach and perfuse the pancreas; these FFAs then directly contact β -cells and exert a combination of effects on the secretory dynamics of these cells. In addition to the total quantity of FFAs, the population of polyunsaturated ω 6 FFAs in the serum of OWMS adolescents significantly differed from that of the NW and ONMS groups. The increase in metabolic intermediaries in the pathway of ω 6 PUFAs is accompanied by an increase in the production of prostaglandins and series two thromboxanes, which stimulate inflammatory processes that can be associated with the pathogenesis of T2DM [24]. For example, prostaglandin E₂ (PGE₂) is a negative regulator of insulin secretion in HIT-T15 cell lines but does not affect cell viability [25].

The current study demonstrates that 24 and 48 hours of treatment with the FFA mixture corresponding to the profile of OWMS adolescents, compared with that of the NW group, decreased the secretory function and viability of MIN6 cells. The precise mechanisms through which FFAs generate functional damage in β -cells have yet to be clarified. Previous studies reported that long-chain and saturated acyl-CoA accumulates in the cytoplasm and potentiates synthesis of ceramides that are implicated in apoptosis and functional damage [26]. Other mechanisms include over-activation of protein kinase C or inhibition of Akt kinase and calpain-10 [27,28]. In addition, activation of the stress kinases PKR (double-stranded RNA-dependent protein kinase), IKK ($\text{I}\kappa\text{B}$ kinase), and JNK (c-Jun N-terminal kinase) has been documented. The

activation of these kinases induces intracellular proinflammatory pathways. For example, in cells treated with saturated FFAs, the expression of nuclear factor- κ B increases up to 10-fold [29].

There was a limitation in this research. In the GSIS assay, the cells were incubated in PBS with 2 mmol/L glucose to reduce basal insulin secretion levels. Afterwards, insulin secretion was stimulated in DMEM with 25 mmol/L glucose. PBS and DMEM are different buffers, and cells treated with PBS with 2 mmol/L glucose may not be the best control for cells treated with DMEM with 25 mmol/L glucose. However this does not affect the percentage of insulin secretion (up to 150%; data not shown).

It is also possible that the FFAs corresponding to the OWMS profile induce acute changes in the fluidity of microsomal membranes, impairing the function of the ER and thus reducing the production and processing of insulin. Saturated FFAs have been implicated as participants in the activation of the unfolded protein response (UPR) that can culminate in apoptosis. The onset of apoptosis has been documented in INS-1 (rat insulinoma) cells, with only 16 hours of incubation with 1 mmol/L palmitate [8].

As mentioned above, either the quantity of fatty acids or the proportions of the different FFAs present in this profile may explain the effect on GSIS and cell viability observed in cells treated with the OWMS FFA mixture. For this reason, experiments were carried out in which the total amount of FFAs from the mixture corresponding to the OWMS profile was reduced to the same levels as the NW profile without changes in the proportions of the FFAs (Table 2). Insulin secretion was only significantly diminished after 24 hours of treatment with the FFA mixture corresponding to the modified OWMS profile, but after 48 hours, the levels returned to those of cells treated with the NW profile. At 48 hours, the cells probably adjusted to the new nutritional conditions and regained their capacity to secrete insulin. Conversely, the viability of the cells was reduced at 48 hours of treatment with the FFA mixture corresponding to the modified OWMS profile, possibly indicating that in addition to the amount of FFA present in the medium, the proportion of ω 6 fatty acids may exert a negative effect on cell viability.

Mitochondrial function was evaluated, and the FFA mixture corresponding to the OWMS profile reduced the membrane potential and the ATP/ADP ratio in MIN6 cells. Saturated FFAs induce negative changes in the mitochondrial dynamics, such as fission, liberation of cytochrome C, an excessive load of calcium in the matrix or an increase in transitional mitochondrial permeability with an acute decrease in membrane potential [15]. ATP is the principal inducer in the depolarisation of β -cells; thus, its decrease has rapid repercussions on secretion during the first stage. Disorders in mitochondria tend to lead to an increase in reactive oxygen and nitrogen species. Due to their instability and rapid disappearance, we quantified the production of lipid peroxidation products as a direct indicator of cellular oxidative stress. The OWMS FFA mixture may induce structural mitochondrial damage that culminates in oxidative stress. The cells were cultivated in conditions of high

glucose (25 mmol/L) to simulate both glucotoxic conditions and other substrates for oxidation; the fatty acids themselves impose a great oxidative demand that can facilitate the generation of ROS and reactive nitrogen species if the damage is maintained in the mitochondrial membranes [23]. The antioxidant defence of β -cells tends to be reduced, making them especially vulnerable to damage by ROS and RNS. The cells treated with the OWMS FFA mixture had a significantly lower capacity to trap peroxynitrite. This change may be due to the acute load on non-enzymatic antioxidant mechanisms and the reduction in the synthesis of antioxidant enzymes in the context of ER stress and the activation of the UPR [22].

When the total amount of FFAs corresponding to the OWMS profile was decreased, a significant reduction was observed in the MIF, at both 24 and 48 hours of treatment; this effect was not correlated with the ATP/ADP ratio but with an increase in ROS and a decrease in the antioxidant capacity at 48 hours of treatment.


Oxidative stress has been closely related to the dysfunction and death of β -cells in the pancreas, and this can partly explain the physiopathology of T2DM [30,31]. In fact, the control of ROS improves the health of β -cells, even in conditions of glucolipotoxicity [32,33]. The results obtained in this study could suggest that an excessive nutritional load or a greater quantity of ω 6 fatty acids could be involved in deterioration in the electronic transport chain, leading to a decrease in the mitochondrial membrane potential. If the electronic transport chain does not work properly, the electrons generated by the oxidation of nutrients can increase the level of ROS.

The mechanisms by which ω 6 fatty acids induce cell damage are not well known. Meng et al. [25] have demonstrated that PGE₂, a metabolic product of arachidonic acid, has a negative effect on insulin secretion. A probable mechanism is to reduce AKT protein activation and stimulate FoxO (Forkhead box O protein) protein activation. This will reduce these cells' proliferation and survival. On the other hand, the results of the same group, confirmed by Wei et al. [34], demonstrated that PGE₂ stimulates the expression of a receptor-associated Gi protein, reduces cyclic adenosine monophosphate levels, and causes a concomitant reduction of insulin secretion [25].

Previous studies conducted on rodents and large-scale population studies have demonstrated that ω 6 eicosanoids are more powerful mediators of inflammation than ω 3. An increase in the ω 6/ ω 3 ratio greatly favors proinflammatory processes, and this contributes significantly to establishing IR and diabetes-related complications. These facts suggest that inflammation control, by reducing ω 6 can have a beneficial effect on a beta cell's secretory function [35].

Several studies have demonstrated that ω 3-generated lipid mediators, including eicosapentaenoic acid (20:5 *n*-3) and DHA (22:6 *n*-3) have anti-inflammatory and protective activities in different pathologies including T2DM [36]. ω 3 Production in transgenic mice that overexpress the fat-1 gene

(an ω_3 desaturase, that catalyzes the ω_6 to ω_3 conversion) favors GSIS and protects β -cells from cytokine-mediated destruction [34]. Studies conducted on streptozotocin-induced diabetic rats demonstrated that a ω_3 -enriched diet reduces hyperglycemia [37]. On the other hand, ω_3 reduces IR and obesity-induced hepatic steatosis through the stimulation of insulin-sensitivity-related gene expression such as type 2 and type 4 glucose transporters, and type 1 and type 2 insulin receptor substrates [38]. In conclusion, modern diets have an elevated content of ω_6 PUFAs and very low levels of ω_3 , leading to very unhealthy ω_6/ω_3 ratios in the blood, with a level of 20:1 instead of the ideal proportion of 1:1 [39]. Adolescents with obesity and MetS have a very elevated risk of developing T2DM. These results may contribute in part to an improvement in the nutritional management of OWMS adolescents by suggesting an increase in the consumption of foods rich in ω_3 fatty acids and thereby reducing the probability of developing T2DM. In a future study, we shall examine the individual effect of each one of the FFAs in the proportion found in the plasma of each experimentation group.

Go to : 

ACKNOWLEDGMENTS

We thank Anne-Lise Haenni for her critical review. This study was supported by COLCIENCIAS (Colombia) grant No. 1115-569-33200.

Go to : 

NOTES

CONFLICTS OF INTEREST: No potential conflict of interest relevant to this article was reported.

Go to : 

REFERENCES

1. NCD Risk Factor Collaboration (NCD-RisC). Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 192 million participants. *Lancet* 2016;387:1377-1396.

[PubMed](#) 

2. Must A, Spadano J, Coakley EH, Field AE, Colditz G, Dietz WH. The disease burden associated with overweight and obesity. *JAMA* 1999;282:1523-1529.

[PubMed](#) 

3. Mokdad AH, Ford ES, Bowman BA, Dietz WH, Vinicor F, Bales VS, Marks JS. Prevalence of obesity, diabetes, and obesity-related health risk factors,

2001. JAMA 2003;289:76-79.

[PubMed](#) 

4. Despres JP, Lemieux I. Abdominal obesity and metabolic syndrome. Nature 2006;444:881-887.

[PubMed](#) 

5. Hotamisligil GS. Inflammation and metabolic disorders. Nature 2006;444:860-867.

[PubMed](#) 

6. Kloting N, Bluher M. Adipocyte dysfunction, inflammation and metabolic syndrome. Rev Endocr Metab Disord 2014;15:277-287.

[PubMed](#) 

7. Furstova V, Kopska T, James RF, Kovar J. Comparison of the effect of individual saturated and unsaturated fatty acids on cell growth and death induction in the human pancreatic beta-cell line NES2Y. Life Sci 2008;82:684-691.

[PubMed](#) 

8. Karaskov E, Scott C, Zhang L, Teodoro T, Ravazzola M, Volchuk A. Chronic palmitate but not oleate exposure induces endoplasmic reticulum stress, which may contribute to INS-1 pancreatic beta-cell apoptosis. Endocrinology 2006;147:3398-3407.

[PubMed](#) 

9. Lai E, Bikopoulos G, Wheeler MB, Rozakis-Adcock M, Volchuk A. Differential activation of ER stress and apoptosis in response to chronically elevated free fatty acids in pancreatic beta-cells. Am J Physiol Endocrinol Metab 2008;294:E540-E550.

[PubMed](#) 

10. Bermudez JA, Velasquez CM. Profile of free fatty acids (FFA) in serum of young Colombians with obesity and metabolic syndrome. Arch Latinoam Nutr 2014;64:248-257.

[PubMed](#) 

11. Berridge MV, Tan AS. Characterization of the cellular reduction of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT): subcellular localization, substrate dependence, and involvement of mitochondrial electron transport in MTT reduction. Arch Biochem Biophys 1993;303:474-482.

[PubMed](#) 

12. Rojano BA, Gaviria CA, Saez JA. Antioxidant activity determination in a lipidic peroxidation model of butter inhibited by isoespintanol. Vitae 2008;15:212-218.

13. Whiteman M, Ketsawatsakul U, Halliwell B. A reassessment of the peroxy-nitrite scavenging activity of uric acid. Ann N Y Acad Sci 2002;962:242-

259.

[PubMed](#) 

14. Maedler K, Spinas GA, Dyntar D, Moritz W, Kaiser N, Donath MY. Distinct effects of saturated and monounsaturated fatty acids on beta-cell turnover and function. *Diabetes* 2001;50:69-76.

[PubMed](#) 

15. Maedler K, Oberholzer J, Bucher P, Spinas GA, Donath MY. Monounsaturated fatty acids prevent the deleterious effects of palmitate and high glucose on human pancreatic beta-cell turnover and function. *Diabetes* 2003;52:726-733.

[PubMed](#) 

16. Kawashima A, Sugawara S, Okita M, Akahane T, Fukui K, Hashiuchi M, Kataoka C, Tsukamoto I. Plasma fatty acid composition, estimated desaturase activities, and intakes of energy and nutrient in Japanese men with abdominal obesity or metabolic syndrome. *J Nutr Sci Vitaminol (Tokyo)* 2009;55:400-406.

[PubMed](#) 

17. Klein-Platat C, Draï J, Oujaa M, Schlienger JL, Simon C. Plasma fatty acid composition is associated with the metabolic syndrome and low-grade inflammation in overweight adolescents. *Am J Clin Nutr* 2005;82:1178-1184.

[PubMed](#) 

18. Malmgren S, Spegel P, Danielsson AP, Nagorny CL, Andersson LE, Nitert MD, Ridderstrale M, Mulder H, Ling C. Coordinate changes in histone modifications, mRNA levels, and metabolite profiles in clonal INS-1 832/13 beta-cells accompany functional adaptations to lipotoxicity. *J Biol Chem* 2013;288:11973-11987.

[PubMed](#)  [PMC](#) 

19. Akash MS, Rehman K, Chen S. Role of inflammatory mechanisms in pathogenesis of type 2 diabetes mellitus. *J Cell Biochem* 2013;114:525-531.

[PubMed](#) 

20. Haber EP, Hirabara SM, Gomes AD, Curi R, Carpinelli AR, Carvalho CR. Palmitate modulates the early steps of insulin signalling pathway in pancreatic islets. *FEBS Lett* 2003;544:185-188.












[PubMed](#) 

21. Pi J, Collins S. Reactive oxygen species and uncoupling protein 2 in pancreatic beta-cell function. *Diabetes Obes Metab* 2010;12(Suppl 2):141-148.

[PubMed](#) 

22. Cnop M, Foufelle F, Velloso LA. Endoplasmic reticulum stress, obesity and diabetes. *Trends Mol Med* 2012;18:59-68.

[PubMed](#) 

23. Lowell BB, Shulman GI. Mitochondrial dysfunction and type 2 diabetes. *Science* 2005;307:384-387.
[PubMed](#) 
24. Luo P, Wang MH. Eicosanoids, beta-cell function, and diabetes. *Prostaglandins Other Lipid Mediat* 2011;95:1-10.
[PubMed](#)  [PMC](#) 
25. Meng ZX, Sun JX, Ling JJ, Lv JH, Zhu DY, Chen Q, Sun YJ, Han X. Prostaglandin E2 regulates Foxo activity via the Akt pathway: implications for pancreatic islet beta cell dysfunction. *Diabetologia* 2006;49:2959-2968.
[PubMed](#) 
26. Lupi R, Dotta F, Marselli L, Del Guerra S, Masini M, Santangelo C, Patane G, Boggi U, Piro S, Anello M, Bergamini E, Mosca F, Di Mario U, Del Prato S, Marchetti P. Prolonged exposure to free fatty acids has cytostatic and pro-apoptotic effects on human pancreatic islets: evidence that beta-cell death is caspase mediated, partially dependent on ceramide pathway, and Bcl-2 regulated. *Diabetes* 2002;51:1437-1442.
[PubMed](#) 
27. Eitel K, Staiger H, Rieger J, Mischak H, Brandhorst H, Brendel MD, Bretzel RG, Haring HU, Kellerer M. Protein kinase C delta activation and translocation to the nucleus are required for fatty acid-induced apoptosis of insulin-secreting cells. *Diabetes* 2003;52:991-997.
[PubMed](#) 
28. Wrede CE, Dickson LM, Lingohr MK, Briaud I, Rhodes CJ. Protein kinase B/Akt prevents fatty acid-induced apoptosis in pancreatic beta-cells (INS-1). *J Biol Chem* 2002;277:49676-49684.
[PubMed](#) 
29. Rakatzi I, Mueller H, Ritzeler O, Tennagels N, Eckel J. Adiponectin counteracts cytokine- and fatty acid-induced apoptosis in the pancreatic beta-cell line INS-1. *Diabetologia* 2004;47:249-258.
[PubMed](#) 
30. Mehmeti I, Gurgul-Convey E, Lenzen S, Lortz S. Induction of the intrinsic apoptosis pathway in insulin-secreting cells is dependent on oxidative damage of mitochondria but independent of caspase-12 activation. *Biochim Biophys Acta* 2011;1813:1827-1835.
[PubMed](#) 
31. Newsholme P, Haber EP, Hirabara SM, Rebelato EL, Procopio J, Morgan D, Oliveira-Emilio HC, Carpinelli AR, Curi R. Diabetes associated cell stress and dysfunction: role of mitochondrial and non-mitochondrial ROS production and activity. *J Physiol* 2007;583(Pt 1):9-24.
[PubMed](#)  [PMC](#) 

32. Lim S, Rashid MA, Jang M, Kim Y, Won H, Lee J, Woo JT, Kim YS, Murphy MP, Ali L, Ha J, Kim SS. Mitochondria-targeted antioxidants protect pancreatic beta-cells against oxidative stress and improve insulin secretion in glucotoxicity and glucolipotoxicity. *Cell Physiol Biochem* 2011;28:873-886.

[PubMed](#) [↗](#)

33. Lin N, Chen H, Zhang H, Wan X, Su Q. Mitochondrial reactive oxygen species (ROS) inhibition ameliorates palmitate-induced INS-1 beta cell death. *Endocrine* 2012;42:107-117.

[PubMed](#) [↗](#)

34. Wei D, Li J, Shen M, Jia W, Chen N, Chen T, Su D, Tian H, Zheng S, Dai Y, Zhao A. Cellular production of n-3 PUFAs and reduction of n-6-to-n-3 ratios in the pancreatic beta-cells and islets enhance insulin secretion and confer protection against cytokine-induced cell death. *Diabetes* 2010;59:471-478.

[PubMed](#) [↗](#)

35. Simopoulos AP. Importance of the ratio of omega-6/omega-3 essential fatty acids: evolutionary aspects. *World Rev Nutr Diet* 2003;92:1-22.

[PubMed](#) [↗](#)

36. Calder PC. n-3 Polyunsaturated fatty acids, inflammation, and inflammatory diseases. *Am J Clin Nutr* 2006;83(6 Suppl):1505S-1519S.

[PubMed](#) [↗](#)

37. Linn T, Noke M, Woehrl M, Kloer HU, Hammes HP, Litzlbauer D, Bretzel RG, Federlin K. Fish oil-enriched diet and reduction of low-dose streptozocin-induced hyperglycemia. Inhibition of macrophage activation. *Diabetes* 1989;38:1402-1411.

[PubMed](#) [↗](#)

38. Gonzalez-Periz A, Horrillo R, Ferre N, Gronert K, Dong B, Moran-Salvador E, Titos E, Martinez-Clemente M, Lopez-Parra M, Arroyo V, Claria J. Obesity-induced insulin resistance and hepatic steatosis are alleviated by omega-3 fatty acids: a role for resolvins and protectins. *FASEB J* 2009;23:1946-1957.

[Article](#) [↗](#) [PubMed](#) [↗](#) [PMC](#) [↗](#)

39. Kang JX. The importance of omega-6/omega-3 fatty acid ratio in cell function. The gene transfer of omega-3 fatty acid desaturase. *World Rev Nutr Diet* 2003;92:23-36.

[Article](#) [↗](#) [PubMed](#) [↗](#)