

Tabla 3

| Tabla 3. Declive neuropsicológico a través de diferentes Pruebas mentales | | | | | | | | | | | | |
|--|-----------|-------------|------------------|-----------|-------------|--------------|-----------|-------------|--------------|-----------|-------------|--------------|
| Variables | BASC | | | HVLTR | | | MAZES | | | FLUENCY | | |
| | Estimates | IC | p | Estimates | IC | p | Estimates | IC | p | Estimates | IC | p |
| Edad | -0.1 | -0.3 – 0.1 | 0.305 | 0.0 | -0.1 – 0.1 | 0.885 | 0.0 | -0.1 – 0.2 | 0.556 | 0.0 | -0.2 – 0.3 | 0.715 |
| Sexo Masculino | -1.6 | -6.8 – 3.5 | 0.538 | -1.4 | -5.7 – 3.0 | 0.542 | -2.9 | -7.8 – 2.1 | 0.256 | -1.6 | -7.0 – 3.8 | 0.564 |
| Escolaridad > 11 años | 1.7 | -1.9 – 5.4 | 0.359 | 3.8 | 0.1 – 7.6 | 0.047 | 2.1 | -1.8 – 6.0 | 0.284 | 1.5 | -2.2 – 5.1 | 0.431 |
| Inicio de enfermedad | -0.0 | -0.3 – 0.3 | 0.928 | 0.1 | -0.2 – 0.3 | 0.634 | 0.1 | -0.1 – 0.3 | 0.300 | 0.1 | -0.3 – 0.5 | 0.609 |
| SANS | -0.1 | -0.2 – -0.1 | <0.001 | -0.0 | -0.1 – 0.0 | 0.081 | -0.1 | -0.2 – -0.1 | 0.001 | -0.1 | -0.2 – -0.0 | 0.005 |
| Antecedente de consumo de SPA | 0.3 | -3.7 – 4.3 | 0.874 | 2.5 | -1.7 – 6.7 | 0.246 | 2.5 | -1.8 – 6.8 | 0.261 | 1.3 | -3.4 – 6.1 | 0.581 |
| Diferencia promedio en el puntaje de la prueba por año desde la primera evaluación | -0.4 | -0.8 – -0.0 | 0.042 | -0.8 | -1.4 – -0.1 | 0.025 | -0.6 | -1.0 – -0.2 | 0.001 | -0.7 | -1.1 – -0.2 | 0.006 |

Abreviaturas
 BASC, Brief Assessment of Cognition in Schizophrenia
 HVTR, revised Hopkins Verbal Learning Test
 MAZES, Neuropsychological Assessment Battery
 FLUENCY, Category Fluency
 IC, Intervalo de confianza

Tabla 4

| Tabla 4. Pruebas neuropsicologicas que no presentaron cambios significativos en el tiempo | | | | | | | | | | | | |
|---|-----------|-------------|--------------|-----------|-------------|------------------|-----------|--------------|------------------|-----------|-------------|--------------|
| Variables | TMT | | | WMS | | | LNS | | | CPT | | |
| | Estimates | IC | p | Estimates | IC | p | Estimates | IC | p | Estimates | IC | p |
| Edad | -0.2 | -0.4 – 0.0 | 0.097 | -0.1 | -0.3 – 0.1 | 0.422 | -0.2 | -0.4 – -0.0 | 0.047 | -0.3 | -0.5 – 0.0 | 0.089 |
| Sexo Masculino | -2.6 | -6.9 – 1.8 | 0.243 | -4.0 | -11.3 – 3.3 | 0.278 | -8.3 | -13.7 – -2.9 | 0.003 | 2.1 | -3.8 – 7.9 | 0.489 |
| Escolaridad > 11 años | 2.8 | -0.9 – 6.6 | 0.141 | 6.0 | 0.1 – 11.9 | 0.046 | 2.3 | -2.4 – 7.0 | 0.336 | 0.6 | -4.4 – 5.5 | 0.814 |
| Inicio de enfermedad | -0.1 | -0.4 – 0.2 | 0.491 | 0.2 | -0.2 – 0.7 | 0.323 | 0.2 | -0.1 – 0.6 | 0.127 | 0.3 | -0.0 – 0.7 | 0.063 |
| SANS | -0.1 | -0.2 – -0.0 | 0.017 | -0.1 | -0.2 – -0.1 | <0.001 | -0.2 | -0.2 – -0.1 | <0.001 | -0.1 | -0.2 – -0.0 | 0.021 |
| Antecedente de consumo de SPA | 1.7 | -2.0 – 5.4 | 0.380 | 4.2 | -0.9 – 9.3 | 0.103 | 3.2 | -1.3 – 7.8 | 0.162 | -1.2 | -8.0 – 5.6 | 0.727 |
| Diferencia promedio en el puntaje de la prueba por año desde la primera evaluación | 0.3 | -0.3 – 1.0 | 0.280 | -0.4 | -0.8 – 0.0 | 0.073 | -0.3 | -0.7 – 0.2 | 0.261 | -0.1 | -1.1 – 0.9 | 0.839 |

Abreviaturas:
TMT, Trail Making Test
WMS, Wechsler Memory Scale
LNS, University of Maryland Letter–Number Span test
CPT, Continuous Performance Test
IC, Intervalo de confianza