



ORIGINAL ARTICLE

Assessment of nasal obstruction: Correlation between subjective and objective techniques

J.J. Yepes-Nuñez^a, J. Bartra^{b,c,d}, R. Muñoz-Cano^{b,c,d}, J. Sánchez-López^b, C. Serrano^e, J. Mullol^{c,d,f}, I. Alobid^{c,d,f}, J. Sastre^{c,g}, C. Picado^{b,c,d}, A. Valero^{b,c,d,*}

^a Grupo de Alergología Clínica y Experimental (GACE), Grupo Académico de Epidemiología Clínica (GRAEPIC), Universidad de Antioquia, Medellín, Colombia

^b Unidad de Alergia, Servicio de Neumología y Alergia Respiratoria, Hospital Clínic, Barcelona, Spain

^c Centro de Investigación Biomédica en red para Enfermedades Respiratorias (CIBERES), Spain

^d Institut d'Investigacions Biomediques August Pi i Sunyer (IDIBAPS), Barcelona, Spain

^e Unidad de Alergia, Fundación Valle del Lili, Cali, Colombia

^f Unitat de Rinologia i Clínica de l'Olfacte, Servei d'Otorinolaringologia, Hospital Clínic, Barcelona, Spain

^g Servicio de Alergia, Fundación Jiménez Díaz, Madrid, Spain

Received 19 April 2012; accepted 17 May 2012

Available online 8 November 2012

KEYWORDS

Nasal obstruction;
Anterior
rhinomanometry;
Acoustic rhinometry;
Peak nasal;
Inspiratory flow;
Visual analogue scale

Abstract

Background: Nasal sensation of airflow describes the perception of the passage of air through the nose. Nasal obstruction can be assessed using subjective techniques (symptom scores and visual analogue scales [VAS]) and objective techniques (anterior rhinomanometry [RMN], acoustic rhinometry [AR], and peak nasal inspiratory flow [PNIF]). Few studies have evaluated the correlation between these techniques.

Objective: The primary objective of our study was to determine the degree of correlation between subjective and objective techniques to assess nasal obstruction.

Materials and methods: Nasal obstruction was assessed using a symptom score, VAS, RMN, AR (minimal cross-sectional area [MCSA] and volume), and PNIF in 184 volunteer physicians. Spearman's rho was recorded. Correlations were considered weak if $r \leq 0.4$, moderate if $0.4 < r < 0.8$, and strong if $r > 0.8$.

Results: Mean (SD) age was 37.1 (6.9) years (range, 25–56 years); 61% were women. We found a strong correlation ($r > 0.8$; $p = 0.001$) between the different parameters of RMN and a moderate correlation between symptom score and VAS ($r = 0.686$; $p = 0.001$) and between MCSA and RMN (resistance) ($r = 0.496$; $p = 0.001$) and PNIF ($r = 0.459$; $p = 0.001$). The correlations were weak or non-significant for the remaining comparisons.

* Corresponding author.

E-mail address: valero@clinic.ub.es (A. Valero).

Conclusion: Nasal obstruction can be assessed using subjective and objective approaches. The correlations between objective techniques were moderate to strong. In addition, between subjective techniques we reported a moderate correlation. Finally, the correlations between the subjective and objective techniques were weak and absent. These findings suggest that each of the techniques assesses different aspects of nasal obstruction, thus making them complementary.

© 2012 SEICAP. Published by Elsevier España, S.L. All rights reserved.

Introduction

The nasal airways humidify, filter, and warm the air we breathe. Consequently, they account for 50% of resistance in the respiratory tract.¹ Nasal obstruction is our perception of air passing through the nasal airways. It has been defined as discomfort caused by insufficient airflow in the nose or as the sensation of increased resistance of airflow through the nostrils. The sensation of airflow through the nose arises from the cooling of the sensory receptors in the nostrils during inspiration. These receptors are innervated by the trigeminal nerve, and anaesthesia or damage of this nerve can produce a sensation of nasal obstruction.²

Nasal obstruction is one of the main symptoms of the various conditions affecting the nose and may be due to anatomical abnormalities, inflammatory processes (eg. rhinitis), nasal polyps, and tumours.

Objective assessment of nasal obstruction involves direct examination of the area or volume of the nasal cavity or indirect analysis of nasal airflow and resistance.

Subjective assessment of nasal symptoms using symptom scores (including those examining nasal obstruction) are highly variable; therefore, the visual analogue scale [VAS] has been proposed as a tool to evaluate severity³. The most widely-used objective methods to assess nasal obstruction are active anterior rhinomanometry, acoustic rhinometry, and measurement of peak nasal inspiratory airflow (PNIF).³

A recent review evaluated recommendations on the use of these approaches and the correlations between them.^{4,5} Few studies have examined the correlation between subjective and objective methods for assessing nasal obstruction.^{4,6}

We analysed the correlation between subjective and objective techniques used to evaluate nasal obstruction.

Materials and methods

Design

We performed a prospective descriptive study based on the voluntary participation of 184 clinicians attending an educational course on rhinitis during the period 2002–2008. Nasal obstruction was subjectively assessed using a symptom score (0, absence of symptoms or mild symptoms; 1, moderate symptoms; and 2, severe symptoms) and a VAS scoring from 0 to 10 cm (0 indicated absence of symptoms and 10 indicated more severe symptoms). Nasal

obstruction was also objectively assessed using active anterior rhinomanometry to record airflow (RMN *Q*) and resistance (RMN *R*) during inspiration and expiration in both nostrils (Rhinospir PRO, Sibel, Barcelona, Spain), acoustic rhinometry (SER 2000, RhinoMetrics, Lynge, Denmark) to determine the minimal cross-sectional area (MCSA) and nasal volume (0–6 cm and 0–12 cm [V_{0-6} and V_{0-12}]) from both nostrils, and measurement of PNIF in L/min (Clement Clark International, Essex, UK) by recording the highest result of three manoeuvres. Throughout the study, the same trained physicians assessed nasal obstruction using these subjective and objective approaches.

Statistical analysis

Qualitative variables were analysed using proportions. Quantitative variables were analysed using measures of position, central tendency, and dispersion. Values were compared using the Mann–Whitney test. The Spearman correlation coefficient (ρ) was calculated. A 95% confidence level was set with a 5% alpha error. Statistical significance was set at $p < 0.05$. Correlations were generally considered weak if $r \leq 0.4$, moderate if $0.4 < r < 0.8$, and strong if $r > 0.8$.⁷

Results

Mean (SD) age was 37.1 (6.9) years with a range of 25–56 years. Women accounted for 61% of the sample. The values for the symptom score, VAS, active anterior rhinomanometry, acoustic rhinometry, and PNIF are shown in [Table 1](#).

The correlations between the subjective and objective methods are shown in [Table 2](#).

- (a) Correlation between subjective methods
Symptom score and VAS were moderately correlated.
- (b) Correlation between objective methods
Correlation between rhinomanometry variables
The values for RMN *Q* and RMN *R* correlated strongly in both inspiration and expiration.
Correlation between acoustic rhinometry variables
The values for V_{0-6} and V_{0-12} were moderately correlated; V_{0-6} and MCSA were weakly correlated.
Correlation between anterior rhinomanometry and acoustic rhinometry

Table 1 Subjective and objective variables in the assessment of nasal obstruction.

	Mean (SD)	Median (IQR)
Symptom score	0.7 (0.6)	1 (0–1)
Visual analogue scale, cm	2.1 (2.5)	1.3 (0–3.4)
Acoustic rhinometry, V_{0-6} cm ³	14.2 (3.9)	13.4 (11.6–16.3)
Acoustic rhinometry, V_{0-12} cm ³	54.4 (19.9)	53.2 (43.6–66.7)
Acoustic rhinometry, MCSA cm ²	0.8 (0.2)	0.8 (0.6–0.9)
RMN <i>Q</i> , inspiration (L/min)	604.9 (252.5)	569 (438–762)
RMN <i>R</i> , inspiration (Pa/L/s)	0.3 (0.2)	0.3 (0.2–0.4)
RMN <i>Q</i> , expiration (L/min)	608.5 (255.8)	568 (445.5–724)
RMN <i>R</i> , expiration (Pa/L/s)	0.3 (0.2)	0.3 (0.2–0.3)
PNIF (L/min)	122.7 (39.9)	120 (93.8–150)

VAS, visual analogue scale; V_{0-6} , volume between 0 and 6 cm; V_{0-12} , volume between 0 and 12 cm; MCSA, minimal cross-sectional area; RMN, anterior rhinomanometry; *Q*, airflow; *R*, resistance; PNIF, peak nasal inspiratory flow. Pa/L/s, pascal/litre/second; L/min, litre/minute.

MCSA correlated moderately with anterior RNM *R* in inspiration and weakly with RMN *Q* and RMN *R*. MCSA correlated weakly with all the anterior rhinomanometry values.

Correlation between acoustic rhinometry variables and PNIF variables

MCSA and PNIF were moderately correlated.

(c) Correlation between subjective and objective methods
Correlation between acoustic rhinometry and symptom score

Symptom score correlated weakly with acoustic rhinometry (V_{0-6} and MCSA).

Analysis of the correlations between patients with greater or less nasal obstruction, depending on whether they had rhinitis at the time of the evaluation or whether they had a symptom score of 0–1 vs. 2–3 or a VAS score of <4 vs. >4, revealed no differences with respect to data obtained from an analysis of the total sample (data not shown).

Discussion

We applied different subjective and objective techniques to assess nasal obstruction. The correlations between these techniques were generally weak or moderate. The most notable results were the strong correlation between the different parameters for active anterior rhinomanometry ($r > 0.8$) and moderate correlation ($0.4 < r < 0.8$) between the symptom score and VAS and between MCSA and RMN *R* and PNIF.

VAS and several symptom scores have been shown to reveal changes in the severity of perceived nasal obstruction⁵ and a certain degree of correlation.⁸ We found correlations to be moderate; therefore, we suggest that both methods are similar when applied for subjective assessment of nasal obstruction.

Opinions on the value of objective assessment of nasal obstruction are divergent.⁴ PNIF has been used in several studies and its sensitivity is comparable to that of acoustic rhinometry and active anterior rhinomanometry.^{9,10} We found a moderate correlation with MCSA and a weak correlation with active anterior rhinomanometry.

Acoustic rhinometry has been compared with active anterior rhinomanometry. Its properties enable it to measure changes that are not assessed using active anterior rhinomanometry. Furthermore, several authors have found differences between the techniques and suggested that they should complement each other.^{11–13} Our weak correlations between the different variables in each technique, despite being moderate between MCSA and resistance, reinforce the concept that these techniques assess different aspects of nasal obstruction and are therefore complementary.

Correlations between subjective and objective measurements have also been investigated. In a study involving 250 volunteers, Jones et al.¹⁴ did not find an association between the subjective sensation of nasal obstruction and active anterior rhinomanometry. Gungor et al.¹⁵ did not find correlations between VAS and acoustic rhinometry during the nasal cycle. In other studies, the subjective nature of nasal obstruction correlates better with measurements of airflow resistance (RMN) than with acoustic rhinometry.^{16,17} A strong correlation has also been observed between PNIF and the subjective sensation of nasal obstruction.¹⁸

Despite the differences between subjective and objective techniques, all the methods provide reliable results and are used in several studies to evaluate the effect of treatment. Available data enable us to say that nasal obstruction should be evaluated using at least one subjective technique and one objective technique; PNIF should be reserved for outpatient monitoring of nasal obstruction.⁵

Although André et al.⁴ suggest that the presence of nasal obstruction could improve correlations between active anterior rhinomanometry and acoustic rhinometry, we did not find differences (data not shown) in the correlations depending on the degree of nasal obstruction. Consequently, correlations do not vary in patients who perceive greater nasal obstruction according to the symptom score, VAS, or presentation of symptoms of rhinitis at the time of the examination.

In conclusion, nasal obstruction can be assessed using subjective and objective approaches. The correlations between objective techniques were moderate to strong. In addition, between subjective techniques we reported a

Table 2 Correlations between subjective and objective variables of nasal obstruction (Spearman rho).

	Symptom score	V ₀₋₆	V ₀₋₁₂	MCSA	RMN Q _i inspiration	RMN R _i inspiration	RMN Q _e expiration	RMN R _e expiration	VAS	PNIF
Symptom score	1.000									
V ₀₋₆	-0.237 ^a	1.000								
V ₀₋₁₂	0.658 ^b	0.658 ^b	1.000							
MCSA	-0.265 ^a	0.248 ^b	-0.136	1.000						
RMN Q _i inspiration	0.023	0.023	0.057	0.300 ^b	1.000					
RMN R _i inspiration	0.184	0.064	0.143	-0.490 ^b	-0.966 ^b	1.000				
RMN Q _e expiration	-0.205	0.021	0.061	0.257 ^b	0.918 ^b	-0.907 ^b	1.000			
RMN R _e expiration	0.130	0.088	0.133	-0.32 ^b	-0.863 ^b	-0.893 ^b	1.000			
VAS	0.686 ^b	-0.114	0.159	-0.151	-0.089	-0.090	-0.041	1.000		
PNIF	-0.152	0.083	-0.065	0.459 ^b	0.257 ^b	0.188 ^a	-0.293 ^b	-0.187	1.000	

V₀₋₆, volume between 0 and 6 cm; V₀₋₁₂, volume between 0 and 12 cm; MCSA, minimal cross-sectional area; RMN, anterior rhinomanometry; Q_i, airflow; R_i, resistance; VAS, visual analogue scale; PNIF, peak nasal inspiratory flow.

^a Correlation is significant at 0.05.

^b Correlation is significant at 0.01.

moderate correlation. Finally, the correlations between the subjective and objective techniques were weak and absent. These findings suggest that each of the techniques assess different aspects of nasal obstruction making them complementary.

Ethical disclosures

Patients' data protection. Confidentiality of data. The authors declare that they have followed the protocols of their work centre on the publication of patient data and that all the patients included in the study have received sufficient information and have given their informed consent in writing to participate in that study.

Right to privacy and informed consent. Right to privacy and informed consent. The authors have obtained the informed consent of the patients and/or subjects mentioned in the article. The author for correspondence is in possession of this document.

Protection of human subjects and animals in research. Protection of human and animal subjects. The authors declare that the procedures followed were in accordance with the regulations of the responsible Clinical Research Ethics Committee and in accordance with those of the World Medical Association and the Helsinki Declaration.

Conflict of interest

The authors have no conflict of interests to declare.

Acknowledgement

We thank Pablo Pons from Content Ed Net Madrid for Editorial Assistance.

References

1. Jones N. The nose and paranasal sinuses physiology and anatomy. *Adv Drug Deliv Rev.* 2001;51:5-19.
2. Hirschberg A. Rhinomanometry: an update. *ORL J Otorhinolaryngol Relat Spec.* 2002;64:263-7.
3. Eccles R. Nasal airway resistance and nasal sensation of airflow. *Rhinol Suppl.* 1992;14:86-90.
4. André RF, Vuyk HD, Ahmed A, Graamans K, Nolst Trenité GJ. Correlation between subjective and objective evaluation of the nasal airway. A systematic review of the highest level of evidence. *Clin Otolaryngol.* 2009;34:518-25.
5. van Spronsen E, Ingels KJ, Jansen AH, Graamans K, Fokkens WJ. Evidence-based recommendations regarding the differential diagnosis and assessment of nasal congestion: using the new GRADE system. *Allergy.* 2008;63:820-33.
6. Kim CS, Moon BK, Jung DH, Min YG. Correlation between nasal obstruction symptoms and objective parameters of acoustic rhinometry and rhinomanometry. *Auris Nasus Larynx.* 1998;25:45-8.
7. Shi R, Conrad SA. Correlation and regression analysis. *Ann Allergy Asthma Immunol.* 2009;103:535-41.
8. Linder A. Symptom scores as measures of the severity of rhinitis. *Clin Allergy.* 1988;18:29-37.

9. Harar RP, Kalan A, Kenyon GS. Assessing the reproducibility of nasal spirometry parameters in the measurement of nasal patency. *Rhinology*. 2001;39:211–4.
10. Holmström M, Scadding GK, Lund VJ, Darby YC. Assessment of nasal obstruction. A comparison between rhinomanometry and nasal inspiratory peak flow. *Rhinology*. 1990;28:191–6.
11. Alu K. The current methodological approaches in assessing nasal breathing function. *Vestn Otorinolaringol*. 1998;4:51–2.
12. Roithmann R, Cole P, Chapnik J, Barreto SM, Szalai JP, Zamel N. Acoustic rhinometry, rhinomanometry, and the sensation of nasal patency: a correlative study. *J Otolaryngol*. 1994;23:454–8.
13. Kesavanathan J, Swift DL, Fitzgerald TK, Permutt T, Bascom R. Evaluation of acoustic rhinometry and posterior rhinomanometry as tools for inhalation challenge studies. *J Toxicol Environ Health*. 1996;48:295–307.
14. Jones A, Willatt D, Durham L. Nasal airflow: resistance and sensation. *J Laryngol Otol*. 1989;103:909–11.
15. Gungor A, Moinuddin R, Nelson RH, Corey JP. Detection of the nasal cycle with acoustic rhinometry: techniques and applications. *Otolaryngol Head Neck Surg*. 1999;120:238–47.
16. Szücs E, Clement PA. Acoustic rhinometry and rhinomanometry in the evaluation of nasal patency of patients with nasal septal deviation. *Am J Rhinol*. 1998;12:345–52.
17. Numminen J, Ahtinen M, Huhtala H, Rautiainen M. Comparison of rhinometric measurement methods in intranasal pathology. *Rhinology*. 2003;41:65–8.
18. Morrissey MS, Alun-Jones T, Hill J. The relationship of peak inspiratory airflow to subjective airflow in the nose. *Clin Otolaryngol Allied Sci*. 1990;15:447–51.