

Amphibians and reptiles from two localities in the northern Andes of Colombia

Adriana Restrepo, Claudia Molina-Zuluaga, Juan P. Hurtado, Carlos M. Marín, Juan M. Daza

Grupo Herpetológico de Antioquia, Instituto de Biología, Universidad de Antioquia, A.A. 1226, Medellín, Colombia.

Corresponding author: Juan M. Daza, jumadaza@gmail.com

Abstract

The northern Andes are a geographically, climatically, and ecologically complex area that harbors a high number of species and presents a high level of endemism. However, the landscape of this region has been highly transformed by human activities, including agriculture, cattle grazing, urbanization, and recently, the construction of large infrastructure projects. This study presents a list of amphibians and reptiles compiled from field surveys carried out between 2006 and 2016 in the protected areas surrounding 2 hydroelectric projects located on the eastern flank of the northern Cordillera Central in Colombia. The checklist is comprised of 101 species (43 amphibians and 58 reptiles), including 17 endemic to Colombia and 4 threatened species. The high species density, high representativeness of the regional biodiversity, and high level of amphibian endemism make these 2 protected areas important for conservation of the regional herpetofauna.

Key words

Endemism; frogs; hydroelectric dam; lizards; snakes; species richness.

Academic editor: Sebastian Lotzkatz | Received 3 December 2015 | Accepted 4 June 2017 | Published 28 July 2017


Citation: Restrepo A, Molina-Zuluaga C, Hurtado JP, Marín CM, Daza JM (2017) Amphibians and reptiles from two localities in the northern Andes of Colombia. *Check List* 13 (4): 203–237. <https://doi.org/10.15560/13.4.203>

Introduction

The Andes mountain range extends for 5000 km along the western coast of South America and 3 main domains are recognized: southern, central and northern Andes (Gregory-Wodzicki 2000). The northern Andes reach their maximum geomorphological complexity in Colombia, where 3 mountain chains (the Western, Central, and Eastern cordilleras), are separated by the sedimentary basins of the Cauca and Magdalena rivers (Kattan et al. 2004). Three major historical events have shaped the biological diversity in this region: (1) the uplift of mountain ranges in a complex series of orogenic processes generating new environments and causing vicariant events, (2)

the final closure of the Isthmus of Panama that allowed the biotic interchange between 2 continents, and (3) the climatic fluctuations during the Pleistocene that lead to cycles of range contraction and expansion resulting in the fragmentation and isolation of natural populations, with subsequent speciation and radiation events (Duellman 1999, Kattan et al. 2004). Thus, the diversity of ecological settings generated by this physiographic complexity has made the northern Andes one of the most biodiverse regions on the planet with an exceptional number of endemic plants and vertebrates (Fjeldsa 1994, Duellman 1999, Myers et al. 2000).

The northern Andes are also under continuous and intense anthropogenic pressure. About 70% of the human



Figures 1. Study area showing the location in Colombia and the northern Cordillera Central. On the right, the two localities (Jaguas and San Carlos) are expanded and black dots represent the sampling sites. Colors represent cover type.

population in Colombia is concentrated in this region and agricultural activities, cattle grazing and infrastructure development (i.e. urbanization, dam construction, and roads) are among the main causes of habitat loss and fragmentation (Etter and van Wyngaarden 2000, Fjeldsa et al. 2005). On the eastern flank of the northern Cordillera Central, several hydroelectric projects involving dam constructions, roads, and power lines were developed in the last 2 decades. As a mitigation strategy, Colombia's environmental legislation demands that these large infrastructure projects compensate the negative impacts through the implementation of effective conservation and restoration plans by preserving and monitoring the biodiversity in the surrounding areas (Minambiente 2012). As a consequence, these habitats, as virtually the only remnants left, represent a last-minute opportunity to make complete species checklists for the northern portion of the central Andes of Colombia.

Based on more than a decade of field surveys, we provide the species list of amphibians and reptiles found in the protected areas of 2 main hydroelectric projects located on the northeastern Cordillera Central in Colombia.

Methods

Study sites. This study was conducted in the northern Andes at 2 sites above 500 m elevation in the Cordillera Central in Antioquia department, Colombia (Fig. 1). The Jaguas site ($06^{\circ}26'06.0''$ N, $075^{\circ}00'57.6''$ W) includes a 4000 ha private protected area with habitats in different successional stages, within the premontane wet forest life zone (bmh-PM, Holdridge 1964). The altitude ranges between 1100 and 1300 m above sea level. The temperature ranges between 18 and 24 °C and the annual rainfall between 2000 and 4000 mm. The San Carlos site ($06^{\circ}12'57.6''$ N, $074^{\circ}51'21.6''$ W) includes a private protected area of 3000 ha, within the tropical moist forest life zone (bh-T, Holdridge 1964). The altitude is between 600 and 950 m above sea level. It has a mean temperature of 24 °C and the annual rainfall ranges between 2000 and 4000 mm. The 2 sites are under a bimodal rainfall regime with 1 rainy season between March and May and 1 between September and November (Poveda et al. 2005).

Data collection. Between 2006 and 2016, the complete species inventory has been an ongoing goal as part of the

monitoring plan implemented by the company ISAGEN S.A. Amphibians and reptiles were searched for in all possible microhabitats including diurnal and nocturnal sampling events during rainy, dry, and transition seasons. In total, 49 sampling points were distributed in different vegetation types.

Collected specimens were euthanized with 2% Roxicain[®], fixed in 10% formalin solution, and preserved in 70% ethanol following the protocol by Cortez et al. (2006). Voucher specimens are deposited in the Museo de Herpetología Universidad de Antioquia (MHUA), in Medellín, Colombia (Instituto Alexander von Humboldt registry # 80, since the year 2000). All procedures were conducted under permit No. 112–0046 granted by the Corporación Autónoma Regional CORNARE. We list the collected and examined specimens in Table 1 (amphibians) and 2 (reptiles). All coordinates use the WGS84 datum.

Amphibian nomenclature follows Frost (2016) and reptile nomenclature follows Uetz and Hošek (2016). We reviewed the original descriptions of the species along with the MHUA reference collection to corroborate the taxonomic identifications. We included in the list only the species having voucher specimens or photographs that allow unambiguous identification of taxa. We follow the species distribution provided by IUCN (2016) for amphibians and Uetz and Hošek (2016) for reptiles.

Results

Amphibians

From the 2 protected areas, we report 43 amphibian species representing 13 families and 25 genera (Table 3). In the protected area of the Jaguas locality, we report 37 amphibian species from 11 families. The most diverse lineages were Hylidae with 12 species (32%) and Craugastoridae with 6 species (16%). Eleven species are endemic to Colombia (30%), and according to the IUCN Red Lists, 1 is classified as Endangered, 2 as Vulnerable, and 1 as Near Threatened (IUCN 2016). In comparison, the San Carlos site has 34 amphibian species grouped in 11 families. Again, the most diverse groups were Hylidae with 11 species (32%) and Craugastoridae with 6 species (18%). Eleven species from San Carlos locality are endemic to Colombia (32%) and 2 are listed as Vulnerable (IUCN 2016). Jaguas and San Carlos localities shared 28 amphibian species.

Rheobates pseudopalmaris (Rivero & Serna, 2000): Figure 2A

Colostethus pseudopalmaris Rivero and Serna 2000: 55.

Rheobates pseudopalmaris — Grant et al. 2006: 159.

Material examined. Table 1.

A moderate-sized frog with cryptic, brown or gray dorsal coloration; dorsal skin granular posteriorly; toe webbing extensive; finger I shorter than finger II; pale oblique lateral stripe present or absent and pale dorso-lateral and ventrolateral stripes absent. Rivero and Serna

(2000) proposed *R. pseudopalmaris* as sister species of *R. palmaris*. However, Grant et al. (2006) question the validity of morphological characters used to distinguish this species from *R. palmaris* and consider that both species could be conspecific given that the type locality of *R. pseudopalmaris* (Amalfi, Antioquia) lies within the known distribution of *R. palmaris* on the eastern slope of the Central Cordillera. Here, we assign our samples to *R. pseudopalmaris* according to Muñoz-Ortiz et al. (2015) who suggested, based on molecular information, that this species could be redescribed restricting *R. pseudopalmaris* to the Central Cordillera and *R. palmaris* to the Eastern Cordillera.

Rhinella alata (Thomiot, 1884): Figure 2B

Bufo alatus Thomiot 1884: 151.

Rhinella alata — Frost et al. 2006: 365

Material examined. Table 1.

Rhinella alata is a small-sized toad with skin of dorsum bearing a mixture of warts, pustules and minute tubercles; supraorbital crests low and thick, continuous with preorbital crests; dorsal coloration ranging from light gray or brown to dark gray or brown, with irregular black and yellowish marks (a mid-dorsal line is usually present) and coloration of ventral surfaces cream to yellowish-cream with irregular darker marks arranged in diverse patterns (dos Santos et al. 2015). Populations of *Rhinella margaritifera* species group distributed on Magdalena river valley have commonly been referred to as *R. margaritifera* (see Burbano et al. 2016). Nevertheless, although *R. alata* is closely related to *R. margaritifera* (distributed in the upper Amazon basin of Peru and Ecuador), this species may be easily distinguished by differences in body size (*R. margaritifera* being significant larger than *R. alata*) and the relative size of the cranial crest (larger in *R. margaritifera* than in *R. alata*).

Rhinella horribilis (Wiegmann, 1833): Figure 2C

Bufo horribilis Wiegmann 1833: 654.

Bufo marinus — Peters 1873: 618.

Rhinella horribilis — Acevedo et al. 2016: 584.

Material examined. Table 1.

Recently, Acevedo et al. (2016) proposed that populations of *Rhinella marina* (Linnaeus, 1758) distributed west of the Andean cordilleras should be considered a different species from the eastern populations. Therefore, the eastern populations should maintain the name *R. marina* and *R. horribilis* should be revalidated for the western populations. According to this concept, the Magdalena river valley populations correspond to *R. horribilis* (Wiegmann, 1833).

Cochranella resplendens (Lynch & Duellman, 1973): Figure 2D

Centrolenella resplendens Lynch and Duellman 1973: 51.

Cochranella resplendens — Ruiz-Carranza and Lynch 1991: 23.

Material examined. Table 1.

Espadarana prosoblepon (Boettger, 1892): Figure 2E

Hyla prosoblepon Boettger 1892: 45.

Centrolene prosoblepon Noble 1924: 66.

Table 1. Voucher list of amphibians of Jaguas and San Carlos hydroelectric protected areas with geographic coordinates.

Taxon	Voucher	Latitude	Longitude	Taxon	Voucher	Latitude	Longitude
AMPHIBIA					MHUA-A 10688	6.19176	-74.80257
ANURA				<i>Hyalinobatrachium</i>	MHUA-A 07075	6.23108	-74.87661
Aromobatidae				<i>aureoguttatum</i>	MHUA-A 07076	6.23108	-74.87661
<i>Rheobates pseudopalmaris</i>	MHUA-A 07845	6.38871	-75.02555	<i>Hyalinobatrachium</i>	MHUA-A 07198	6.21734	-74.81262
	MHUA-A 08485	6.36725	-75.02723	<i>fleischmanni</i>			
	MHUA-A 08491	6.39128	-75.03640	<i>Rulyrana susatamai</i>	MHUA-A 09539	6.36725	-75.02723
	MHUA-A 09425	6.39128	-75.03640	<i>Sachatamia punctulata</i>	MHUA-A 09728	6.32866	-75.01155
	MHUA-A 09426	6.36725	-75.02723		MHUA-A 09764	6.29817	-74.91981
	MHUA-A 09538	6.39128	-75.03640		MHUA-A 10803	6.28806	-74.82722
	MHUA-A 09758	6.29817	-74.91981		MHUA-A 10804	6.28806	-74.82722
	MHUA-A 10599	6.15750	-75.04325	<i>Sachatamia</i> sp.	MHUA-A 07123	6.37141	-75.04253
	MHUA-A 10641	6.15750	-75.04325		MHUA-A 07842	6.38871	-75.02555
	MHUA-A 10643	6.15829	-75.04531		MHUA-A 07844	6.38871	-75.02555
	MHUA-A 10644	6.15829	-75.04531		MHUA-A 08205	6.39202	-75.03650
	MHUA-A 10646	6.15829	-75.04531		MHUA-A 08216	6.38539	-75.02202
	MHUA-A 10650	6.15829	-75.04531		MHUA-A 08217	6.39170	-75.03626
	MHUA-A 10651	6.15750	-75.04325		MHUA-A 08229	6.39202	-75.03650
	MHUA-A 10652	6.15763	-75.04316		MHUA-A 08231	6.39202	-75.03650
	MHUA-A 10653	6.15763	-75.04316		MHUA-A 08232	6.39202	-75.03650
	MHUA-A 10654	6.15829	-75.04531		MHUA-A 08233	6.39202	-75.03650
	MHUA-A 10662	6.15829	-75.04531		MHUA-A 08336	6.39115	-75.03641
	MHUA-A 10726	6.20908	-74.85436		MHUA-A 08337	6.39115	-75.03641
					MHUA-A 10615	6.37172	-75.04534
					MHUA-A 10682	6.36662	-75.02798
Bufonidae				Craugastoridae			
<i>Rhinella alata</i>	MHUA-A 09731	6.32711	-75.01186	<i>Craugastor metriosistus</i>	MHUA-A 10468	6.24144	-74.86976
	MHUA-A 09732	6.32758	-75.01176		MHUA-A 10507	6.23974	-74.86880
	MHUA-A 09739	6.27667	-74.93102		MHUA-A 10586	6.15763	-75.04316
	MHUA-A 09766	6.29231	-75.03969		MHUA-A 10587	6.36767	-75.02635
	MHUA-A 08492	6.39128	-75.03640		MHUA-A 10588	6.38518	-75.02140
	MHUA-A 06979	6.30032	-75.06698		MHUA-A 10589	6.24144	-74.86976
	MHUA-A 07930	6.38871	-75.02555		MHUA-A 10590	6.20938	-74.85352
	MHUA-A 08225	6.36725	-75.02723		MHUA-A 10656	6.20908	-74.85436
	MHUA-A 08330	6.36725	-75.02723		MHUA-A 10660	6.20908	-74.85436
	MHUA-A 08486	6.39128	-75.03640		MHUA-A 10800	6.28806	-74.82722
	MHUA-A 10671	6.18324	-74.79362		MHUA-A 10808	6.15844	-75.04433
	MHUA-A 10673	6.18325	-74.79362		MHUA-A 10809	6.15865	-75.04447
	MHUA-A 10674	6.18280	-74.19330	<i>Pristimantis fallax</i>	MHUA-A 07125	6.20671	-74.85500
	MHUA-A 10676	6.19224	-74.80183		MHUA-A 10523	6.20908	-74.85436
	MHUA-A 10677	6.15671	-75.04108	<i>Pristimantis gaigei</i>	MHUA-A 07118	6.24320	-74.87138
	MHUA-A 10679	6.18336	-74.79308		MHUA-A 07208	6.21854	-74.83727
	MHUA-A 10722	6.13321	-75.09676		MHUA-A 07209	6.21626	-74.83435
<i>Rhinella horribilis</i>					MHUA-A 08348	6.19062	-74.80474
Centrolenidae					MHUA-A 08497	6.36725	-75.02723
<i>Cochranella resplendens</i>	MHUA-A 9540	6.36725	-75.02723		MHUA-A 09750	6.27681	-74.93975
<i>Espadarana prosoblepon</i>	MHUA-A 06978	6.30032	-75.06698		MHUA-A 09751	6.27679	-74.93085
	MHUA-A 07124	6.36699	-75.02892		MHUA-A 10495	6.17844	-74.80273
	MHUA-A 07420	6.36930	-75.02520		MHUA-A 10516	6.19224	-74.80183
	MHUA-A 07421	6.36930	-75.02520		MHUA-A 10802	6.28806	-74.82722
	MHUA-A 08212	6.38871	-75.02335	<i>Pristimantis jaguensis</i>	MHUA-A 07026	6.38871	-75.02335
	MHUA-A 08213	6.38871	-75.02335		MHUA-A 07027	6.38871	-75.02335
	MHUA-A 08214	6.38871	-75.02335		MHUA-A 07028	6.38871	-75.02335
	MHUA-A 08226	6.36725	-75.02723		MHUA-A 07030	6.38871	-75.02335
	MHUA-A 08230	6.39202	-75.03650		MHUA-A 07031	6.38871	-75.02335
	MHUA-A 08329	6.36725	-75.02723		MHUA-A 07032	6.38871	-75.02335
	MHUA-A 08338	6.39115	-75.03641		MHUA-A 07033	6.38871	-75.02335
	MHUA-A 08493	6.39128	-75.03640		MHUA-A 07238	6.38871	-75.02335
	MHUA-A 09723	6.32845	-75.01171		MHUA-A 07246	6.38871	-75.02335
	MHUA-A 09724	6.32734	-75.01160		MHUA-A 07248	6.38871	-75.02335
	MHUA-A 09729	6.32734	-75.01160		MHUA-A 07249	6.38871	-75.02335
	MHUA-A 09730	6.32734	-75.01160		MHUA-A 07250	6.38871	-75.02335
	MHUA-A 10629	6.38518	-75.02140		MHUA-A 07251	6.38871	-75.02335
	MHUA-A 10631	6.37172	-75.04534		MHUA-A 07252	6.38871	-75.02335
	MHUA-A 10632	6.35150	-74.99528		MHUA-A 07253	6.38871	-75.02335
	MHUA-A 10633	6.35150	-74.99528		MHUA-A 07254	6.38871	-75.02335
	MHUA-A 10634	6.35150	-74.99528		MHUA-A 07255	6.38871	-75.02335
	MHUA-A 10635	6.19176	-74.80257		MHUA-A 07256	6.35812	-74.99367
	MHUA-A 10636	6.38518	-75.02140		MHUA-A 07257	6.35812	-74.99367
	MHUA-A 10638	6.35150	-74.99528				
	MHUA-A 10681	6.36662	-75.02798				
	MHUA-A 10686	6.35150	-74.99528				

Continued

Table 1. Continued.

Taxon	Voucher	Latitude	Longitude	Taxon	Voucher	Latitude	Longitude
<i>Pristimantis penelopus</i>	MHUA-A 07415	6.38871	-75.02335	<i>Pristimantis viejas</i>	MHUA-A 10503	6.15801	-75.04480
	MHUA-A 07417	6.38871	-75.02335		MHUA-A 10522	6.15829	-75.04379
	MHUA-A 07837	6.38871	-75.02555		MHUA-A 10525	6.15829	-75.04379
	MHUA-A 10518	6.36662	-75.02798		MHUA-A 10591	6.24279	-74.86818
	MHUA-A 07034	6.34818	-75.00237		MHUA-A 07058	6.34818	-75.00237
	MHUA-A 07035	6.34818	-75.00237		MHUA-A 07135	6.24739	-74.86938
	MHUA-A 07424	6.36930	-75.02520		MHUA-A 08335	6.39202	-75.03650
	MHUA-A 08588	6.36728	-75.02697		MHUA-A 08341	6.38528	-75.02158
	MHUA-A 09727	6.32860	-75.01146		MHUA-A 08538	6.40999	-75.14157
	MHUA-A 09733	6.32831	-75.01188		MHUA-A 09742	6.29610	-74.91944
	MHUA-A 09734	6.32866	-75.01155		MHUA-A 09743	6.27721	-74.92910
	MHUA-A 09754	6.29647	-74.91934		MHUA-A 09744	6.27721	-74.92910
	MHUA-A 09756	6.29771	-74.91969		MHUA-A 09755	6.29631	-74.91931
	MHUA-A 09760	6.29817	-74.91981		MHUA-A 09759	6.29648	-74.91931
	MHUA-A 10461	6.15750	-75.04325		MHUA-A 10465	6.38518	-75.02140
	MHUA-A 10462	6.36697	-75.02733	MHUA-A 10474	6.37198	-75.04549	
	MHUA-A 10463	6.36701	-75.02718	MHUA-A 10475	6.15750	-75.04325	
	MHUA-A 10464	6.37194	-75.04556	MHUA-A 10521	6.37198	-75.04549	
	MHUA-A 10466	6.36662	-75.02798	Dendrobatidae			
	MHUA-A 10467	6.15728	-75.42740	<i>Colostethus inguinalis</i>	MHUA-A 00223	6.21591	-74.85079
	MHUA-A 10469	6.35150	-74.99528		MHUA-A 07072	6.22342	-74.84733
	MHUA-A 10470	6.35150	-74.99528		MHUA-A 10645	6.18324	-74.79362
	MHUA-A 10476	6.35150	-74.99528		MHUA-A 10600	6.18370	-74.79353
	MHUA-A 10478	6.36662	-75.02798		MHUA-A 10642	6.18370	-74.79353
	MHUA-A 10479	6.37188	-75.04533		MHUA-A 10647	6.19976	-74.80432
	MHUA-A 10480	6.35150	-74.99528		MHUA-A 10648	6.18324	-74.79362
	MHUA-A 10482	6.36697	-75.02733		MHUA-A 10649	6.24307	-74.86771
	MHUA-A 10483	6.38518	-75.02140		MHUA-A 10661	6.18370	-74.79353
	MHUA-A 10485	6.15822	-75.04457	<i>Colostethus aff. fraterdanieli</i>	MHUA-A 07396	6.38871	-75.02335
	MHUA-A 10486	6.15723	-75.04282		MHUA-A 07411	6.38871	-75.02335
	MHUA-A 10487	6.36701	-75.02718		MHUA-A 07416	6.36930	-75.02520
	MHUA-A 10488	6.37172	-75.04534		MHUA-A 07419	6.38871	-75.02335
	MHUA-A 10490	6.15754	-75.04295		MHUA-A 08472	6.38528	-75.02158
MHUA-A 10491	6.36662	-75.02798		MHUA-A 08587	6.36728	-75.02697	
MHUA-A 10494	6.37188	-75.04533		MHUA-A 09533	6.36725	-75.02723	
MHUA-A 10496	6.15829	-75.04531		MHUA-A 09534	6.36725	-75.02723	
MHUA-A 10497	6.15773	-75.04398		MHUA-A 09535	6.36725	-75.02723	
MHUA-A 10500	6.37158	-75.04247	<i>Dendrobates truncatus</i>	MHUA-A 09536	6.38537	-75.02166	
MHUA-A 10501	6.38518	-75.02140		MHUA-A 08223	6.19055	-74.80480	
MHUA-A 10502	6.36662	-75.02798		MHUA-A 07410	6.22020	-74.84713	
MHUA-A 10504	6.20908	-74.85436		MHUA-A 08210	6.19570	-74.81670	
MHUA-A 10505	6.20908	-74.85436		MHUA-A 09740	6.27738	-74.92780	
MHUA-A 10506	6.15849	-75.04496		MHUA-A 10695	6.18360	-74.79343	
MHUA-A 10510	6.15750	-75.04325	Eleutherodactylidae				
MHUA-A 10511	6.15829	-75.04531	<i>Diasporus anthrax</i>	MHUA-A 07237	6.34818	-75.00237	
MHUA-A 10513	6.35150	-74.99528	<i>Diasporus gularis</i>	MHUA-A 07258	6.38871	-75.02335	
MHUA-A 10515	6.15723	-75.04282		MHUA-A 07405	6.38871	-75.02335	
MHUA-A 10517	6.15784	-75.04414		MHUA-A 07406	6.38871	-75.02335	
MHUA-A 10519	6.37188	-75.04533		MHUA-A 07407	6.38871	-75.02335	
MHUA-A 10520	6.36662	-75.02798		MHUA-A 07412	6.38871	-75.02335	
MHUA-A 10527	6.36697	-75.02733		MHUA-A 07413	6.38871	-75.02335	
MHUA-A 10559	6.15829	-75.04531		MHUA-A 07839	6.38871	-75.02555	
MHUA-A 10560	6.37194	-75.04556		MHUA-A 08211	6.38871	-75.02335	
MHUA-A 10592	6.20908	-74.85436		MHUA-A 08494	6.38528	-75.02158	
MHUA-A 10724	6.20908	-74.85436		MHUA-A 08496	6.39128	-75.03640	
MHUA-A 10725	6.20908	-74.85436		MHUA-A 10593	6.38518	-75.02140	
MHUA-A 10811	6.13846	-75.06631		MHUA-A 10594	6.20908	-74.85436	
<i>Pristimantis taeniatus</i>	MHUA-A 07211	6.18737	-74.80460	MHUA-A 10595	6.20908	-74.85436	
	MHUA-A 07212	6.19743	-74.80363	MHUA-A 10794	6.37341	-75.02509	
	MHUA-A 07215	6.21569	-74.81371	Hyllidae			
	MHUA-A 07234	6.35447	-74.98836	<i>Boana boans</i>	MHUA-A 07193	6.22895	-74.87296
	MHUA-A 07423	6.36930	-75.02520		MHUA-A 07838	6.38871	-75.02555
	MHUA-A 08484	6.36725	-75.02723		MHUA-A 07840	6.36664	-75.02746
	MHUA-A 09537	6.37540	-74.98122		MHUA-A 09505	6.36081	-74.98657
	MHUA-A 09757	6.29681	-74.91933		MHUA-A 09752	6.28320	-74.92413
	MHUA-A 09769	6.29371	-75.04156		MHUA-A 10564	6.19175	-74.80257
	MHUA-A 09770	6.29371	-75.04156	<i>Boana xerophylla</i>	MHUA-A 10567	6.19725	-74.80433
	MHUA-A 10471	6.15829	-75.04379		MHUA-A 10566	6.19725	-74.80433

Continued

Table 1. Continued.

Taxon	Voucher	Latitude	Longitude	Taxon	Voucher	Latitude	Longitude	
<i>Dendropsophus ebraccatus</i>	MHUA-A 10568	6.19725	-74.80433		MHUA-A 10556	6.19725	-74.80433	
	MHUA-A 10569	6.19725	-74.80433		MHUA-A 10581	6.19725	-74.80433	
	MHUA-A 10571	6.18406	-74.79338		MHUA-A 10582	6.19725	-74.80433	
	MHUA-A 07196	6.19744	-74.80397		MHUA-A 10583	6.19725	-74.80433	
	MHUA-A 10692	6.19725	-74.80433		MHUA-A 10584	6.19725	-74.80433	
	MHUA-A 10528	6.19725	-74.80433		MHUA-A 10585	6.15781	-75.04087	
	MHUA-A 10529	6.35171	-74.99379	<i>Scinax</i> sp.	MHUA-A 07843	6.38871	-75.02555	
	MHUA-A 10532	6.35171	-74.99379		MHUA-A 08228	6.36081	-74.98657	
	MHUA-A 10533	6.35171	-74.99379	<i>Scinax rostratus</i>	MHUA-A 07233	6.35812	-74.99367	
	MHUA-A 10535	6.19725	-74.80433		MHUA-A 10799	6.28806	-74.82722	
	MHUA-A 10538	6.19725	-74.80433	<i>Scinax ruber</i>	MHUA-A 01685	6.49061	-74.83683	
	MHUA-A 10542	6.19725	-74.80433		MHUA-A 01686	6.49061	-74.83683	
	MHUA-A 10544	6.19725	-74.80433		MHUA-A 09500	6.36081	-74.98657	
	MHUA-A 10551	6.19725	-74.80433	<i>Smilisca phaeota</i>	MHUA-A 08539	6.40999	-75.14157	
	MHUA-A 10553	6.19725	-74.80433		MHUA-A 09503	6.36081	-74.98657	
	MHUA-A 10555	6.35171	-74.99379		MHUA-A 09746	6.27700	-74.92966	
	MHUA-A 10557	6.35171	-74.99379		MHUA-A 09761	6.29143	-74.93671	
	<i>Dendropsophus norandinus</i>	MHUA-A 08235	6.39202	-75.03650		MHUA-A 09762	6.29143	-74.93671
		MHUA-A 08488	6.39128	-75.03640		MHUA-A 09767	6.29243	-75.04211
MHUA-A 08589		6.39214	-75.03689		MHUA-A 10572	6.15743	-75.04177	
MHUA-A 08590		6.39214	-75.03689		MHUA-A 10573	6.38518	-75.02140	
MHUA-A 08591		6.39214	-75.03689		MHUA-A 10574	6.24045	-74.86977	
MHUA-A 10530		6.15784	-75.04414		MHUA-A 10575	6.24045	-74.86977	
MHUA-A 10531		6.15784	-75.04414		MHUA-A 10576	6.35171	-74.99379	
MHUA-A 10534		6.15750	-75.04325		MHUA-A 10577	6.24045	-74.86977	
MHUA-A 10536		6.15801	-75.04480		MHUA-A 10578	6.35171	-74.99379	
MHUA-A 10537		6.15784	-75.04414		MHUA-A 10579	6.35171	-74.99379	
MHUA-A 10539		6.36798	-74.99639		MHUA-A 10580	6.15850	-75.04436	
MHUA-A 10540		6.36798	-74.99639		MHUA-A 10727	6.20908	-74.85436	
MHUA-A 10541		6.15750	-75.04325					
MHUA-A 10543		6.15850	-75.04436	Leptodactylidae				
MHUA-A 10545		6.15829	-75.04531	<i>Engystomops pustulosus</i>	MHUA-A 09726	6.32913	-75.01398	
MHUA-A 10546		6.15742	-75.04264		MHUA-A 10602	6.18406	-74.79338	
MHUA-A 10547		6.15750	-75.04325		MHUA-A 10603	6.19725	-74.80433	
MHUA-A 10548		6.15754	-75.04295		MHUA-A 10604	6.19725	-74.80433	
MHUA-A 10549		6.36798	-74.99639		MHUA-A 10605	6.19725	-74.80433	
MHUA-A 10550		6.36798	-74.99639		MHUA-A 10606	6.19725	-74.80433	
MHUA-A 10554	6.36798	-74.99639	<i>Leptodactylus colombiensis</i>	MHUA-A 07197	6.19144	-74.80397		
MHUA-A 10561	6.15850	-75.04436		MHUA-A 09501	6.39128	-75.03640		
MHUA-A 10562	6.15784	-75.04414		MHUA-A 10806	6.15651	-75.04146		
<i>Dendropsophus subocularis</i>	MHUA-A 07408	6.38871	-75.02335		MHUA-A 10807	6.15651	-75.04146	
	MHUA-A 07409	6.38871	-75.02335	<i>Leptodactylus fuscus</i>	MHUA-A 01687	6.49061	-74.83683	
	MHUA-A 07834	6.38871	-75.02555		MHUA-A 10598	6.38747	-74.99274	
	MHUA-A 08206	6.39202	-75.03650		MHUA-A 10790	6.38788	-74.99348	
	MHUA-A 08339	6.39128	-75.03640		MHUA-A 10791	6.38788	-74.99348	
	MHUA-A 08340	6.39128	-75.03640					
	MHUA-A 08490	6.39128	-75.03640	CAUDATA				
				Plethodontidae				
<i>Hyloscirtus palmeri</i>	MHUA-A 06977	6.30032	-75.06698	<i>Bolitoglossa ramosi</i>	MHUA-A 07899	6.39100	-75.02100	
	MHUA-A 07073	6.23108	-74.84733					
	MHUA-A 09504	6.36081	-74.98657	GYMNOPHIONA				
	MHUA-A 10563	6.15766	-75.04323	Caeciliidae				
	MHUA-A 10693	6.23974	-74.86880	<i>Caecilia thompsoni</i>	MHUA-A 07192	6.62890	-75.66244	
	MHUA-A 10805	6.15665	-75.04128		MHUA-A 07115	6.51013	-75.04172	
	MHUA-A 10810	6.15865	-75.04447	Siphonopidae				
				<i>Microcaecilia pricei</i>	MHUA-A 10728	6.29648	-74.82229	
<i>Phyllomedusa venusta</i>	MHUA-A 07121	6.21284	-74.82631	Rhinatreumatidae				
	MHUA-A 07195	6.22895	-74.87296	<i>Epicrionops parkeri</i>	MHUA-A 07191	6.22895	-74.87296	
				MHUA-A 10720	6.20938	-74.85352		

Espadarana prosoblepon — Guayasamin et al. 2009: 33.

Material examined. Table 1.

Hyalinobatrachium aureoguttatum (Barrera-Rodríguez & Ruíz-Carranza, 1989): Figure 2F

Centrolenella aureoguttata Barrera-Rodríguez and Ruíz-Carranza 1989: 77.

Hyalinobatrachium aureoguttatum — Ruíz-Carranza and Lynch 1991: 23.

Material examined. Table 1.

Small glass frog which exhibits complete ventral transparency; transparent pericardium; white hepatic and visceral peritonea; white bones; iris silvery at the center, yellow at the edges with numerous black spots (Guayasamin et al. 2009). *Hyalinobatrachium aureoguttatum* is superficially similar to *H. valeroi*. Nevertheless, in preservative *H. aureoguttatum* differs from *H. valeroi* in having usually either well-defined rounded spots or elongate blotches of bright white on yellow background




Figure 2. Amphibian species present in Jaguas and San Carlos hydroelectric protected areas. **A:** *Rheobates pseudopalmatus* MHUAA 8491. **B:** *Rhinella alata* MHUAA 8486. **C:** *Rhinella horribilis* MHUAA 10722. **D:** *Cochranella resplendens* MHUAA 9540. **E:** *Espadarana prosoblepon* MHUAA 8493. **F:** *Hyalinobatrachium aureoguttatum* MHUAA 7075. **G:** *Hyalinobatrachium fleischmanni* MHUAA 8176. **H:** *Rulyrana susatamai* MHUAA 10684.

which correspond to clumps of iridophores (absent in *H. valeroi*).

Hyalinobatrachium fleischmanni (Boettger, 1893):

Figure 2

Hylella fleischmanni Boettger 1893: 251.

Hyalinobatrachium fleischmanni — Ruiz-Carranza and Lynch 1991: 24.

Material examined. Table 1.

Rulyrana susatamai (Ruíz-Carranza & Lynch, 1995):

Figure 2H

Cochranella susatamai Ruíz-Carranza and Lynch 1995: 62.

Rulyrana susatamai — Guayasamin et al. 2009: 34.

Material examined. Table 1.

Sachatamia punctulata (Ruiz-Carranza & Lynch, 1995):

Figure 3A

Cochranella punctulata Ruiz-Carranza and Lynch 1995: 4.

Sachatamia punctulata — Guayasamin et al. 2009: 36.

Material examined. Table 1.

***Sachatamia* sp.:** Figure 3B

Material examined. Table 1.

Craugastor metriosistus Ospina-Sarria, Angarita-Sierra & Pedroza-Banda, 2015: Figure 3C

Craugastor metriosistus Ospina-Sarria et al. 2015: 165.

Material examined. Table 1.

A medium-sized frog characterized by having extensive toe webbing and a uniformly reddish brown coloration on the posterior surfaces of the thighs. *Craugastor metriosistus* differ from *C. longirostris* and *C. raniformis* (the latter also distributed in the Magdalena river valley) in having the throat with median white gular stripe and dark coloration on either side (throat with spotted pattern or short lines in *C. longirostris* and white to cream in *C. raniformis*), in having posterior surfaces of the thighs reddish brown (dark to reddish brown with small and conspicuous pale spots in *C. raniformis*), and in having webbing extending to the proximal portion of the distal subarticular tubercle on the outer side of toe III (webbing extends to middle portion between basal and distal subarticular tubercle on the outer side of toe III in *C. raniformis* and *C. longirostris*; Ospina-Sarria et al. 2015).

Pristimantis fallax (Lynch & Rueda-Almonacid, 1999):

Figure 3D

Eleutherodactylus fallax Lynch and Rueda-Almonacid 1999: 308.

Pristimantis fallax — Hedges et al. 2008: 118.

Material examined. Table 1.

Pristimantis fallax is a moderate size frog with a shagreen dorsum; finger I longer than finger II; toe V slightly longer than toe III; absence of toe webbing and broad disks on outer fingers. The most distinctive feature of *P. fallax* is the coloration pattern of the throat consisting in a lateral portion reticulated with gray to black, defining cream central raphe, a feature shared with *Craugastor fitzingeri*. Although superficially resembling *C. fitzingeri*, *P. fallax* is easily distinguished in having toe V slightly longer than toe III (toe III slightly longer than toe V in *C. fitzingeri*), absence of webbing on toes (basally webbed

in *C. fitzingeri*), and by having the posterior surfaces of the thighs unicolor (with pale spots in *C. fitzingeri*).

Pristimantis gaigei (Dunn, 1931): Figure 3E

Lithodytes gaigei Dun 1931: 387.

Pristimantis gaigeae — Hedges et al. 2008: 120.

Material examined. Table 1.

Pristimantis gaigei is a small to medium-sized frog characterized by having a distinct tympanum; annulus partially concealed dorsally and posteriorly by supra-tympanic fold; skin of dorsal surfaces shagreened; skin of venter smooth or feebly granular along lateral and posterior borders of ventral disk; discoidal folds and prominent arm slender (Lynch 1980). This species is easily distinguished from related species (i.e., *P. penelopus*) also distributed in the Magdalena river valley by having smooth and dark gray ventral surface, a bright yellow band on the forearm, and no calcar tubercle (in contrast with brown ventral surface with cream flecking, absence of yellow band and presence of calcar tubercle in *P. penelopus*).

Pristimantis jaguensis Rivera-Prieto, Rivera-Correa & Daza, 2014: Figure 3F

Pristimantis jaguensis Rivera-Prieto, Rivera-Correa & Daza 2014: 225.

Material examined. Table 1.

A moderated-sized frog characterized by a wide variation in color; absence of nuptial pads, discoidal fold and conical calcar tubercles; flanks and belly white to cream without blotches as well as iris yellow ocher to copper with thick brown reticulation and cream sclera (Rivera et al. 2014). This species co-occurs in the Magdalena river valley with *Pristimantis penelopus* and *P. viejas* and may be distinguished from both by having smooth skin and lacking a calcar tubercle (dorsum with tubercles including the heel in *P. penelopus* and *P. viejas*) and the concealed surfaces of the thighs yellow without blotches (orange spots on the concealed surfaces of the limbs in *P. viejas*; yellow spots on the concealed surfaces of the limbs, sometimes forming a reticle, in *P. penelopus*).

Pristimantis penelopus (Lynch & Rueda-Almonacid, 1999): Figure 3G

Eleutherodactylus penelopus Lynch and Rueda-Almonacid 1999: 310.

Pristimantis penelopus — Hedges et al. 2008: 128.

Material examined. Table 1.

A moderated-sized frog with a shagreen dorsum; tubercles posteriorly and on flanks; no dorsolateral folds; skin of venter brown with cream flecking; discoidal folds prominent; subanal tubercles only slightly larger than granules on underside of thighs; and iris copper with black reticulation. Males have spots on concealed surfaces of the thighs, a feature shared with *Pristimantis viejas*; nevertheless, these spots are usually yellow (sometimes forming a reticle) in *P. penelopus* and orange in *P. viejas*. *Pristimantis penelopus* is also similar to *P. latidiscus* from which it may distinguished in having a brown venter with cream flecking (in contrast to a cream venter with brown reticulation) (Lynch and Rueda-Almonacid 1999, Restrepo et al. 2017).




Figure 3. Amphibian species present in Jaguas and San Carlos hydroelectric protected areas. **A:** *Sachatamia punctulata* MHUAA 10689. **B:** *Sachatamia* sp. MHUAA 8205. **C:** *Craugastor metriosistus* MHUAA 10589. **D:** *Pristimantis fallax* MHUAA 10523. **E:** *Pristimantis gaigei* MHUAA 8497. **F:** *Pristimantis jaguensis*. **G:** *Pristimantis penelopus* MHUAA 8487. **H:** *Pristimantis taeniatus* MHUAA 8483.

Pristimantis taeniatus (Boulenger, 1912): Figure 3H

Eleutherodactylus taeniatus Boulenger 1912: 188.

Pristimantis taeniatus — Hedges et al. 2008: 121.

Material examined. Table 1.

Pristimantis viejas (Lynch & Rueda-Almonacid, 1999): Figure 4A

Eleutherodactylus viejas Lynch and Rueda-Almonacid 1999: 311.

Pristimantis viejas — Hedges et al. 2008: 121.

Material examined. Table 1.

A small-sized frog with numerous non-conical tubercles on dorsum; absence of dorsolateral folds; finger I shorter than finger II; toe V very long; brown above with darker brown markings; venter cream stippled with brown and posterior surfaces of the thighs brown with large cream spots. Superficially similar to *Pristimantis taeniatus* in lacking a canthal stripe, but readily distinguished from that species in having orange spots on the concealed surfaces of the limbs (absent in *P. taeniatus*; Lynch and Rueda-Almonacid 1999).

Colostethus inguinalis (Cope, 1868): Figure 4B

Prostherapis inguinalis Cope 1868: 10.

Colostethus inguinalis — Savage 1968: 745.

Material examined. Table 1.

A moderate-sized dendrobatid with a pale oblique lateral stripe extending from groin only to around midbody; pale ventrolateral stripes present; pale dorsolateral stripes absent and toes moderately webbed. *Colostethus inguinalis* may be distinguished from *C. aff. fraterdanieli* (also distributed in the Magdalena river valley) by having an incomplete pale oblique lateral stripe (complete in *C. aff. fraterdanieli*) and by the throat coloration of adults (black in adult males, white unpigmented or faintly pigmented gray or brown in adult females in *C. inguinalis*; spotted in adult males and immaculate in adult females in *C. aff. fraterdanieli*). It differs from *C. pratti* both in toe webbing (absent in *C. pratti*) and adult male throat coloration (pale gray or brown with irregular white spots, forming a faint, mottled or reticulated pattern in *C. pratti*).

Colostethus aff. fraterdanieli: Figure 4C

Material examined. Table 1.

A moderate-sized dendrobatid with dorsal skin texture granular posteriorly; pale dorsolateral stripe absent; pale oblique lateral stripe complete (extending from groin to eye); pale ventrolateral stripe present; coloration pattern of throat spotted in males, immaculate in females (Tolosa et al. 2014). *Colostethus aff. fraterdanieli* differs from *C. fraterdanieli* in having a pale ventrolateral stripe (absent in *C. fraterdanieli*) and a continuous dark band running along the lateral surfaces of body and head (discontinuous in *C. fraterdanieli*). It may be distinguished from *C. inguinalis* by having a complete oblique lateral stripe (partial in *C. inguinalis*) and from *C. pratti* by lacking pale dorsolateral stripe (present in *C. pratti*).

Dendrobates truncatus (Cope, 1861): Figure 4D

Phyllobates truncatus Cope 1861a: 372.

Dendrobates truncatus — Cope 1867: 197.

Material examined. Table 1.

Diasporus anthrax (Lynch, 2001): Figure 4E

Eleutherodactylus anthrax Lynch 2001: 292.

Diasporus anthrax — Hedges et al. 2008: 47.

Material examined. Table 1.

Small frog with skin of dorsum smooth; tympanum round and prominent; snout subacuminate in dorsal view, rounded in lateral view; all fingers and toes with expanded disks; pads triangular on toes III–IV; dorsum brown with darker blotches and venter dark brown with whitish flecks. *Diasporus anthrax* differs from other species of the genus by its coloration which consist in a dark venter with white flecks and pale blotches (red in life) on the concealed surfaces of the limb (orange above, venter immaculate, and lacking distinctive coloration marks on limbs in *D. gularis*; brown above, venter white with brown marbling, and lacking distinctive coloration marks on limbs in *D. tinkler*) and the rounded disk pads and disk covers except for finger III and toes III–IV in contrast with all fingers bearing narrow disks and disk cover of finger III bearing elongate papilla in *D. tinkler*; unornamented disk covers and knobbed disk cover occur in about equal frequency in *D. gularis* (Lynch 2001).

Diasporus gularis (Boulenger, 1898): Figure 4F

Hylodes gularis Boulenger 1898: 121.

Diasporus gularis — Hedges et al. 2008: 47.

Material examined. Table 1.

A small frog with short body; snout nearly truncate at tip when viewed from above, truncate in profile; nostrils strongly projecting laterally; fingers very short, with lateral ridges, pointed disks, and very small basal webbings; skin of upper surfaces smooth with a few short weak glandular lines above shoulders and along sides. The best diagnostic characters to recognize *Diasporus gularis* are its very short hands and feet with pointed fingers and toes, of which the entire terminal disk is dark gray, and the enormous U-shaped gular pouch of the males (Cochran and Goin 1970). *Diasporus gularis* is most readily distinguished from other *Diasporus* (i.e., *D. anthrax* and *D. tinkler*) by its coloration which is orange above with a transverse dark bar between eyes and immaculate venter (dorsum brown with darker blotches and venter dark brown with whitish flecks in *D. anthrax*; brown above with a tan interorbital bar, dorsolateral flank stripes and venter white with brown marbling in *D. tinkler*; Lynch 2001).

Hemiphractus fasciatus Peters, 1862: Figure 4G

Hemiphractus fasciatus Peters 1862: 169.

Boana boans (Linnaeus, 1758): Figure 4H

Rana boans Linnaeus 1758: 213.

Hypsiboas boans — Faivovich et al. 2005: 89.

Boana boans — Dubois 2017: 28.

Material examined. Table 1.

Boana xerophylla (Duméril & Bibron, 1841): Figure 5A

Hyla xerophylla Duméril and Bibron 1841: 549.

Hypsiboas xerophyllus — Orrico et al. 2017: 107.

Boana xerophylla — Dubois 2017: 28.

Material examined. Table 1.




Figure 4. Amphibian species present in Jaguas and San Carlos hydroelectric protected areas. **A:** *Pristimantis viejas* MHUAA 9742. **B:** *Colostethus inguinalis* MHUAA. 10645. **C:** *Colostethus* aff. *fraterdanieli* MHUAA 8140. **D:** *Dendrobates truncatus* MHUAA 9740. **E:** *Diasporus anthrax* MHUAA 7237. **F:** *Diasporus gularis* MHUAA 8496. **G:** *Hemiphractus fasciatus*. **H:** *Boana boans* MHUAA 10564.

This species was recently separated from *Boana crepitans* (Orrico et al. 2017). A large-sized tree frog having its dorsal surfaces pinkish tan with some darker brown spots, blotches or other markings; dark bars on the upper surfaces of the thighs extending onto the rear surfaces of the thighs; ventral surface orange, except for the throat and chest, which are white with some brown flecking. *Boana xerophylla* may be confused with *B. boans* and *B. pugnax* (species distributed in the Magdalena river valley) due to similarities in body size and coloration. However, it may be distinguished by its horizontal pupil and its iris that is pale gray at the center and greenish yellow at the edges (in contrast to a rhomboid pupil with an iris that is black at the center and light orange at the edges in *B. boans*, and a horizontal pupil with an iris that is black at the center and greenish in the upper portion in *B. pugnax*).

Dendropsophus ebraccatus (Cope, 1874): Figure 5B

Hyla ebraccata Cope 1874: 69.

Dendropsophus ebraccatus — Faivovich et al. 2005: 91.

Material examined. Table 1.

Dendropsophus microcephalus (Cope, 1886): Figure 5C

Hyla microcephala Cope 1886: 281.

Dendropsophus microcephalus — Faivovich et al. 2005: 92.

Dendropsophus norandinus Rivera-Correa & Guitérrez-Cárdenas, 2012: Figure 5D

Dendropsophus norandinus Rivera-Correa & Guitérrez-Cárdenas 2012: 51.

Material examined. Table 1.

A small hylid frog with a snout that is short, rounded in dorsal view and slightly rounded in lateral view; head wider than long; skin on dorsal surfaces smooth, with some minute scattered tubercles; axillary membrane present and developed; dorsum reddish brown in females and cream or silver gray in males (Rivera-Correa and Guitérrez-Cárdenas 2012). *Dendropsophus norandinus* may be distinguished from *D. bogerti* by having the head wider than long (longer than wide in *D. bogerti*) and by having finger webbing that extends to the base of the disks of toes I, II, and III (finger webbing absent in *D. bogerti*), from *D. subocularis* by lacking light bands connecting eye to edge of lips, and from *D. microcephalus* by its webbed fingers (fingers free in *D. microcephalus*).

Dendropsophus subocularis (Dunn, 1934): Figure 5E

Hyla subocularis Dunn 1934: 2.

Dendropsophus subocularis — Faivovich et al. 2005: 93.

Material examined. Table 1.

A small hylid frog with short snout; canthus rostralis very marked; fingers webbed at base about toes I–III; toes webbed nearly to disks of III and V; skin of dorsum smooth and skin of vent rugose (Dunn 1934). *Dendropsophus subocularis* co-occurs with *D. microcephalus* and *D. norandinus* in the Magdalena river valley. It may be distinguished from both by the presence of 2 light bands from the eye to the edge of the lips, enclosing a bar of dark between them (absent in the latter 2 species), and from *D. microcephalus* by its webbed fingers (fingers free in *D. microcephalus*).

Hyloscirtus palmeri (Boulenger, 1908): Figure 5F

Hyla palmeri Boulenger 1908: 515.

Hyloscirtus palmeri — Faivovich et al. 2005: 85.

Material examined. Table 1.

Phyllomedusa venusta Duellman & Trueb, 1967: Figure 5G

Phyllomedusa venusta Duellman and Trueb 1967: 128.

Material examined. Table 1.

***Scinax* sp.:** Figure 5H

Material examined. Table 1.

Our *Scinax* sp. corresponds to *Scinax* “A”, an undescribed species proposed by Nieto-Castro (1999) using morphological traits, morphometric and chromatic analyses. From that study, the name began to be used to refer to populations of *Scinax* distributed in Antioquia and Cundinamarca, characterized by having a golden color with a dorsal spotted pattern.

Scinax rostratus (Peters, 1863): Figure 6A

Hyla rostrata Peters 1863a: 466.

Scinax rostratus — Köhler and Böhme 1996: 139.

Material examined. Table 1.

A moderate-sized hylid frog characterized by a conspicuous acuminate and projecting snout in dorsal view, protruding in lateral view (Duellman 1970). *Scinax rostratus* is readily distinguished from *Scinax ruber* and *Smilisca sila* (both species occurring in the Magdalena river valley) by the combination of its characteristic snout and coloration pattern (surfaces of thighs yellow or pale orange with black or dark brown markings in *S. rostratus*; distinct black and white markings on posterior thighs in *S. ruber*; blue spots on flanks and posterior surfaces of the thighs in *S. sila*).

Scinax ruber (Laurenti, 1768): Figure 6B

Hyla rubra Laurenti 1768: 35.

Scinax ruber — Köhler and Böhme 1996: 139.

Material examined. Table 1.

A medium-sized frog with toes webbed except for the innermost; dorsal surfaces generally olive brown or orange-brown; distinct black and white markings on posterior surfaces of thighs. Related species which may be confused with *Scinax ruber* are *S. Boulengeri* and *S. rostratus* (also distributed in the Magdalena river valley) from both of these it differs by its smaller size and the presence of black markings on the thighs in form of vermiculations instead of broad and vertical dark bands (Cochran and Goin 1970). *Scinax ruber* is also similar to *Smilisca phaeota* from which it differs by lacking green coloration in the rostral region, and to *Smilisca sila* from which it differs by the absence of blue spots on the flanks and posterior surfaces of the thighs.

Smilisca phaeota (Cope, 1862): Figure 6C

Hyla phaeota Cope 1862: 358.

Smilisca phaeota — Starrett 1960: 303.

Material examined. Table 1.

Engystomops pustulosus (Cope, 1864): Figure 6D

Paludicola pustulosa Cope 1864: 180.

Engystomops pustulosus — Boulenger 1882: 276.




Figure 5. Amphibian species present in Jaguas and San Carlos hydroelectric protected areas. **A:** *Boana xerophyllus* MHUAA 10567. **B:** *Dendropsophus ebraccatus* MHUAA 10692. **C:** *Dendropsophus microcephalus* MHUAA 8667. **D:** *Dendropsophus norandinus* MHUAA 8589. **E:** *Dendropsophus subocularis* MHUAA 8490. **F:** *Hyloscirtus palmeri* MHUAA 9504. **G:** *Phyllomedusa venusta* MHUAA 10581. **H:** *Scinax* sp. MHUAA 8228.




Figure 6. Amphibian species present in Jaguas and San Carlos hydroelectric protected areas. **A:** *Scinax rostratus* MHUAA 7233. **B:** *Scinax ruber* MHUAA 9500. **C:** *Smilisca phaeota* MHUAA 9503. **D:** *Engystomops pustulosus* MHUAA 10602. **E:** *Leptodactylus colombiensis* MHUAA 8818. **F:** *Leptodactylus fuscus* MHUAA 10598. **G:** *Leptodactylus savagei* MHUAA 9768. **H:** *Bolitoglossa ramosi*.




Figure 7. Amphibian species present in Jaguas and San Carlos hydroelectric protected areas. **A:** *Caecilia thompsoni* MHUAA 7192. **B:** *Microcaecilia pricei* MHUA-A 10728. **C:** *Epicrionops parkeri* MHUA-A 10720.

Physalaemus pustulosus — Lynch 1970: 488.

Material examined. Table 1.

Leptodactylus colombiensis Heyer, 1994: Figure 6E

Leptodactylus colombiensis Heyer 1994: 82.

Material examined. Table 1.

Leptodactylus fuscus (Schneider, 1799): Figure 6F

Rana fusca Schneider 1799: 130.

Leptodactylus fuscus — Heyer 1968: 160.

Material examined. Table 1.

Leptodactylus savagei Heyer, 2005: Figure 6G

Leptodactylus savagei Heyer 2005: 330.

Bolitoglossa ramosi Brame & Wake, 1972: Figure 6H

Bolitoglossa ramosi Brame and Wake 1972: 9.

Material examined. Table 1.

Caecilia thompsoni Boulenger, 1902: Figure 7A

Caecilia thompsoni Boulenger 1902: 152.

Material examined. Table 1.

A large caecilian with generally cylindrical body; eye in an open orbit and tentacular aperture below nostril, close to the anterior portion of the face (Lynch 1999). *Caecilia thompsoni* is readily distinguished from *C. subnigricans* (another species known from Magdalena river valley) by having a longer body (up to 1.5 m in *C. thompsoni*; 0.35–0.37 m in *C. subnigricans*) and more primary (187–240 in *C. thompsoni*; 145–169 in *C. subnigricans*) and secondary annuli (26–30 in *C. thompsoni*; 6–27 in *C. subnigricans*)

Microcaecilia pricei (Dunn, 1944): Figure 7B

Gymnopsis pricei Dunn 1944: 473.

Microcaecilia pricei — Wilkinson, O'Connor and Nassbaum 2013: 7.

Material examined. Table 1.

Epicrionops parkeri (Dunn, 1942): Figure 7C

Rhinatrema parkeri Dunn 1942: 458.

Epicrionops parkeri — Taylor 1968: 216.

Material examined. Table 1.

A bicolored caecilian with well-developed tail; lateral stripe yellow, cream or white from head or second collar onto tail; stripe ending at posterior level of vent; subcaudal region dark lilac to blackish (Taylor 1968). *Epicrionops parkeri* differs from *E. bicolor* (a species with similar coloration pattern distributed in Colombia) by having a larger body length and 198 primary annuli (in contrast to 117–135 primary annuli in *E. bicolor*).

Reptiles

We report 58 reptile species comprising 47 genera in 16 families (Table 2). The protected area of Jaguas hydroelectric project has 38 species from 14 families, where the most speciose families were Dipsadidae and Colubridae with 10 and 8 species, respectively (26 and 21%). One species (*Anolis sulcifrons* Daudin, 1802) is endemic to Colombia and only 10 (26%) have been evaluated by the IUCN, 1 listed as Near Threatened and 9 as Least Concern (Table 4). In contrast, in San Carlos we found 57 reptile species comprising 16 families. Again, the most speciose families at this site were Dipsadidae with 15 species (26%) and Colubridae with 11 species (19%). Two species, *Anolis sulcifrons* Daudin, 1802 and *Ninia teresitae* Angarita-Sierra & Lynch, 2017 are endemic to

Colombia (4%) and 16 have been evaluated by the IUCN (2016), all of them categorized as Least Concern. Jaguas and San Carlos localities shared 36 reptile species.

Basiliscus galeritus Duméril, 1851: Figure 8A

Basiliscus galeritus Duméril 1851: 61.

Material examined. Table 2.

Adults of this large semiaquatic lizard are easily recognized by their gray to green color and the presence of a single, rounded head crest that is larger in males. Similar species that are found sympatric with *B. galeritus* are *Corytophanes cristatus* and *Basiliscus basiliscus*. The first is usually brownish and has 2 head crests that converge posteriorly. *Basiliscus basiliscus* has a brown or gray coloration and a single pointed head crest that is larger in males. Males of *B. basiliscus* also have a large crest in the back that resembles a fin (Maturana 1962; Savage 2002; Köhler 2008).

Corytophanes cristatus (Merrem, 1820): Figure 8B

Agama cristata Merrem 1820: 50.

Corytophanes cristatus — Boie in Schlegel 1827: 290.

Material examined. Table 2.

Anolis auratus Daudin, 1802: Figure 8C

Anolis auratus Daudin 1802: 89.

Material examined. Table 2.

Anolis fuscoauratus D'Orbigny, 1837

Anolis fuscoauratus D'Orbigny in Duméril and Bibron 1837: 110.

Material examined. Table 2.

Anolis* aff. *granuliceps: Figure 8D

Material examined. Table 2.

Anolis sulcifrons Cope, 1899: Figure 8E

Anolis sulcifrons Cope 1899: 6.

Material examined. Table 2.

Anolis tropidogaster Hallowell, 1956: Figure 8F

Anolis tropidogaster Hallowell 1956: 224.

Anolis vittigerus Cope, 1862

Anolis vittigerus Cope 1862: 179.

Material examined. Table 2.

Anadia* gr. *ocellata Gray, 1845: Figure 8G

Material examined. Table 2.

This small lizard may be distinguished from most other mid- to low-elevation gymnophthalmid lizards in having an acuminate snouth and elongate body in combination with smooth head and dorsal scales. Its color pattern includes a series of lateral white spots between

the fore and hind limbs (Oftedal 1974, Savage 2002). Usually, anterior most spots are bordered by black, forming ocelli. Although there is no evidence of *Anadia ocellata* for the Magdalena river valley, we assigned the specimen MHUAR 12694 to *Anadia* gr. *ocellata* because its dorsal coloration and 55 transversal rows in contrast to the 44–48 dorsal rows found in *A. rhombifera*, another species present in the Magdalena river valley (Oftedal 1974, Savage 2002).

***Cercosaura* sp.**: Figure 8H

Material examined. Table 2.

Echinosaura palmeri Boulenger, 1911: Figure 9A

Echinosaura palmeri Boulenger 1911: 23.

Material examined. Table 2.

Loxopholis rugiceps Cope, 1869

Loxopholis rugiceps Cope 1869: 305.

Material examined. Table 2.

Loxopholis southi (Ruthven & Gaige, 1924): Figure 9B

Leposoma southi Ruthven and Gaige 1924: 1.

Loxopholis southi — Goicoechea et al. 2016: 32.

Material examined. Table 2.

This species is easily differentiated from other gymnophthalmid lizards in the region by its strongly keeled head and dorsal scales and by having the ventral scales large, keeled, mucronate, and quadrangular to rectangular. It differs from *Loxopholis rugiceps*, the only other *Loxopholis* in trans-Andean Colombia, by having the frontonasal divided, instead of single (Köhler 2008).

***Ptychoglossus* sp.**: Figure 9C

Material examined. Table 2.

Thecadactylus rapicauda (Houttuyn, 1782): Figure 9D

Gekko rapicauda Houttuyn 1782: 323.

Thecadactylus rapicauda — Boulenger 1885: 111.

Material examined. Table 2.

Polychrus guttuerosus Berthold, 1846: Figure 9E

Polychrus guttuerosus Berthold 1846: 11.

This large blunt-headed lizard is immediately recognizable by its extremely long round tail that is at least twice the snouth–vent length, its robust body and relatively thin legs, and the presence of a large subtympnic scale (Peters and Donoso-Barros 1970, Savage 2002). In addition, it lacks toe pads, nuchal, dorsal and caudal crests (Peters and Donoso-Barros 1970, Savage 2002). It differs from *Polychrus marmoratus*, the other species that could be sympatric, because the latter has a row of

Table 2. Voucher list of reptiles of Jaguas and San Carlos hydroelectric compensation areas with geographic coordinates.

Taxon	Voucher	Latitude	Longitude	Taxon	Voucher	Latitude	Longitude
REPTILIA					MHUA-R 13006	6.29922	-74.96633
SQUAMATA					MHUA-R 13173	6.3515	-74.99528
SAURIA				<i>Corytophanes cristatus</i>	MHUA-R 12468	6.21854	-74.83727
Corytophanidae					MHUA-R 12429	6.24537	-74.86961
<i>Basiliscus galeritus</i>	MHUA-R 13010	6.277	-74.93104		MHUA-R 13015	6.28288	-74.92422
	MHUA-R 13009	6.2832	-74.92413		MHUA-R 13191	6.19012	-74.80477
	MHUA-R 13016	6.29256	-75.04247				

Continued

Table 2. Continued.

Taxon	Voucher	Latitude	Longitude	Taxon	Voucher	Latitude	Longitude
Dactyloidae					MHUA-R 13174	6.36704	-75.02715
<i>Anolis auratus</i>	MHUA-R 12432	6.2199	-74.84463	Amphisbaenidae			
	MHUA-R 13216	6.20643	-74.83891	<i>Amphisbaena varia</i>	MHUA-R 12276	6.19073	-74.80437
	MHUA-R 13217	6.20643	-74.83891	Colubridae			
	MHUA-R 13218	6.20643	-74.83891	<i>Chironius grandisquamis</i>	MHUA-R 15011	6.36728	-75.02697
<i>Anolis fuscoauratus</i>	MHUA-R 12433	6.21537	-74.83533	<i>Dendrophidion percarinatum</i>	MHUA-R 14764	6.19073	-74.80437
<i>Anolis aff. granuliceps</i>	MHUA-R 12726	6.19062	-74.80474	<i>Drymobius rhombifer</i>	MHUA-R 15012	6.39058	-75.03704
	MHUA-R 12792	6.19086	-74.80454	<i>Mastigodryas boddaerti</i>	MHUA-R 15054	6.19062	-74.80474
	MHUA-R 12514	6.1957	-74.8167		MHUA-R 15078	6.28296	-74.92384
	MHUA-R 12482	6.21768	-74.83748		MHUA-R 14866	6.22879	-74.87466
	MHUA-R 13013	6.32754	-75.01631	<i>Phrynonax poecilonotus</i>	MHUA-R 15013	6.19675	-74.80444
	MHUA-R 13011	6.32781	-75.01151	<i>Scaphiodontophis venustissimus</i>	MHUA-R 14871	6.2189	-74.8477
	MHUA-R 13014	6.32791	-75.01171	<i>Tantilla melanocephala</i>	MHUA-R 14855	6.21854	-74.83727
	MHUA-R 12725	6.36725	-75.02723		MHUA-R 14842	6.22148	-74.8418
	MHUA-R 12513	6.3693	-75.0252	Dipsadidae			
	MHUA-R 12445	6.37236	-75.04374	<i>Atractus occipitoalbus</i>	MHUA-R 15131	6.36767	-75.02635
	MHUA-R 12745	6.38528	-75.02158	<i>Coniophanes fissidens</i>	MHUA-R 14865	6.21691	-74.81272
	MHUA-R 12695	6.38871	-75.02335		MHUA-R 14879	6.2189	-74.8477
	MHUA-R 12746	6.39128	-75.0364		MHUA-R 14846	6.3852	-74.99314
	MHUA-R 12438	6.39677	-75.01162		MHUA-R 15133	6.20908	-74.85436
	MHUA-R 12436	6.40714	-75.02618	<i>Erythrolamprus epinephelus</i>	MHUA-R 14841	6.20874	-74.81525
	MHUA-R 12442	6.21611	-75.00078	<i>Erythrolamprus pseudocorallus</i>	MHUA-R 14826	6.349358	-75.002699
	MHUA-R 12441	6.37088	-75.04567		MHUA-R 14856	6.20716	-74.81999
	MHUA-R 12440	6.37187	-75.04636	<i>Imantodes cenchoa</i>	MHUA-R 14978	6.36725	-75.02723
	MHUA-R 12435	6.37588	-75.03902		MHUA-R 15134	6.36704	-75.02715
	MHUA-R 12019	6.8906	-75.5067	<i>Leptodeira septentrionalis</i>	MHUA-R 15077	6.32938	-75.01601
	MHUA-R 12393	7.393528	-75.036633		MHUA-R 14597	6.348145	-75.005172
	MHUA-R 13211	6.39912	-75.02728	<i>Ninia atrata</i>	MHUA-R 14847	6.2199	-74.84463
	MHUA-R 13226	6.1401	-75.06688		MHUA-A 15130	6.23974	-74.8688
	MHUA-R 13227	6.14229	-75.06267	<i>Ninia teresitae</i>	MHUA-R 14860	6.21626	-74.83435
<i>Anolis sulcifrons</i>	MHUA-R 12696	6.19055	-74.8048	<i>Oxyrhopus petolarius</i>	MHUA-R 14843	6.21146	-74.82359
	MHUA-R 12431	6.20801	-74.8157		MHUA-R 14693	6.8906	-75.5067
	MHUA-R 12434	6.21146	-74.82359		MHUA-R 15150	6.38927	-74.99146
	MHUA-R 12464	6.21872	-74.84776		MHUA-R 15151	6.25277	-74.82888
	MHUA-R 12443	6.24739	-74.86933	<i>Rhadinaea decorata</i>	MHUA-R 14859	6.2167	-74.86468
<i>Anolis vittigerus</i>	MHUA-R 12512	6.1957	-74.8167	<i>Sibon nebulatus</i>	MHUA-R 14857	6.20716	-74.81999
Gymnophthalmidae					MHUA-R 15139	6.38747	-74.99274
<i>Anadia gr. ocellata</i>	MHUA-R 12694	6.1957	-74.8167		MHUA-R 15149	6.38927	-74.99151
<i>Cercosaura sp.</i>	MHUA-R 12762	6.409987	-75.14157		MHUA-R 15154	6.15764	-75.0416
	MHUA-R 13221	6.287778	-74.828611	<i>Siphlophis compressus</i>	MHUA-R 14733	6.1738	-74.7513
<i>Echinosaura palmeri</i>	MHUA-R 13212	6.3995	-75.02823	<i>Urotheca fulviceps</i>	MHUA-R 14979	6.19062	-74.80474
	MHUA-R 13213	6.3994	-74.0284		MHUA-R 15055	6.19062	-74.80474
	MHUA-R 13223	6.15816	-75.04485		MHUA-R 15132	6.37516	-74.98231
	MHUA-R 13224	6.15813	-75.04476	Elapidae			
	MHUA-R 13225	6.1584	-75.04494	<i>Micrurus dumerilii</i>	MHUA-R 15137	6.19032	-74.80515
<i>Loxopholis rucigeps</i>	MHUA-R 12107	6.2247	-74.8477		MHUA-R 15138	6.19032	-74.80515
<i>Loxopholis southi</i>	MHUA-R 12430	6.20671	-74.855	<i>Micrurus mipartitus</i>	MHUA-R 14958	6.25141	-74.82854
	MHUA-R 12465	6.21841	-74.84771		MHUA-R 15152	6.19228	-74.80225
	MHUA-R 12105	6.2275	-74.8525	<i>Leptotyphlopidae</i>			
	MHUA-R 12466	6.22787	-74.872264	<i>Trilepida macrolepis</i>	MHUA-R 14858	6.21734	-74.81262
	MHUA-R 12106	6.2275	-74.8525	Viperidae			
<i>Ptychoglossus sp.</i>	MHUA-R 12602	6.36665	-75.02802	<i>Bothrops asper</i>	MHUA-R 14039	6.20453	-74.8949
Phyllodactylidae					MHUA-R 14691	6.8906	-75.5067
<i>Thecadactylus rapicauda</i>	MHUA-R 12601	6.19132	-74.80448	<i>Porthidium lansbergii</i>	MHUA-R 14054	6.20453	-74.8949
Sphaerodactylidae					MHUA-R 14516	6.224778	-74.847778
<i>Gonatodes albogularis</i>	MHUA-R 12423	6.231083	-74.876611		MHUA-R 15079	6.27689	-74.93108
<i>Lepidoblepharis xanthostigma</i>	MHUA-R 12467	6.2072	-74.81239		MHUA-R 14514	6.352361	-75.012139
	MHUA-R 12471	6.2167	-74.86468		MHUA-R 14515	6.354778	-75.008278
	MHUA-R 12422	6.231083	-74.876611		MHUA-R 14956	6.38553	-75.0218
	MHUA-R 13012	6.29719	-74.91953		MHUA-R 14694	6.511907	-74.910799
	MHUA-R 13008	6.29826	-74.91976		MHUA-R 14513	6.399989	-75.010364
	MHUA-R 12698	6.39202	-75.0365	TESTUDINES			
<i>Sphaerodactylus lineolatus</i>	MHUA-R 12470	6.18822	-74.8045	Kinosternidae			
	MHUA-R 12728	6.19062	-74.80474	<i>Kinosternon leucostomum</i>	MHUA-R 18526	6.38871	-75.02555
	MHUA-R 12469	6.21856	-74.83739		MHUA-R 18310	6.1897	-74.9969
Teiidae							
<i>Holcosus festivus</i>	MHUA-R 12444	6.21537	-74.83533				
	MHUA-R 12018	6.8906	-75.5067				




Figure 8. Reptile species present in Jaguas and San Carlos hydroelectric protected areas. **A:** *Basiliscus galeritus* MHUA-R 13009. **B:** *Corytophanes cristatus* MHUA-R 13015. **C:** *Anolis auratus* MHUA-R 13217. **D:** *Anolis* aff. *granuliceps* MHUA-R 13011. **E:** *Anolis sulcifrons* MHUA-R 12464. **F:** *Anolis tropidogaster*. **G:** *Anadia* gr. *ocellata* MHUA-R 12694. **H:** *Cercosaura* sp. MHUA-R 13162.




Figure 9. Reptile species present in Jaguas and San Carlos hydroelectric protected areas. **A:** *Echinosauro palmeri* MHUA-R 13158. **B:** *Loxopholis southi* MHUA-R 12466. **C:** *Ptychoglossus* sp. MHUA-R 12602. **D:** *Thecadactylus rapicauda* MHUA-R 12601. **E:** *Polychrus gutturosus*. **F:** *Marisora* sp. MHUA-R 13210. **G:** *Gonatodes albogularis* MHUA-R 12424. **H:** *Lepidoblepharis xanthostigma* MHUA-R 13012.




Figure 10. Reptile species present in Jaguas and San Carlos hydroelectric protected areas. **A:** *Sphaerodactylus lineolatus* MHUA-R 12470. **B:** *Ameiva* sp. **C:** *Cnemidophorus lemniscatus*. **D:** *Holcosus festivus* MHUA-R 13174. **E:** *Amphisbaena varia*. **F:** *Boa constrictor*. **G:** *Chironius exoletus*. **H:** *Chironius grandisquamis* MHUA-R 15011.

enlarged scales, forming a crest, in the mental and gular regions, which is absent in *P. guttuosus* (Peters and Donoso-Barros 1970).

Gonatodes albogularis (Duméril & Bibron, 1836):

Figure 9G

Gymnodactylus albogularis Duméril and Bibron 1836: 415.

Gonatodes albogularis — Boulenger 1885: 59.

Material examined. Table 2.

Lepidoblepharis xanthostigma (Noble, 1916): Figure 9H

Lathrogecko xanthostigma Noble 1916: 87.

Lepidoblepharis xanthostigma — Vanzolini 1953: 264.

Material examined. Table 2.

This species differs from other *Lepidoblepharis* species by having granular dorsals (cycloid in *L. sanctaemartae*), 12–14 lamellae under the fourth toe (<10 in *L. victormartinezi* and *L. emberawoundule*; Batista et al. 2015), median subcaudal scales more than twice as wide as the laterally adjacent subcaudals (less than 2 times in *L. ruficularis*; Batista et al. 2015).

Sphaerodactylus lineolatus Lichtenstein & von Martens 1856: Figure 10A

Sphaerodactylus lineolatus Lichtenstein & von Martens in Lichtenstein 1856: 6.

Material examined. Table 2.

Cnemidophorus lemniscatus (Linnaeus, 1758): Figure 10C

Lacerta lemniscata Linnaeus 1758: 209.

Cnemidophorus lemniscatus — Duméril and Bibron 1839: 129.

Holcosus festivus (Lichtenstein, 1856): Figure 10D

Cnemidophorus festivus Lichtenstein 1856: 13.

Holcosus festivus Harvey et al. 2012: 118.

Material examined. Table 2.

Amphisbaena varia Laurenti, 1768: Figure 10E

Amphisbaena varia Laurenti 1768: 66.

Material examined. Table 2.

This legless lizard is distinguished by its bullet-like head and rectangular body scales arranged in transversal rings (annuli). It has a whitish or pink ground with a large amount of irregular brown blotches, often obscuring ground color (Vanzolini 2002). *Amphisbaena varia* has traditionally been considered a subspecies of *A. fuliginosa* (Vanzolini 1951, 2002, Köhler 2008), but Gans (2005) elevated it to species level without further explanation.

Boa constrictor Linnaeus, 1758: Figure 10F

Boa constrictor Linnaeus 1758: 215.

Chironius exoletus (Linnaeus, 1758): Figure 10G

Coluber exoletus Linnaeus 1758: 223.

Chironius exoletus — Hoge, Romano and Cordeiro 1978: 41.

Chironius grandisquamis (Peters, 1869): Figure 10H

Spilotes grandisquamis Peters 1869: 451.

Chironius grandisquamis — Savage 2002: 650.

Material examined. Table 2.

This large species has a black or dark brown ground color and may have reddish or brown transversal bands (Dixon et al. 1993, Savage 2002). This species may be

separated from its sympatric congeners by having 10 rows of dorsal scales with the paravertebral scales keeled (12 rows in *Chironius exoletus* and *C. carinatus*), by having its dorsum black, or brown with transversal bands (dorsally green in *C. exoletus* and yellowish or olive ground color with 2 broad reddish brown stripes in *C. carinatus*).

Dendrophidion percarinatum (Cope, 1893): Figure 11A

Drymobius percarinatum Cope 1893: 344.

Dendrophidion percarinatum — Smith 1941: 73

Material examined. Table 2

Drymobius rhombifer (Günther, 1860): Figure 11B

Coryphodon rhombifer Günther 1860: 236.

Drymobius rhombifer — Boulenger 1894: 14.

Material examined. Table 2.

Leptophis ahaetulla (Linnaeus, 1758): Figure 11C

Coluber ahaetulla Linnaeus 1758: 225.

Leptophis ahaetulla — Bell 1825: 328.

Mastigodryas boddaerti (Sentzen, 1796): Figure 11D

Coluber boddaerti Sentzen 1796: 59.

Mastigodryas boddaerti — Montingelli et al. 2011: 189.

Material examined. Table 2.

Oxybelis aeneus (Wagler, 1824): Figure 11E

Dryinus aeneus Wagler in Spix 1824: 12.

Oxybelis aeneus — Duméril and Bibron 1854: 819.

Phrynonax poecilonotus (Günther, 1858): Figure 11F

Spilotes poecilonotus Günther 1858: 100.

Phrynonax poecilonotus — Boulenger 1894: 20.

Material examined. Table 2.

Adults have either red or yellow ground color, with a variable dorsal pattern that may consist of black transversal bands or several black dots, mainly located in the posterior region of the scales. Other than coloration, the species differs from other sympatric snakes by having 21–25 dorsal rows at midbody, with the uppermost (3–19) rows keeled (Savage 2002). Some authors recognize an additional species in trans-Andean Colombia, *Phrynonax shropshirei* (Taylor 1954, Peters et al. 1970, Pérez-Santos and Moreno 1988), but according to Savage (2002), it seems to be a confusion given the wide color variation in *P. poecilonotus*. Likewise, the profound genetic distinction between *P. poecilonotus* and *P. polylepis* (Peters, 1876) revealed by Jadin et al. (2014), who accordingly resurrected the latter species, does apparently not correspond to any morphological distinction. Since neither Jadin et al. (2014) nor the original description (Peters 1876) offer any morphological characters to discriminate between these 2 species, we must assume such distinctions to be obscured by the exuberant variability in *Phrynonax* for now. Unable to answer the interesting question to which species our specimen actually pertains, at this point we choose a conservative approach and tentatively assign it to *P. poecilonotus*.

Scaphiodontophis venustissimus (Günther, 1893):

Figure 11G

Henicognathus venustissimus Günther 1893: 144.

Scaphiodontophis venustissimus — Taylor and Smith 1943: 309.




Figure 11. Amphibian species present in Jaguas and San Carlos hydroelectric protected areas. **A:** *Dendrophidion percarinatum*. **B:** *Drymobius rhombifer* MHUA-R 15012. **C:** *Leptophis ahaetulla*. **D:** *Mastigodryas boddaerti* MHUA-R 15054. **E:** *Oxybelis aeneus*. **F:** *Phrynonax poecilonotus* MHUA-R 15013. **G:** *Scaphiodontophis venustissimus* MHUA-R 14871. **H:** *Spilotes pullatus*.

Material examined. Table 2.***Spilotes pullatus*** (Linnaeus, 1758): Figure 11H*Coluber pullatus* Linnaeus 1758: 225.*Spilotes pullatus* — Wagler 1830: 23.

This large snake reaches up to 2.5 m long. It differs from other snake species in the area by having a dorsal pattern of transversal yellow and black bands. Often, the posterior portion of the body is completely black (Savage 2002). It also differs from similar species in the region by having 14–18 dorsal rows of scales at the midbody with the uppermost keeled (Savage 2002).

Tantilla melanocephala (Linnaeus, 1758): Figure 12A*Coluber melanocephalus* Linnaeus 1758: 218.*Tantilla melanocephala* — Wilson and Mena 1980: 6.**Material examined.** Table Appendix 2.***Atractus occipitoalbus*** (Jan, 1862): Figure 12B*Rhabdosoma occipitulum* Jan 1862: 16.*Atractus occipitoalbus* — Boulenger 1894: 310.**Material examined.** Table 2.***Clelia clelia*** (Daudin, 1803): Figure 12C*Coluber clelia* Daudin 1803: 330.*Clelia clelia* — Zaher 1996: 301.***Coniophanes fissidens*** (Günther, 1858): Figure 12D*Coronella fissidens* Günther 1858: 36.*Coniophanes fissidens* — Pérez-Santos and Moreno 1988: 127.**Material examined.** Table 2.***Erythrolamprus epinephelus*** (Cope, 1862): Figure 12E*Liophis epinephalus* Cope 1862: 78.*Erythrolamprus epinephelus* — Grazziotin et al. 2012: 21.**Material examined.** Table 2.***Erythrolamprus melanotus*** (Shaw, 1802): Figure 12F*Coluber melanotus* Shaw 1802: 534.*Erythrolamprus melanotus* — Grazziotin et al. 2012: 21.

This medium-sized species may be recognized by having 17-17-15 smooth dorsal scale rows. Its dorsal ground color is light brown and yellowish, with 3 dorsal dark brown lines that extend from the neck to the tail. The 2 more lateral lines, running along the fourth and fifth dorsal rows, begin as series of points and the vertebral line one is about 5 scale rows wide (Dixon and Michaud 1992). Ventral coloration is white or light yellow (Dixon and Michaud 1992). The most similar sympatric species is *Erythrolamprus epinephelus*, which differs in having a grayish ground color with mainly transversal blotches or bands, 2 lateral dark lines that begin only in the posterior portion of the body extending to the tail, and frequently a reddish to yellowish venter with black blotches (Dixon 1983).

Erythrolamprus pseudocorallus Roze, 1959: Figure 12G*Erythrolamprus pseudocorallus* Roze 1959: 530.**Material examined.** Table 2.***Imantodes cenchoa*** (Linnaeus, 1758): Figure 12H*Coluber cenchoa* Linnaeus 1758: 226.**Material examined.** Table 2.***Leptodeira septentrionalis*** Kennicott, 1859: Figure 13A*Leptodeira septentrionalis* Kennicott in Baird 1859: 16.**Material examined.** Table 2.***Ninia atrata*** (Hallowell, 1845): Figure 13B*Coluber atratus* Hallowell 1845: 245.*Ninia atrata* — Cope 1860: 340.**Material examined.** Table 2.***Ninia tereseitae*** Angarita-Sierra & Lynch, 2017: Figure 13C*Ninia tereseitae* Angarita-Sierra and Lynch (2017): 478.**Material examined.** Table 2.***Oxyrhopus petolarius*** (Linnaeus, 1758): Figure 13D*Coluber petola* Linnaeus 1758: 225.*Oxyrhopus petolarius* — Duméril and Bibron 1854: 1033.**Material examined.** Table 2.***Rhadinaea decorata*** (Günther, 1858): Figure 13E*Coronella decorata* Günther 1858: 35.*Rhadinaea decorata* — Taylor 1949: 196.**Material examined.** Table 2.

This small, brown snake has a white, black-bordered line that goes from the postoculars through the fifth and sixth dorsal rows to the tip of the tail. The vertebral region is a little paler than below the lines. Specimens of *Rhadinaea decorata* often have a subpreocular scale and 110–134 ventral scales (Myers 1974). The most similar species in the region is *Urotheca fulviceps*, which has a reddish or purple head band, a pale line running along the first and second dorsal scale rows and more than 136 ventrals.

Sibon nebulatus (Linnaeus, 1758): Figure 13F*Coluber sibon* Linnaeus 1758: 222.*Sibon nebulatus* — Fitzinger 1826: 60.**Material examined.** Table 2.

This species has a pattern of gray, white and black bands. Gray bands are larger and with dispersed small black blotches, followed by a thin white band and then by a medium-sized black band (Savage 2002). Ventrally, ground is white with black blotches or bands. *Sibon nebulatus* has a large head and eyes like species of the genus *Imantodes*, but it can be easily recognized by its color pattern (Savage 2002).

Siphlophis compressus (Daudin, 1803): Figure 13G*Coluber compressus* Daudin 1803: 247.*Siphlophis compressus* — Zaher and Prudente 1999: 699.**Material examined.** Table 2.***Urotheca fulviceps*** (Cope, 1886): Figure 13H*Rhadinaea fulviceps* Cope 1886: 279.*Urotheca fulviceps* — Savage and Crother 1989: 343.**Material examined.** Table 2.***Xenodon rabdocephalus*** (Wied-Neuwied, 1824):

Figure 14A

Coluber rabdocephalus Wied-Neuwied 1824: 668.*Xenodon rabdocephalus* — Duméril et al. 1854: 758.***Micrurus dumerilii*** Jan, 1858: Figure 14B*Micrurus dumerilii* Jan 1858: 522.**Material examined.** Table 2.



Figure 12. Amphibian species present in Jaguas and San Carlos hydroelectric protected areas. **A:** *Tantilla melanocephala* MHUA-R 14855. **B:** *Atractus occipitoalbus* MHUA-R 15131. **C:** *Clelia clelia*. **D:** *Coniophanes fissidens* MHUA-R 14879. **E:** *Erythrolamprus epinephelus* MHUA-R 14841. **F:** *Erythrolamprus melanotus*. **G:** *Erythrolamprus pseudocorallus* MHUA-R 14856. **H:** *Imantodes cenchoa* MHUA-R 15134.



Figure 13. Amphibian species present in Jaguas and San Carlos hydroelectric protected areas. **A:** *Leptodeira septentrionalis*. **B:** *Ninia atrata* MHUA-R 14847. **C:** *Ninia teresitae*. MHUA-R 14860. **D:** *Oxyrhopus petolaris* MHUA-R 14843. **E:** *Rhadinaea decorata* MHUA-R 14859. **F:** *Sibon nebulatus* MHUA-R 15154. **G:** *Siphlophis compressus* MHUA-R 14733. **H:** *Urotheca fulviceps* MHUA-R 15132.

Micrurus mipartitus (Duméril, Bibron & Duméril, 1854):

Figure 14C

Elaps mipartitus Duméril et al. 1854: 1220.

Micrurus mipartitus — Amaral 1926: 66.

Material examined. Table 2.

Trilepida macrolepis (Peters, 1857): Figure 14D

Stenostoma macrolepis Peters 1857: 402.

Trilepida macrolepis — Hedges 2011: 63 [by implication].

Material examined. Table 2.

This is a very small species that may be differentiated from other snakes by having body scales of the same size (without enlarged ventral), arranged in 14 rows around body (Köhler 2008). It has 4 supralabial scales and 10 scale rows around the middle of the tail. Individuals are dark brown dorsally and lighter ventrally, with whitish scale borders (Pinto et al. 2010). *Epictia goudotii*, a species potentially sympatric with *Trilepida macrolepis*, has 2 supralabials, a darker color pattern with stripes, and a yellow rostral scale (Pinto et al. 2010).

Bothrops asper (Garman, 1883): Figure 14E

Trigonocephalus asper Garman 1883: 124.

Bothrops asper — Hoge 1966: 113.

Material examined. Table 2.

Porthidium lansbergii (Schlegel, 1841): Figure 14F

Trigonocephalus lansbergii Schlegel 1841: 1.

Porthidium lansbergii — McDiarmid et al. 1999: 320.

Material examined. Table 2.

Kinosternon leucostomum (Duméril, Bibron & Duméril, 1851): Figure 14G

Cinosternon leucostomum Duméril et al. 1851: 17.

Cinosternon postinguinale Cope 1887: 23.

Material examined. Table 2.

This small freshwater cryptodiran turtle is yellowish brown. It may be recognized by having a single carapace keel and the axillar and inguinal scutes usually separated, whereas *Kinosternon scorpioides* has 3 carapace keels and the axillar and inguinal scutes usually in contact (Köhler 2008).

Discussion

The 43 amphibian species recorded in this study represent 5.2% of the Colombian amphibian species (Frost 2016), and 15% of the species reported from the Cordillera Central in Colombia (IUCN 2016). The 58 reptile species represent 6% of the total Colombian reptiles species (Uetz and Hošek 2016), and 44% of the reptile fauna reported for the Cordillera Central. The Cordillera Central in Colombia is recognized for its high diversity in amphibians and reptiles (Lynch et al. 1997) and the 2 sites we studied represent a high proportion of this herpetofauna.

Fifteen (35%) of the amphibian species found during the monitoring are endemic to Colombia. Most of them are restricted to the eastern flank of the Cordillera Central: *Bolitoglossa ramosi*, *Dendropsophus norandinus*, *Diasporus anthrax*, *Epicrionops parkeri*,

Pristimantis fallax, *Pristimantis jaguensis*, *Pristimantis viejas*, *Rheobates pseudopalmaris*, *Rulyrana susatamai*, and *Sachatamia punctulata*. The distribution of the other endemic species extends to the Magdalena river valley (*Microcaecilia pricei*, *Caecilia thompsoni* and *Craugastor metriosistus*), the Chocoan Pacific lowland (*Colostethus inguinalis*; Grant and Lynch 2004) or the Caribbean lowland and foothills of the Sierra Nevada de Santa Marta (*Dendrobates truncatus*). Among the reptiles, *Anolis sulcifrons* was the only recorded species with a distribution restricted to Colombia (Uetz and Hošek 2016).

Five records are of particular interest in this study. *Pristimantis jaguensis* is an endemic frog from Colombia, with only 1 known locality, the surroundings of the Jaguas hydroelectric project (Rivera-Prieto et al. 2014). *Hyalinobatrachium aureoguttatum* is a frog previously recorded from the departments of Valle del Cauca, Risaralda, Chocó and Antioquia on the western flank of the Cordillera Occidental in Colombia, between 45 and 1570 m above sea level from the provinces of Esmeralda and Imbabura in Ecuador, and from the eastern cordilleras in southwestern Darién province in Panama (IUCN 2016). This is the first record of this species in the Magdalena river valley in Colombia. The presence of *Rhadinaea decorata* in Colombia was suggested by Myers (1974), and we present a voucher of this species from the San Carlos site, which thus represents the first verified record from Colombia. *Anadia ocellata* is distributed on both slopes of the cordilleras in Costa Rica and western Panama (Savage 2002). Batista et al. (2016) mention the occurrence in the Pacific lowlands of Colombia, contrary to Oftedal (1974), Savage (2002), and Köhler (2008), who indicated a southern range limit in Panama. We conservatively assign our voucher specimen from the Magdalena river valley to the *A. ocellata* group (sensu Oftedal 1974). The distribution of *Rhinella alata* in eastern cordillera was suggested by Pramuck (2006), although her observation lacked a voucher. Here, we present vouchers of *R. alata* that confirm the inter-Andean distribution of this species with records from Magdalena river valley.

Historically, the herpetofauna from the eastern side of the Andes has been considered very distinct from the trans-Andean region (Duellman 1979). Nevertheless, as part of the long-term sampling carried out in this study, we have recently reported on the occurrence of species distributed in the eastern flank of the northern Cordillera Central but previously known from the Amazon basin: *Atractus occipitoalbus* (Marin et al. 2017), *Anolis fuscoauratus* (Grisales-Martínez et al. in press) and *Cochranella resplendens* (Molina-Zuluaga et al. 2017). These findings urge the need for thorough sampling and a detailed taxonomic revision of species distributed in the Magdalena river valley to elucidate the biogeographic connection between both areas.

We were not able to assign 7 records to any valid species. The record of *Marisora* sp. (Figure 9F) may




Figure 14. Amphibian species present in Jaguas and San Carlos hydroelectric protected areas. **A:** *Xenodon rabdocephalus*. **B:** *Micrurus dumerillii* MHUA-R 15138. **C:** *Micrurus mipartitus* MHUA-R 15152. **D:** *Trilepida macrolepis* MHUA-R 14858. **E:** *Bothrops asper*. **F:** *Porthidium lansbergii* MHUA-R 14956. **G:** *Kinosternon leucostomum*.

correspond to *Marisora (Mabuya)* candidate species IV suggested by Pinto-Sánchez et al. (2015). The second record refers to an *Ameiva* sp. In Colombia, there are 3 *Ameiva* species: *Ameiva ameiva*, *A. bifrontata*, and *A. praesignis* (Harvey et al. 2012). As we only have a

specimen photograph (Figure 10B) we cannot assign it to any of these species. The vouchers of *Ptychoglossus* sp., *Sachatamia* sp., *Colostethus* aff. *fraterdanieli*, and *Cercosaura* sp. may represent undescribed taxa.

Pristimantis fallax is the only species reported in this

Table 3. Taxonomic list of amphibians reported in the Jaguas and San Carlos Hydroelectric protected areas. Asterisks indicate species endemic to Colombia. IUCN Conservation status: CR = Critically Endangered, EN = Endangered, VU = Vulnerable, NT = Near Threatened, LC = Least Concern, DD = Data Deficient; NE = Not Evaluated. Geographic distribution follows Lynch et al. (1997): A = Chocoan lowland forests, B = Caribbean lowlands and inter-Andean valleys, C = Llanos, D and E = Amazonia, F = Cordillera Occidental, G = Cordillera Central, H = Cordillera Oriental, I = Sierra Nevada de Santa Marta.

Taxon	Jaguas	San Carlos	IUCN	Distribution
AMPHIBIA				
ANURA				
Aromobatidae				
* <i>Rheobates pseudopalmaris</i> (Rivero & Serna, 2000)	X	X	DD	G
Bufoidea				
Bufoidea				
<i>Rhinella alata</i> (Thomiot, 1884)	X	X	DD	A
<i>Rhinella horribilis</i> (Wiegmann, 1833)	X	X	NE	A-B, F-I
Centrolenidae				
<i>Cochranella resplendens</i> (Lynch & Duellman, 1973)	X		LC	G, D-E,
<i>Espadarana prosoblepon</i> (Boettger, 1892)	X	X	LC	A-B, F-H
<i>Hyalinobatrachium aureoguttatum</i> (Barrera-Rodríguez & Ruiz-Carranza, 1989)		X	NT	A-B, G
<i>Hyalinobatrachium fleischmanni</i> (Boettger, 1893)		X	LC	A-B, F-H
* <i>Rulyrana susatamai</i> (Ruiz-Carranza & Lynch, 1995)	X		VU	B, G
* <i>Sachatamia punctulata</i> (Ruiz-Carranza & Lynch, 1995)	X	X	EN	B, G
<i>Sachatamia</i> sp.	X			
Craugastoridae				
* <i>Craugastor metriosistus</i> Ospina-Sarria, Angarita-Sierra & Pedroza-Banda, 2015	X	X	NE	B
* <i>Pristimantis fallax</i> (Lynch & Rueda, 1999)		X	VU	G
<i>Pristimantis gaigei</i> (Dunn, 1931)	X	X	LC	A-B, F-H
* <i>Pristimantis jaguensis</i> Rivera-Prieto, Rivera-Correa & Daza, 2014	X		NE	G
<i>Pristimantis penelopus</i> (Lynch & Rueda, 1999)	X	X	VU	B, G-H
<i>Pristimantis taeniatus</i> (Boulenger, 1912)	X	X	LC	A-B, F-H
* <i>Pristimantis viejas</i> (Lynch & Rueda-Almonacid, 1999)	X	X	LC	B, G-H
Dendrobatidae				
* <i>Colostethus inguinalis</i> (Cope, 1868)		X	LC	A-B, G
<i>Colostethus</i> aff. <i>fraterdanieli</i>	X			
* <i>Dendrobates truncatus</i> (Cope, 1861)	X	X	LC	A-B, G-I
Eleutherodactylidae				
* <i>Diasporus anthrax</i> (Lynch, 2001)	X	X	DD	B, G
<i>Diasporus gularis</i> (Boulenger, 1898)	X	X	LC	A-B, F-G
Hemiphractidae				
<i>Hemiphractus fasciatus</i> Peters, 1862	X		NT	A-B, F
Hylidae				
<i>Boana boans</i> (Linnaeus, 1758)	X	X	LC	A-I
<i>Boana xerophylla</i> (Duméril & Bibron, 1841)	X	X	NE	B-C, F-I
<i>Dendropsophus ebraccatus</i> (Cope, 1874)	X	X	LC	A-B, G-H
<i>Dendropsophus microcephalus</i> (Cope, 1886)	X	X	LC	A-C, G-I
* <i>Dendropsophus norandinus</i> Rivera-Correa & Gutiérrez-Cárdenas, 2012	X		NE	G
<i>Dendropsophus subocularis</i> (Dunn, 1934)	X	X	LC	A-B, F-H
<i>Hyloscirtus palmeri</i> (Boulenger, 1908)	X	X	LC	A-B, F-H
<i>Phyllomedusa venusta</i> Duellman & Trueb, 1967	X	X	LC	B, G-I
<i>Scinax</i> sp.	X	X		
<i>Scinax rostratus</i> (Peters, 1863)	X	X	LC	A-C, F-I
<i>Scinax ruber</i> (Laurenti, 1768)	X	X	LC	A-I
<i>Smilisca phaeota</i> (Cope, 1862)	X	X	LC	A-B, F-G
Leptodactylidae				
<i>Engystomops pustulosus</i> (Cope, 1864)	X	X	LC	A-C, G-I
<i>Leptodactylus colombiensis</i> Heyer, 1994	X	X	LC	B-C, F-H
<i>Leptodactylus fuscus</i> (Schneider, 1799)	X	X	LC	B-I
<i>Leptodactylus savagei</i> Heyer, 2005	X		LC	B,F-G, I
CAUDATA				
Plethodontidae				
* <i>Bolitoglossa ramosi</i> (Brame & Wake, 1972)	X		LC	G
GYMNOPHIONA				
Caeciliidae				
* <i>Caecilia thompsoni</i> (Boulenger, 1902)	X	X	DD	B, G-H
Siphonopidae				
* <i>Microcaecilia pricei</i>		X	LC	B,F
Rhinatrematidae				
* <i>Epicrionops parkeri</i> (Dunn, 1942)		X	DD	G

Table 4. Taxonomic list of reptiles reported in the Jaguas and San Carlos Hydroelectric protected areas. Asterisks indicate species endemic to Colombia. IUCN Conservation status is shown for both: a global assessment and a national assessment (IUCN 2016/Morales-Betancourt et al. 2015). CR= Critically Endangered, EN = Endangered, VU = Vulnerable, NT = Near Threatened, LC = Least Concern, DD = Data Deficient; NE = Not Evaluated. Geographic distribution follows Lynch et al. (1997): A = Chocoan lowland forests, B = Caribbean lowlands and inter-Andean valleys, C = Llanos, D and E = Amazonia, F = Cordillera Occidental, G = Cordillera Central, H = Cordillera Oriental, I = Sierra Nevada de Santa Marta.

Taxon	Jaguas	San Carlos	IUCN (Global/ National)	Distribution
REPTILIA				
SQUAMATA				
SAURIA				
Corytophanidae				
<i>Basiliscus galeritus</i> Dumeril, 1851	X	X	NE/LC	A–B, G
<i>Corytophanes cristatus</i> (Merrem, 1820)	X	X	LC/LC	A–B, G
Dactyloidae				
<i>Anolis auratus</i> Daudin, 1802		X	NE/LC	A–H
<i>Anolis</i> aff. <i>granuliceps</i>	X	X		
<i>Anolis fuscoauratus</i> D'Orbigny, 1837		X	NE/LC	
* <i>Anolis sulcifrons</i> Cope, 1899	X	X	NE/NT	B, G–H
<i>Anolis tropidogaster</i> Hallowell, 1856		X	NE/LC	
<i>Anolis vittigerus</i> Cope 1862		X	NE/LC	A–B, G–H
Gymnophthalmidae				
<i>Anadia</i> gr. <i>ocellata</i> Gray, 1845		X	NE/NT	A–B, G
<i>Cercosaura</i> sp.	X	X		
<i>Echinosaura palmeri</i> Boulenger, 1911	X	X	LC/NE	A, G
<i>Loxopholis rugiceps</i> Cope, 1869		X	LC/LC	A–B, G–I
<i>Loxopholis southi</i> Ruthven & Gaige, 1924		X	LC/LC	A, G
<i>Ptychoglossus</i> sp.	X	X		
Phyllodactylidae				
<i>Thecadactylus rapicauda</i> (Houttuyn, 1782)	X	X	NE/LC	A–C, E, G–H
Polychrotidae				
<i>Polychrus gutturosus</i> Berthold, 1846	X	X	NE/LC	A–B, F–H
Scincidae				
<i>Marisora</i> sp.	X	X		
Sphaerodactylidae				
<i>Gonatodes albogularis</i> (Duméril & Bibron, 1836)	X	X	NE/LC	A–I
<i>Lepidoblepharis xanthostigma</i> (Noble, 1916)	X	X	LC/LC	A–B, G
<i>Sphaerodactylus lineolatus</i> Lichtenstein & Von, 1856		X	LC/LC	A–B, G
Teiidae				
<i>Ameiva</i> sp.	X	X		
<i>Cnemidophorus lemniscatus</i> (Linnaeus, 1758)	X	X	NE/NE	A–B, D–E, G
<i>Holcosus festivus</i> (Lichtenstein, 1856)	X	X	LC/LC	A–B, G–H
AMPHISBAENIA				
Amphisbaenidae				
<i>Amphisbaena varia</i> Laurenti, 1768	X	X	NE/LC	A–H
OPHIDIA				
Boidae				
<i>Boa constrictor</i> Linnaeus, 1758	X	X	NE/LC	A–C, E, G–H
Colubridae				
<i>Chironius exoletus</i> (Linnaeus, 1758)	X	X	NE/LC	A–H
<i>Chironius grandisquamis</i> (Peters, 1869)	X	X	LC/LC	A–B, F–G
<i>Dendrophidion percarinatum</i> (Cope, 1893)	X	X	LC/LC	A–B, F–G
<i>Drymobius rhombifer</i> (Günther, 1860)		X	LC/LC	A–I
<i>Leptophis ahaetulla</i> (Linnaeus, 1758)	X	X	NE/LC	A–I
<i>Mastigodryas boddaerti</i> (Sentzen, 1796)	X	X	NE/LC	A–I
<i>Oxybelis aeneus</i> (Wagler, 1824)	X	X	NE/LC	A–H
<i>Phrynonax poecilonotus</i> (Günther, 1858)	X	X	LC/NE	A–B, F–H
<i>Scaphiodontophis venustissimus</i> (Gunther, 1893)		X	LC/NE	G
<i>Spilotes pullatus</i> (Linnaeus, 1758)	X	X	NE/LC	A–I
<i>Tantilla melanocephala</i> (Linnaeus, 1758)		X	NE/LC	A–C, E–H
Dipsadidae				
<i>Atractus occipitoalbus</i> (Jan, 1862)	X		NT/NT	E, G
<i>Clelia clelia</i> (Daudin, 1803)	X	X	NE/LC	A–I
<i>Coniophanes fissidens</i> (Gunther, 1858)	X	X	NE/LC	B, F–G
<i>Erythrolamprus epinephelus</i> (Cope, 1862)		X	NE/LC	F–I
<i>Erythrolamprus melanotus</i> (Shaw, 1802)		X	LC/LC	A–I
<i>Erythrolamprus pseudocorallus</i> Roze, 1959	X	X	LC/LC	B, G–H
<i>Imantodes cenchoa</i> Linnaeus, 1758	X	X	NE/LC	A–H

Continued

Table 4. Continued.

Taxon	Jaguas	San Carlos	IUCN (Global/ National)	Distribution
<i>Leptodeira septentrionalis</i> Kennicott, 1859	X	X	NE/LC	A–B, F–H
<i>Ninia atrata</i> (Hallowell, 1845)	X	X	LC/LC	B–C, E–H
* <i>Ninia teresitae</i> Angarita-Sierra & Lynch, 2017		X	NE/NE	B
<i>Oxyrhopus petolarius</i> (Linnaeus, 1758)	X	X	NE/LC	A–H
<i>Rhadinaea decorata</i> (Gunther, 1858)	X	X	NE/NE	G
<i>Sibon nebulatus</i> (Linnaeus, 1758)	X	X	NE/LC	A–C, F–I
<i>Siphlophis compressus</i> (Daudin, 1803)		X	LC/LC	A–E, G
<i>Urotheca fulviceps</i> (Cope, 1886)		X	NE/DD	A–B, H
<i>Xenodon rabdocephalus</i> (Wied-Neuwied, 1824)		X	NE/LC	A–B, E–H
Elapidae				
<i>Micrurus dumerilii</i> Jan, 1858		X	NE/LC	A–B, F–I
<i>Micrurus mipartitus</i> (Duméril, Bribon & Duméril, 1854)		X	NE/LC	A–B, F–I
Leptotyphlopidae				
<i>Trilepida macrolepis</i> (Peters, 1857)		X	NE/LC	A–B, D–E, G–H
Viperidae				
<i>Bothrops asper</i> (Garman, 1883)	X	X	NE/LC	A–B, G–H
<i>Porthidium lansbergii</i> (Schlegel, 1841)	X	X	NE/LC	A–B, F–H
TESTUDINES				
Kinosternidae				
<i>Kinosternon leucostomum</i> (Duméril, Bibron & Duméril, 1851)	X	X	NE/LC	A–B, G–H

study included as Vulnerable at the country level in the Resolución 192 del 2014 del Ministerio de Ambiente y Desarrollo Sostenible. At the global level, *Sachatamia punctulata* is listed by the IUCN as Endangered because its extent of occurrence is less than 5000 km², its distribution is severely fragmented, and there is a continuing decline in the extent and quality of its habitat (IUCN 2016). *Pristimantis fallax*, *P. penelopus* and *Rulyrana susatamai* are listed as Vulnerable because of continuous decline in the extent and quality of their habitat (Castro et al. 2004, IUCN 2016). Lastly, 2 species are categorized as Near Threatened: *Hemiphractus fasciatus* Peters, 1862 and *Hyalinobatrachium aureoguttatum* (Coloma et al. 2008, Solís et al. 2010; Table 3).

Recently, the conservation status of Colombia's reptiles has been assessed and 43 species are now under one of the 3 threat categories (Morales-Betancourt et al. 2015). However, none of them correspond to species present in our study areas. It is well known that reptiles around the world are declining due to a number of causes directly related to human activities (Gibbons et al. 2000). In addition to habitat loss and degradation, another threat is present in Colombia: the cultural view that reptiles, in particular snakes, are dangerous and nasty animals that should be exterminated (Castaño-Mora 2002, Lynch 2012). This view has probably driven several snake species to the brink of extinction at least locally (Lynch 2012).

To build complete species lists for Colombian amphibians and reptiles evokes large challenges for the following reasons: (1) the high levels of species diversity of amphibians and reptiles in the area, (2) the extreme cryptic diversity in several genera (e.g., *Pristimantis*, *Anolis*, *Atractus*), and (3) the lack of well-identified reference collections. However, continued specimen collection throughout different seasons and years in a specific local-

ity will give us a better picture of the regional diversity and will impact future studies aiming to understand the origin and maintenance of this rich fauna.

The high species density, high representativeness of the regional biodiversity, and the high levels of amphibian endemism make these 2 private protected areas very important for conservation of the regional herpetofauna. As shown here, compensation areas of large infrastructure projects (i.e., hydroelectric projects) represent a potential opportunity to preserve the already highly degraded ecosystems in the northern Andes of Colombia.

Acknowledgements

During almost a decade, ISAGEN and the Universidad de Antioquia have funded herpetofaunal studies in this region. This manuscript was partially funded under Isagen project 47/574. We thank the Grupo Herpetológico de Antioquia and all the herpetologists and students who over the years have been collecting information about the amphibian and reptile diversity in this highly diverse region of the Andes. M. Rivera-Correa provided useful comments to the manuscript.

Authors' Contributions

AR, CMZ and JMD conceived the idea and wrote the text. JMD, CMM and JPH confirmed the taxonomic identity of all specimens.

References

- Acevedo AA, Lampo M, Cipriani R (2016) The cane or marine toad, *Rhinella marina* (Anura, Bufonidae): two genetically and morphologically distinct species. *Zootaxa* 4103 (6): 574–586. <http://doi.org/10.11646/zootaxa.4103.6.7>
- Angarita-Sierra T, Lynch JD (2017) A new species of *Ninia* (Serpentes)

- tes: Dipsadidae) from Chocó-Magdalena biogeographical province, western Colombia. *Zootaxa* 4244 (4): 478–492. <http://doi.org/10.11646/zootaxa.4244.4.2>
- Amaral AD (1926) Studies on Neotropical ophidia. II. On *Micrurus mipartius* and allied forms. Proceedings of the New England Zoological Club 9: 61–66.
- Baird SF (1859) Reptiles of the boundary. In: United States and Mexican Boundary Survey under the Order of Lieut. Col. W. H. Emory, Major First Cavalry, and United States Commissioner. 2, Rept, Pt 2. Department of the Interior, Washington DC, 35 pp.
- Barrera-Rodríguez M, Ruíz-Carranza PM (1989) Una nueva especie del género *Centrolenella* Noble, 1920 (Amphibia: Anura; Centrolenidae) de la Cordillera Occidental de Colombia. *Trianea* 3: 77–84.
- Batista A, García-Rodríguez A, Saborío G, Vargas-Álvarez V, Acosta-Chaves V, Arredondo JC (2016) *Anadia ocellata*. The IUCN Red List of Threatened Species 2016. <https://doi.org/10.2305/iucn.uk.2016-1.rlts.t203048A2759434.en>. Accessed on: 2016-10-1.
- Batista A, Ponce M, Vesely M, Mebert K, Hertz A, Köhler G, Carrizo A, Lotzkat S (2015) Revision of the genus *Lepidoblepharis* (Reptilia: Squamata: Sphaerodactylidae) in Central America, with the description of three new species. *Zootaxa* 3994 (2): 187–221. <https://doi.org/10.11646/zootaxa.3994.2.2>
- Bell T (1825) On Leptophina, a group of serpents comprising the genus *Dyinus* of Merrem, and a newly formed genus proposed to be named *Leptophis*. *Zoological Journal* 2: 322–329, pl. 12.
- Berthold AA (1846) Über verschiedene neue order seltene Reptilien aus Neu-Granada und Crustaceen aus China. *Abhandlungen der Königlichen Gesellschaft der Wissenschaften zu Göttingen* 3: 3–32.
- Boettger O (1892) Katalog der Batrachier-Sammlung im Museum der Senckenbergischen Naturforschenden Gessellschaft in Frankfurt am Main. Frankfurt a. M. Gebrüder-Knauer, Frankfurt am Main, 73 pp.
- Boettger O (1893) Ein neuer Laubfrosch aus Costa Rica. Bericht der Senckenbergischen Naturforschenden Gesellschaft in Frankfurt am Main 1893: 251–252.
- Boulenger GA (1882) Catalogue of the Batrachia Salientias. Ecaudata in the Collection of the British Museum. 2nd ed. Taylor and Francis, London, 588 pp. <https://doi.org/10.5962/bhl.title.20903>
- Boulenger GA (1894) Catalogue of the Snakes in the British Museum (Natural History). Volume II, Containing the Conclusion of the Colubridæ Aglyphæ. British Museum (Natural History), London, xi, 382 pp.
- Boulenger GA (1911) Descriptions of new reptiles from the Andes of South America, preserved in the British Museum. *Annals and Magazine of Natural History*, 8th series 7: 19–25.
- Boulenger GA (1885) Catalogue of the Lizards in the British Museum (Natural History). Vol. 1. Geckonidae, Eublepharidae, Uroplatidae, Pygopodidae, Agamidae. British Museum (Natural History), London, 450 pp. <https://doi.org/10.5962/bhl.title.21097>
- Boulenger GA (1894) Catalogue of the Snakes in the British Museum (Natural History). Vol. 2. Containing the Conclusion of the Colubridæ Aglyphæ. British Museum (Natural History), London, xi, 382 pp. <https://doi.org/10.5962/bhl.title.21097>
- Boulenger GA (1898) An account of the reptiles and batrachians collected by Mr. W.F.H. Rosenberg in Western Ecuador. Proceedings of the Zoological Society of London 1898: 107–126.
- Boulenger GA (1902) Descriptions of two new South-American apodal batrachians. *Annals and Magazine of Natural History*, 7th series 10: 152–153.
- Boulenger GA (1908) Descriptions of new batrachians and reptiles discovered by Mr. M.G. Palmer in south-western Colombia. *Annals and Magazine of Natural History*, 8th series 2: 515–522.
- Boulenger GA (1912) Descriptions of new batrachians from the Andes of South America, preserved in the British Museum. *Annals and Magazine of Natural History*, 8th series 10: 185–191.
- Brame AH, Wake DB (1972) New species of salamanders (genus *Bolitoglossa*) from Colombia, Ecuador, and Panama. *Contributions in Science, Natural History Museum of Los Angeles County* 219: 1–34.
- Burbano-Yandi CE, Gómez-Días MA, Figueroa AG, Velásquez-Tujillo DA, Bolívar-García W (2016) Amphibians assemblage present in a tropical dry forest and in production systems, in Middle Magdalena Valley, Victoria and La Dorada, Caldas, Colombia. *Revista de Ciencias* 20 (2): 81–93.
- Castaño-Mora OV (Ed.) (2002) Libro rojo de reptiles de Colombia. Instituto de Ciencias Naturales/Universidad Nacional de Colombia/Ministerio del Medio Ambiente/Conservación Internacional, Bogotá, 166 pp.
- Castro F, Herrera I, Lynch JD (2004) *Pristimantis penelopus*. The IUCN Red List of Threatened species. Version 2014.2. <http://www.iucnredlist.org>. Accessed on: 2014-9-1.
- Cochran DM, Goin DJ (1970) Frogs of Colombia. *Bulletin of the United States National Museum* 288: 1–655. <https://doi.org/10.5479/si.03629236.288.1>
- Coloma LA, Ron SR, Jungfer K, Solís F, Ibáñez R, Jaramillo C, Fuenmayor Q, Bolaños F (2008) *Hemiphysalis fasciatus*. The IUCN Red List of Threatened species. Version 2014.2. <http://www.iucnredlist.org>. Accessed on: 2014-9-1.
- Cope ED (1861a) Descriptions of reptiles from tropical America and Asia. *Proceedings of the Academy of Natural Sciences of Philadelphia* 12: 368–374.
- Cope ED (1861b) Notes and descriptions of new and little-known species of American reptiles. *Proceedings of the Academy of Natural Sciences of Philadelphia* 12: 339–345 [1860].
- Cope ED (1861c) Catalogue of the colubrids in the museum of the Academy of Natural Sciences of Philadelphia. *Proceedings of the Academy of Natural Sciences of Philadelphia* 12: 368–374.
- Cope ED (1862) Catalogues of the reptiles obtained during the explorations of the Parana, Paraguay, Vermeio and Uruguay Rivers, by Capt. Thos. J. Page, U. S. N.; and of those procured by Lieut. N. Michler, U. S. Top. Eng., Commander of the expedition conducting the survey of the Atrato River. *Proceedings of the Academy of Natural Sciences of Philadelphia* 14: 346–359.
- Cope ED (1864) Contributions to the herpetology of tropical America. *Proceedings of the Academy of Natural Sciences of Philadelphia* 16: 166–181.
- Cope ED (1867) On the families of the raniform Anura. *Journal of the Academy of Natural Sciences of Philadelphia*, 2nd series 6: 189–206.
- Cope ED (1868) An examination of the Reptilia and Batrachia obtained by the Orton Expedition to Equador and the upper Amazon, with notes on others species. *Proceedings of the Academy of Natural Sciences of Philadelphia* 20: 96–140.
- Cope ED (1869) Sixth contribution to the herpetology of tropical America. *Proceedings of the Academy of Natural Sciences of Philadelphia* 20: 305–313 [1868].
- Cope ED (1874) Description of some species of reptiles obtained by Dr John F. Bransford, Assistant Surgeon United States Navy, while attached to the Nicaraguan Surveying Expedition in 1873. *Proceedings of the Academy of Natural Sciences of Philadelphia* 26: 64–72.
- Cope ED (1886) Thirteenth contribution to the herpetology of tropical America. *Proceedings of the American Philosophical Society* 23: 271–287.
- Cope ED (1887) Catalogue of batrachians and reptiles of Central America and México. *Bulletin of the United States National Museum* 32: 1–98. <https://doi.org/10.5479/si.03629236.32>
- Cope ED (1893) Second addition to the knowledge of the Batrachia and Reptilia of Costa Rica. *Proceedings of the American Philosophical Society* 31: 333–347.
- Cope ED (1899) Contributions to the herpetology of New Granada and Argentina, with descriptions of new forms. *Philadelphia Museums Science Bulletin* 1: 1–19.
- Cortez C, Suárez-Mayorga AM, López-López FJ (2006) Preparación y preservación de material científico. In: Angulo A, Rueda-Almonacid JV, Rodríguez-Mahecha JV, La Marca E (Eds) *Técnicas de Inventario y Monitoreo para los Anfibios de la Región Tropical*

- Andina. Conservación Internacional, Bogotá, 173–220.
- Daudin FM (1802) Histoire Naturelle, Générale et Particulière des Reptiles, Vol 4. F. Dufart, Paris, 426 pp. <https://doi.org/10.5962/bhl.title.60678>
- Daudin FM (1803) Histoire Naturelle Generale et Particuliere des Reptiles. Vol. 6. F. Dufart, Paris, 426 pp. <https://doi.org/10.5962/bhl.title.60678>
- Dubois A (2017) The nomenclatural status of *Hysaplesia*, *Hylaplesia*, *Dendrobates* and related nomina (Amphibia, Anura), with general comments on zoological nomenclature and its governance, as well as on taxonomic databases and websites. *Bionomina* 11: 1–48. <https://doi.org/10.11646/bionomina.11.1.1>
- Dixon JR (1983) A first record for an Amazonian snake similar to the Eastern Brazilian *Liophis*. *Copeia* 1983 (3): 791–802. <https://doi.org/10.2307/1441003>
- Dixon JR, Wiest JA Jr, Cei JM (1993) Revision of the Neotropical snake genus *Chironius* Fitzinger (Serpentes, Colubridae). *Museo Regionale di Scienze Naturali Monografie* (Turin) 13: 1–280.
- Dixon JR, Michaud EJ (1992) Shaw's black-backed snake (*Liophis melanotus*) (Serpentes: Colubridae) of northern South America. *Journal of Herpetology* 26 (3): 250–259. <https://doi.org/10.2307/1564878>
- dos Santos, PS, Ibáñez R, Ron RS (2015) Systematics of the *Rhinella margaritifera* complex (Anura, Bufonidae) from western Ecuador and Panama with insights in the biogeography of *Rhinella alata*. *ZooKeys* 501: 109–145. <https://doi.org/10.3897/zookeys.501.8604>
- Duellman WE (1970) The hylid frogs of Middle America. Monograph of the Museum of Natural History, the University of Kansas 1: 1–753. <https://doi.org/10.5962/bhl.title.2835>
- Duellman WE (1979) The South American herpetofauna: its origin, evolution, and dispersal. Monograph of the Museum of Natural History, the University of Kansas. No. 7. Museum of Natural History, University of Kansas, Lawrence, Kansas, 485 pp. <https://doi.org/10.5962/bhl.title.3207>
- Duellman WE (1999) Patterns and distribution of amphibians: a global perspective. The Johns Hopkins University Press, Baltimore, 633 pp.
- Duellman WE, Trueb L (1967) Two new species of tree frogs (genus *Phyllomedusa*) from Panamá. *Copeia* 1967: 125–131. <https://doi.org/10.2307/1442184>
- Duméril AMC, Bibron G (1836) Erpétologie Générale ou Histoire Naturelle Complete des Reptiles. Vol. 3. Libr. Encyclopédique Roret, Paris, 528 pp. <https://doi.org/10.5962/bhl.title.45973>
- Duméril AMC, Bibron G (1837) Erpétologie Générale ou Histoire Naturelle Complete des Reptiles. Vol. 4. Libr. Encyclopédique Roret, Paris, 570 pp.
- Duméril AMC, Bibron G (1839) Erpétologie Générale on Histoire Naturelle Complète des Reptiles. Vol. 5. Roret/Fain et Thunot, Paris, 871 pp.
- Duméril AMC, Bibron G (1841) Erpétologie générale ou Histoire naturelle complète des reptiles. Vol. 8. Librarie Encyclope de Roret, Paris, 818 pp..
- Duméril AMC, Bibron G, Duméril AHA (1854) Erpétologie générale ou Histoire Naturelle complete des Reptiles. Vol. 7. Librarie Encyclope de Roret, Paris, 488 pp.
- Duméril AMC, Duméril AHA (1851) Catalogue Méthodique de la Collection des Reptiles du Muséum d'Histoire Naturelle de Paris. Gide et Baudry/Roret, Paris, 224 pp.
- Dunn ER (1931) New frogs from Panama and Costa Rica. Occasional Papers of the Boston Society of Natural History 5: 385–401.
- Dunn ER (1934) Two new frogs from Darien. *American Museum Novitates* 747: 1–2.
- Dunn ER (1942) The American caecilians. *Bulletin of the Museum of Comparative Zoology, Harvard University* 91: 439–540.
- Dunn ER (1944) Notes in Colombian herpetology, III. *Caldasia* 2: 473–474.
- Etter A, van Wyngaarden W (2000) Patterns of landscape transformation in Colombia, with emphasis in the Andean region. *AMBIO: A Journal of the Human Environment* 29 (7): 432–439. <https://doi.org/10.1579/0044-7447-29.7.432>
- Faivovich J, Haddad CFB, Garcia PCA, Frost DR, Campbell JA, Wheeler WC (2005) Systematic review of the frog family Hylidae, with special reference to Hylinae: a phylogenetic analysis and taxonomic revision. *Bulletin of the American Museum of Natural History* 294: 1–240.
- Fitzinger LJ (1826) Neue Classification der Reptilien nach ihren Natürlichen Verwandtschaften nebst einer Verwandtschafts-Tafel und einem Verzeichnisse der Reptilien-Sammlung des K. K. Zoologisch Museums zu Wien. J. G. Heubner, Vienna, 88 pp. <https://doi.org/10.5962/bhl.title.4683>
- Fjeldsa J (1994) Geographical patterns for relict and young species of birds in Africa and South America and implications for conservation priorities. *Biodiversity and Conservation* 3 (3): 207–226. <https://doi.org/10.1007/BF00055939>
- Fjeldsa J, Álvarez MD, Lazcano JM, León B (2005) Illicit crops and armed conflict as constraints on biodiversity conservation in the Andes region. *AMBIO: A Journal of Human Environment* 34 (3): 205–211. <https://doi.org/10.1579/0044-7447-34.3.205>
- Frost DR, Grant T, Faivovich J, Bain RH, Haas A, Haddad CFB, de Sá RO, Channing A, Wilkinson M, Donnellan SC, Raxworthy CJ, Campbell JA, Blotto BL, Moler PE, Drewes RC, Nussbaum RA, Lynch JD, Green DM, Wheeler WC (2006) The amphibian tree of life. *Bulletin of the American Museum of Natural History* 297: 1–370. [https://doi.org/10.1206/0003-0090\(2006\)297\[0001:TATOL\]2.0.CO;2](https://doi.org/10.1206/0003-0090(2006)297[0001:TATOL]2.0.CO;2)
- Frost D 2016. Amphibian Species of the World 6.0. <http://research.amnh.org/vz/herpetology/amphibia>. Accessed on: 2016-10-1
- Gans C (2005) Checklist and bibliography of the Amphisbaenia of the world. *Bulletin of the American Museum of Natural History* 289: 1–130. [https://doi.org/10.1206/0003-0090\(2005\)289<0001:CABO>2.0.CO;2](https://doi.org/10.1206/0003-0090(2005)289<0001:CABO>2.0.CO;2)
- Garman S (1883) The reptiles and batrachians of North America. *Memoirs of the Museum of Comparative Zoölogy at Harvard College* Vol. 8 (3). Frankfort, Ky. Yeoman Press, 258 pp. <https://doi.org/10.5962/bhl.title.50435>
- Gibbons JW, Scott DE, Ryan TJ, Buhlmann KA, Tuberville TD, Metts BS, Greene JL, Mills T, Leiden Y, Poppy S, Winne CT (2000) The global decline of reptiles, déjà vu amphibians: reptile species are declining on a global scale. Six significant threats to reptile populations are habitat loss and degradation, introduced invasive species, environmental pollution, disease, unsustainable use, and global climate change. *BioScience* 50 (8): 653–666. [https://doi.org/10.1641/0006-3568\(2000\)050\[0653:tgddord\]2.0.co;2](https://doi.org/10.1641/0006-3568(2000)050[0653:tgddord]2.0.co;2)
- Goicoechea N, Frost DR, de la Riva I, Pellegrino KCM, Sites J, Rodrigues MT, Padial JM (2016) Molecular systematics of teioid lizards (Teioidea/Gymnophthalmoidea: Squamata) based on the analysis of 48 loci under tree-alignment and similarity-alignment. *Cladistics* 32 (6): 624–671. <https://doi.org/10.1111/cla.12150>
- Grant T, Frost DR, Caldwell JP, Gagliardo R, Haddad CFB, Kok PJR, Means DB, Noonan BP, Schargel WE, Wheeler WC (2006) Phylogenetic systematics of dart-poison frogs and their relatives (Amphibia: Athesphatanura: Dendrobatidae). *Bulletin of the American Museum of Natural History* 299: 1–262. <https://doi.org/10.5531/sd.sp.14>
- Gray JE (1845) Catalogue of the Specimens of Lizards in the Collection of the British Museum. Trustees of die British Museum/Edward Newman, London, 289 pp.
- Grazziotin FG, Zaher H., Murphy RW, Scrocchi G, Benavides MA, Zhang Y, Bonatto SL (2012) Molecular phylogeny of the New World Dipsadidae (Serpentes: Colubroidea): a reappraisal. *Cladistics* 28 (5): 437–459. <https://doi.org/10.1111/j.1096-0031.2012.00393.x>
- Gregory-Wodzicki KM (2000) Uplift history of the Central and Northern Andes: a review. *Geological Society of America Bulletin* 112 (7): 1091–1105. [https://doi.org/10.1130/0016-7606\(2000\)112<1091:UHOTCA>2.0.CO;2](https://doi.org/10.1130/0016-7606(2000)112<1091:UHOTCA>2.0.CO;2)
- Guayasamin JM, Castroviejo-Fisher S, Trueb L, Ayarzagüena J, Rada M, Vilá C (2009) Phylogenetic systematics of Glassfrogs

- (Amphibia: Centrolenidae) and their sister taxon *Allophryne ruthveni*. *Zootaxa* 2100: 1–97.
- Günther A (1858) Catalogue of Colubrine Snakes of the British Museum. London, Printed by order of the Trustees, 318 pp. <https://doi.org/10.5962/bhl.title.13272>
- Günther A (1860) Third list of the cold-blooded Vertebrata collected by Mr. Fraser in Ecuador. *Proceedings of the Zoological Society of London* 28: 233–241.
- Günther A (1885) *Biologia Centrali-Americana. Reptilia and Batrachia*. Taylor, & Francis, London, 402 pp. <https://doi.org/10.5962/bhl.title.730>
- Hallowell E (1845) Descriptions of reptiles from South America, supposed to be new. *Proceedings of the Academy of Natural Sciences of Philadelphia* 2: 241–247.
- Hallowell E (1856) Notes on reptiles in the collection of the Academy of Natural Sciences of Philadelphia. *Proceedings of the Academy of Natural Sciences of Philadelphia* 8: 221–238.
- Harvey MB, Ugueto GN, Gutberlet RL Jr (2012) Review of teiid morphology with revised taxonomy and phylogeny of the Teiidae (Lepidosauria: Squamata). *Zootaxa* 3459: 1–156.
- Hedges SB, Duellman WE, Heinicke MP (2008) New World direct-developing frogs (Anura: Terrarana): molecular phylogeny, classification, biogeography, and conservation. *Zootaxa* 1737: 1–182.
- Hedges S, Blair S (2011) The type species of the threadsnake genus *Tricheilostoma* Jan revisited (Squamata, Leptotyphlopidae). *Zootaxa* 3027: 63–66.
- Heyer WR (1968) The proper name for the type-species of the genus *Leptodactylus*. *Copeia* 1968: 160–162.
- Heyer WR (1994) Variation within the *Leptodactylus podicipinus-wagneri* complex of frogs (Amphibia: Leptodactylidae). *Smithsonian Contributions to Zoology* 546: 1–124.
- Heyer WR (2005) Variation and taxonomic clarification of the large species of the *Leptodactylus pentadactylus* species group (Amphibia: Leptodactylidae) from Middle America, northern South America, and Amazonia. *Arquivos de Zoologia* 37: 269–348.
- Hoge AR, Romano SA, Cordeiro CL (1978) Contribuição ao conhecimento das serpentes do Maranhão, Brasil. *Serpentes, Boidae, Colubridae, e Viperidae. Memórias do Instituto de Butantan* 40/41: 37–52.
- Hoge AR (1966) Preliminary account on Neotropical Crotalinae (Serpentes: Viperidae). *Memórias do Instituto de Butantan* 32: 109–184.
- Holdridge L (1964) *Life Zone Ecology*. Tropical Science Center, San José, 206 pp.
- Houttuyn M (1782) Het onderscheid der salamanderen van de haagdissen in 't algemeen, en van de gekkos in 't byzonder aangetoond. *Venhandelingen Uitgegeven door het Zeeuwsch Genootschap der Wetenschappen te Vlissingen, serie 1* 9: 305–336.
- IUCN (2016) The IUCN Red List of Threatened species. Version 2014.2. <http://www.iucnredlist.org>. Accessed on: 2016-10-1
- Jadin RC, Burbrink FT, Rivas GA, Vitt LJ, Barrio-Amorós CL, Guralnick RP (2014) Finding arboreal snakes in an evolutionary tree: phylogenetic placement and systematic revision of the Neotropical birdsnakes. *Journal of Zoological Systematics and Evolutionary Research* 52: 257–264. <https://doi.org/10.1111/jzs.12055>
- Jan G (1858) Plan d'une iconographie descriptive des phidiens et description sommaire de nouvelles espèces des serpents. *Revue et Magasin de Zoologie Paris* 10: 438–449, 514–527.
- Jan G (1862) Enumerazione sistematico delle specie d'ofidi del gruppo Calamaridae. *Archivio per la zoologia, l'anatomia e la fisiologia* 2: 1–76.
- Kattan GH, Franco P, Rojas V, Morales G (2004) Biological diversification in a complex region: a spatial analysis of faunistic diversity and biogeography of the Andes of Colombia. *Journal of Biogeography* 31(11): 1829–1839. <https://doi.org/10.1111/j.1365-2699.2004.01109.x>
- Köhler G (2008) *Reptiles of Central America*, 2nd ed. Herpeton Verlag, Offenbach, 400 pp.
- Köhler J, Böhme W (1996) Anuran amphibians from the region of Pre-Cambrian rock outcrops (inselbergs) in northeastern Bolivia, with a note on the gender of *Scinax* Wagler, 1830 (Hylidae). *Revue Française d'Aquariologie, Herpétologie* 23: 133–140.
- Laurenti JN (1768) *Specimen Medicum, Exhibens Synopsin Reptilium Emendatum cum Experimentis Circa Venena et Antidota Reptilium Austriacorum*. Joan. Thom. Nob. De Trattner, Wien, 214 pp.
- Lichtenstein H, Martens von E (1856) *Nomenclator Reptilium et Amphibiorum Musei Zoologici Beronlinensis. Namenverzeichnis der in der zoologischen Sammlung der Königlichen Universität zu Berlin aufgestellten Arten von Reptilien und Amphibien nach ihren Ordnungen, Familien und Gattungen*. Königliche Akademie der Wissenschaften, Berlin, 64 pp. <https://doi.org/10.5962/bhl.title.45245>
- Linnaeus C (1758) *Systema Naturae per Regna Tria Naturae, Secundum Classes, Ordines, Genera, Species, cum Characteribus, Differentiis, Synonymis, Locis*. 10th Edition. Volume 1.L. Salvii. Stockholm, 881 pp. <https://doi.org/10.5962/bhl.title.542>
- Lynch JD (1970) Systematic status of the American leptodactylid frog genera *Engystomops*, *Eupemphix*, and *Physalaemus*. *Copeia* 1970: 488–496.
- Lynch JD (1980) Systematic status and distribution of some poorly known frogs of the genus *Eleutherodactylus* from the Chocó lowlands of South America. *Herpetologica* 36(2): 175–189.
- Lynch JD (2001) Three new rainfrogs of the *Eleutherodactylus diastema* group from Colombia and Panama. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 25: 287–297.
- Lynch JD (2012) El contexto de las serpientes de Colombia con un análisis de las amenazas en contra de su conservación. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 36(140): 435–449.
- Lynch JD, Duellman WE (1973) A review of the centrolenid frogs of Ecuador, with descriptions of new species. *Occasional Papers of the Museum of Natural History, University of Kansas* 16: 1–66.
- Lynch JD, Rueda-Almonacid JV (1999) New species of frogs from low and moderate elevations from the Caldas transect of the eastern flank of the Cordillera Central. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 23: 307–314.
- Lynch JD, Ruiz-Carranza PM, Ardila-Robayo MC (1997) Biogeographic patterns of Colombian frogs and toads. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 21(80): 237–248.
- Marin CM, Toro FA, Daza JM (2017) First trans-Andean record of *Atractus occipitoalbus* (Jan, 1862), (Squamata, Dipsadidae), from Colombia. *Herpetology Notes* 10: 49–51.
- Maturana HR (1962) A study of the species of the genus *Basiliscus*. *Bulletin of the Museum of Comparative Zoology at Harvard College* 16: 1–34.
- McDiarmid RW, Campbell JA, Touré TA (1999) *Snake Species of the World*. Vol. 1. Herpetologists' League, Washington, DC, 511 pp.
- Merrem B (1820) Versuch eines Systems der Amphibien I (Tentamen Systematis Amphibiorum). J. C. Kriegeri, Marburg, 191 pp.
- Minambiente (2012) *Manual para la Asignación de Compensaciones por Pérdida de Biodiversidad*. Ministerio de Ambiente y Desarrollo Sostenible, República de Colombia, Bogotá, 52 pp.
- Molina-Zuluaga CM, Cano E, Restrepo A, Rada M, Daza JM (2017) Out of Amazonia: the unexpected trans-Andean distribution of *Cochranella resplendens* (Lynch and Duellman, 1978) (Anura: Centrolenidae). *Zootaxa* 4238(2): 268–274. <https://doi.org/10.11646/zootaxa.4238.2.8>
- Montingelli GG, Valencia JH, Benavides MA, Zaher H (2011) Revalidation of *Herpetodryas reticulata* (Peters, 1863) (Serpentes: Colubridae) from Ecuador. *South American Journal of Herpetology* 6: 189–197. <https://doi.org/10.2994/057.006.0304>
- Morales-Betancourt MA, Lasso CA, Páez VP, Bock BC (2015) *Libro rojo de reptiles de Colombia* (2015). Instituto de Investigación de Recursos Biológicos Alexander von Humboldt (IAvH), Universidad de Antioquia. Colombia, Bogotá DC, 258 pp.

- Muñoz-Ortiz A, Velásquez-Álvarez AA, Guarnizo CE, Crawford AJ (2015) Of peaks and valleys: testing the roles of orogeny and habitat heterogeneity in driving allopatry in mid-elevation frogs (Aromobatidae: *Rheobates*) of the northern Andes. *Journal of Biogeography* 42: 193–205. <https://doi.org/10.1111/jbi.12409>
- Myers CW (1974) The systematics of *Rhadinaea* (Colubridae), a genus of new world snakes. *Bulletin of the American Museum of Natural History* 153 (1): 1–262.
- Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GA, Kent J (2000) Biodiversity hotspots for conservation priorities. *Nature* 403 (6772): 852–858. <https://doi.org/10.1038/35002501>
- Nieto-Castro MJ (1999) Estudio preliminar de las especies del género *Scinax* (Amphibia: Anura: Hylidae) en Colombia. *Revista de la Academia Colombiana de Ciencias* 23 (suplemento especial): 339–346.
- Noble GK (1916) Description of a new eublepharid lizard from Costa Rica. *Proceedings of the Biological Society of Washington* 29: 87–88.
- Noble GK (1924) Some neotropical batrachians preserved in the United States National Museum with a note on the secondary sexual characters of these and other amphibians. *Proceedings of the Biological Society of Washington* 37: 65–72.
- Ofedal OT (1974) A revision of the genus *Anadia* (Sauria, Teiidae). *Archivos de Zoología* 25 (4): 203–265.
- Orrico VGD, Nunes I, Mattedi C, Fouquet A, Lemos AW, Rivera-Correa M, Lyra ML, Loebmann D, Pimenta BVS, Caramaschi U, Rodrigues MT, Haddad CFB (2017) Integrative taxonomy supports the existence of two distinct species within *Hypsiboas crepitans* (Anura: Hylidae). *Salamandra* 53: 99–113.
- Ospina-Sarría JJ, Angarita-Sierra T, Pedroza-Banda R (2015) A new species of *Craugastor* (Anura: Craugastoridae) from the Magdalena river valley, Colombia, with evaluation of the characters used to identify species of the *Craugastor fitzingeri* group. *South American Journal of Herpetology* 10: 165–177. <https://doi.org/10.2994/sajh-d-14-00014.1>
- Pérez-Santos C, Moreno AG (1988) Ofidios de Colombia. *Museo Regionale di Scienze Naturali*. Torino, 517 pp.
- Peters, J.A. and R. Donoso-Barros. 1970. Catalogue of the Neotropical Squamata: Part II. Lizards and Amphisbaenians. *United States National Museum Bulletin* 297 (2): 1–293. <https://doi.org/10.5479/si.03629236.297.1>
- Peters JA, Orejas-Miranda B, Donoso-Barros R (1970) Catalogue of the Neotropical Squamata: Part I. *United States National Museum Bulletin* 297. <http://dx.doi.org/10.5962/bhl.title.46653>
- Peters WCH (1857) Vier neue amerikanische Schlangen aus der Familie der Typhlopinen und darüber einige vorläufige Mittheilungen. *Monatsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin* 1857: 402.
- Peters WCH (1862) Über die Batrachier-Gattung Hemiphractus. *Monatsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin* 1862: 144–152.
- Peters WCH (1863) Fernere Mittheilungen über neue Batrachier. *Monatsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin* 1863: 445–470.
- Peters WCH (1867) Über Flederthiere (*Pteropus gouldii*, *Rhinolophus deckenii*, *Vespertilio lobipes*, *Vesperugo temminckii*) und Amphibien (*Hypsilurus godeffroyi*, *Lygosoma scutatum*, *Stenostoma narisostre*, *Onychocephalus unguirostris*, *Ahaetulla poylepis*, *Pseudechis scutellatus*, *Hoplobatrachus reinhardtii*, *Hyla coriacea*). *Monatsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin* 1867: 703–712.
- Peters WCH (1869) Über eine neue Nagergattung, *Chiropodomys penicillatus*, sowie über einige neue oder weniger bekannte Amphibien und Fische. *Monatsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin* 1868: 448–461.
- Peters WCH (1873) Über eine neue Schildkrötenart, *Cinosternon Effeldtii* und einige andere neue oder weniger bekannte Amphibien. *Monatsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin* 1873: 603–618.
- Pinto-Sánchez NR, Calderón-Espinosa ML, Miralles A, Crawford AJ, Ramírez-Pinilla MP (2015) Molecular phylogenetics and biogeography of the Neotropical skink genus *Mabuya* Fitzinger (Squamata: Scincidae) with emphasis on Colombian populations. *Molecular phylogenetics and Evolution* 93: 188–211. <https://doi.org/10.1016/j.ympev.2015.07.016>
- Pinto RR, Passos P, Caicedo-Portilla JR, Arredondo JC, Fernández R (2010) Taxonomy of the threadsnakes of the tribe Epictini (Squamata: Serpentes: Leptotyphlopidae) in Colombia. *Zootaxa* 2724: 1–28. <https://doi.org/10.5281/zenodo.199951>
- Poveda G, Mesa OJ, Salazar LF, Arias PA, Moreno HA, Vieira SC, Agudelo PA, Toro VG, Álvarez JF (2005) The diurnal cycle of precipitation in the tropical Andes of Colombia. *Monthly Weather Review* 133 (1): 228–240. doi: <https://doi.org/10.1175/MWR-2853.1>
- Pramuck JB (2006) Phylogeny of South American *Bufo* (Anura: Bufonidae) inferred from combined evidence. *Zoological Journal of the Linnean Society* 146: 407–452. <https://doi.org/10.1111/j.1096-3642.2006.00212.x>
- Restrepo A, Velasco J, Daza JM (2017) Extinction risk or lack of sampling in a threatened species: Genetic structure and environmental suitability of the Neotropical frog *Pristimantis penelopus* (Anura: Craugastoridae). *Papeis Avulsos de Zoologia* 57 (1): 1–15. <https://doi.org/10.11606/0031-1049.2017.57.01>
- Rivera-Correa M, Gutiérrez-Cárdenas PDA (2012) A new highland species of treefrog of the *Dendropsophus columbianus* group (Anura: Hylidae) from the Andes of Colombia. *Zootaxa* 3486: 50–62.
- Rivera-Prieto DA, Rivera-Correa M, Daza JM (2014) A new colorful species of *Pristimantis* (Anura: Craugastoridae) from the eastern flank of the Cordillera Central in Colombia. *Zootaxa* 3900 (2): 223–242. <http://dx.doi.org/10.11646/zootaxa.3900.2.3>
- Rivero JA, Serna MA (2000) Nuevos *Colostethus* (Amphibia, Dendrobatiidae) del Departamento de Antioquia, Colombia, con la descripción del renacuajo de *Colostethus fraterdanieli*. *Revista de Ecología Latino-Americana* 2: 45–58.
- Roze JA (1959) El genero *Erythrolamprus* Wagler (Serpentes; Colubridae) en Venezuela. *Acta Biológica Venezolánica* 2: 523–534.
- Ruiz-Carranza PM Lynch JD (1995) Ranas Centrolenidae de Colombia V. Cuatro nuevas especies de *Cochranella* de la Cordillera Central. *Lozania* 62: 1–23.
- Ruiz-Carranza PM Lynch JD (1991) Ranas Centrolenidae de Colombia I. Propuesta de una nueva clasificación genérica. *Lozania* 57: 1–30.
- Ruthven AG Gaige HT (1924) A new *Leposoma* from Panama. *Occasional Papers of the Museum of Zoology, University of Michigan* 147: 1–3.
- Savage JM (1968) The dendrobatid frogs of Central America. *Copeia* 1968: 745–776.
- Savage JM (2002) The Amphibians and Reptiles of Costa Rica: A Herpetofauna between Two Continents, between Two Seas. *University of Chicago Press, Chicago*, 934 pp.
- Savage JM, Crother BI (1989) The status of *Pliocercus* and *Urotheca* (Serpentes: Colubridae), with a review of included species of coral snake mimics. *Zoological Journal of the Linnean Society* 95: 335–362. <https://doi.org/10.1111/j.1096-3642.1989.tb02315.x>
- Schlegel H (1827) Herpetologische Nachrichten. *Isis von Oken* 20: 281–294.
- Schlegel H (1841) Description d'une nouvelle espece du genre *Trigonocephale* (*Trigonocephalus lansbergii*). *Magazine of Zoology Report* 1–3: 1.
- Schneider JG (1799) *Historia Amphibiorum Naturalis et Literariae, Fasciculus Primus. Continens Ranas, Calamitas, Bufones, Salamandras et Hydros in Genera et Species Descriptos Notisque suis Distinctos*. Friederici Frommanni, Jenna, 290 pp. <http://dx.doi.org/10.5962/bhl.title.4270>
- Sentzen UJ (1796) Ophiologische Fragmente. *Meyer's Zoological Archives* 2: 49–61.
- Shaw G (1802) *General Zoology, or Systematic Natural History*. Vol. 3,

- part 2. London, G. Kearsley, Thomas Davison, 313–615.
- Smith HM (1941) A new name for the Mexican snakes of the genus *Dendrophidion*. Proceedings of the Biological Society of Washington 54: 73–76
- Solís F, Ibáñez R, Jaramillo C, Fuenmayor Q, Castro F, Grant T (2010) *Hyalinobatrachium aureoguttatum*. The IUCN Red List of Threatened Species 2010: e.T55003A11234997. <https://doi.org/10.2305/iucn.uk.2010-2.rlts.t55003a11234997.en>. Accessed on: 2015-11-27.
- Spix JB von (1824) Serpentes brasiliensium species novae ou histoire naturelle des espèces nouvelles de serpens, recueillies et observées pendant le voyage dans l'intérieur du Brésil dans les années 1817, 1818, 1819, 1820, exécuté par ordre de sa majesté le Roi de Bavi Typis Franc. Seraph. Monachii, Hübschmanni, viii, 75 pp.
- Starrett PH (1960) A redefinition of the genus *Smilisca*. Copeia 1960: 300–304.
- Taylor EH (1968) The Caecilians of the World: A Taxonomic Review. University of Kansas Press, Lawrence, 848 pp.
- Taylor EH, Smith HM (1943) A review of American sibynophine snakes, with a proposal of a new genus. University of Kansas Science Bulletin 29 (6): 301–337.
- Taylor EH (1949) A preliminary account of the herpetology of the state of San Luis Potosi, Mexico. University of Kansas Science Bulletin 33 (2): 169–215.
- Thomiot A (1884) Note sur un batracien d'espèce nouvelle provenant de Panama. Bulletin de la Société Philomathique de Paris, série 7–8: 151–152.
- Tolosa Y, Molina-Zuluaga, C, Restrepo A, Daza JM (2014) Sexual maturity and sexual dimorphism in a population of the rocket-frog *Colostethus* aff. *fraterdanieli* (Anura: Dendrobatidae) on the north-eastern Cordillera Central of Colombia. Actualidades Biológicas 37 (102): 287–294.
- Uetz P, Hošek J (2016) The Reptile Database. <http://www.reptile-database.org/>. Accessed on: 2016-10-1.
- Vanzolini PE (1953) Sobre a diferenciação geografica de *Gymnodactylus geckoides* (Sauria, Gekkonidae) and Sobre a Presença do genero *Lepidoblepharis* no Brasil (Sauria, Gekkonidae). Papéis Avulsos Zoologia 11 (14/15): 225–270.
- Vanzolini PE (2002) An aid to the identification of the South American species of *Amphisbaena* (Squamata, Amphisbaenidae). Papéis Avulsos de Zoologia 42 (15): 351–362. <https://doi.org/10.1590/S0031-10492002001500001>
- Vanzolini PE (1951) *Amphisbaena fuliginosa*. Contribution to the knowledge of the Brazilian lizards of the family Amphisbaenidae Gray, 1825. 6. On the geographical distribution and differentiation of *Amphisbaena fuliginosa* Linné. Bulletin of the Museum of Comparative Zoology at Harvard College 106 (1): 1–67.
- Wied-Neuwied M (1824) Verzeichniss der Amphibien, welche im zweyten Bande der Naturgeschichte Brasiliens vom Prinz Max von Neuwied warden beschrieben warden. Isis von Oken 14: 661–673.
- Wiegmann AF (1833) Herpetologischen Beyträge. I. Weber die mexicanischen Kröten nebst bemerkungen über ihren verwandte Arten anderer Weltgegenden. Isis von Oken 26: 651–662.
- Wilkinson M, O'Connor A, Nussbaum RA (2013) Taxonomic status of the Neotropical caecilian genera *Brasilotyphlus* Taylor, 1968, *Microcaecilia* Taylor, 1968 and *Parvicaecilia* Taylor, 1968 (Amphibia: Gymnophiona: Siphonopidae). Occasional Papers of the Museum of Zoology, University of Michigan 744: 1–12.
- Wilson LD, Mena CE (1980) Systematics of the *melanocephala* group of the colubrid snake genus *Tantilla*. Memoirs of the San Diego Society of Natural History 11: 5–58.
- Zaher H (1996) A new genus and species of pseudoboine snake, with a revision of the genus *Clelia* (Serpentes, Xenodontinae). Bolletino dei Musei Regionale Scienza naturali di Torino 14: 289–337.
- Zaher H Prudente LC (1999) Intraspecific variation of the hemipenis in *Siphlophis* and *Tripanurgos*. Journal of Herpetology 33 (4): 698–702. <https://doi.org/10.2307/1565589>