

Yeast Primer

# New insights into a complex fungal pathogen: the case of *Paracoccidioides* spp.

Angel Gonzalez\* and Orville Hernandez

Basic and Applied Microbiology Research Group (MICROBA), School of Microbiology, Universidad de Antioquia, Medellín, Colombia

\*Correspondence to:

A. González, Basic and Applied Microbiology Research Group (MICROBA), School of Microbiology, Universidad de Antioquia, Calle 70 52–72, Of. 607, Medellín, Colombia.  
E-mail: angel.gonzalez@udea.edu.co

## Abstract

**Paracoccidioidomycosis is a systemic mycosis endemic to Latin America, with *Paracoccidioides brasiliensis* and *P. lutzii* being the causal agents of this disorder. Several issues have been raised in the 100 years since its discovery and in this article we discuss features of this fascinating fungal pathogen, including its biology, eco-epidemiology and aspects of its pathogenicity. We also consider some of its virulence determinants, the most recent advances in the study of its metabolic pathways and the molecular and genetic research tools developed for this research. We also review the animal models used to study host–fungal interactions and how the host defence mechanisms against this pathogen work. Copyright © 2015 John Wiley & Sons, Ltd.**

Received: 25 June 2015  
Accepted: 30 November 2015

**Keywords:** *Paracoccidioides brasiliensis*; *Paracoccidioides lutzii*; paracoccidioidomycosis; virulence factors; metabolic pathways; host defense mechanisms

## Introduction

The fungus *Paracoccidioides* spp. is the causal agent of paracoccidioidomycosis (PCM), a systemic mycosis endemic to Latin America with Brazil, Colombia, Argentina and Venezuela being the countries with the highest numbers of reported cases. An estimated 10 million people have been infected with this fungus to date (Brummer *et al.*, 1993; Restrepo *et al.*, 2015).


*Paracoccidioides* is a thermally dimorphic fungal pathogen that exhibits two morphotypes, i.e. a mould occurring at temperatures < 28 °C, composed of thin septated hyphae that in turn produce conidia (considered as the infectious propagules) (Restrepo *et al.*, 2011, 2015) and a yeast found in cultures or in the host at 37 °C composed of variably-sized oval to round cells (blastoconidia), characterized by the formation of larger mother cells surrounded by multiple daughter cells, resembling a pilot wheel (Figure 1) (Restrepo *et al.*, 2011, 2015; Marques, 2013).

The genus *Paracoccidioides* belongs to the Phylum Ascomycota, Class Euromycetes, Order

Onygenales and Family Ajellomycetaceae (Onygenaceae) (Bagagli *et al.*, 2008). It comprises two species: *P. brasiliensis*, considered to be a complex of four distinct phylogenetic lineages (S1, PS2, PS3 and PS4), and the recently described *P. lutzii* (formerly 'Pb01-like') (Matute *et al.*, 2006; Teixeira *et al.*, 2009; Salgado-Salazar *et al.*, 2010; Theodoro *et al.*, 2012; Teixeira *et al.*, 2013).

Clinical presentations of PCM include subclinical or asymptomatic infection and the symptomatic or clinical manifested disease which cause an acute/subacute or a chronic form, the latter involving the lungs as well as other organs (Restrepo *et al.*, 2015; Bocca *et al.*, 2013; Restrepo *et al.*, 2012).

Various functional (*in vitro* and *in vivo*) and molecular studies have been performed in order to evaluate and identify the role of several genes and proteins involved in fungus–host interactions; more recently, new strategies using anti-sense RNA technology have allowed the presence of virulence factors to be demonstrated, as well as that of molecules involved in both the oxidative and metabolic pathways (Menino *et al.*, 2012). Several animal models have led to an understanding of the



**Figure 1.** (A) *Paracoccidioides brasiliensis* yeast surrounded by budding daughter cells; wet mount from culture. (B) *P. brasiliensis* in mouse tissue; GMS stain; magnification  $\approx \times 1000$

pathogenesis of this fungal infection, from the initial infection process through the chronic stages, using both the conidia and yeast cells (González *et al.*, 2008b; Calich *et al.*, 1985; Singer-Vermes *et al.*, 1993a).

Here we present an overview of some of the scientific breakthroughs achieved in studies of *Paracoccidioides* spp. and briefly touch on some key aspects of its biology, including its eco-epidemiology, pathogenesis, host defence mechanisms, genome, genetics, metabolism and the use of new molecular tools to understand the biology of this fungal pathogen.

### Paracoccidioidomycosis: the disease

PCM is acquired after inhalation of infectious propagules in the environment, leading to a primary pulmonary infection. If the infection is not controlled by the immune response, patients may develop clinical manifestations, mainly as an acute/subacute or a chronic form (Restrepo *et al.*, 2012, 2015).

Although subclinical or asymptomatic infection is not associated with any specific clinical characteristics, in these patients the fungus may remain latent in the tissue for years without any manifestation of disease. The acute/subacute or juvenile-type disease evolves rapidly and mainly affects children or young adults, representing about 10% of all cases. This clinical form is characterized by involvement of the reticulo-endothelial organs (lymph nodes, liver, spleen) as well as skin and bone; digestive tract manifestations are also

common (Restrepo *et al.*, 2012, 2015). The chronic or adult-type form is the most common clinical presentation and accounts for approximately 80% of cases. This clinical form is characterized by infection of the lungs and extrapulmonary organs (mainly skin and mucous membranes), accompanied by the development of ulcerative, granulomatous and infiltrated lesions. The adrenal glands and CNS may also be involved. Pulmonary fibrosis, the main sequela of this disease, may develop in 50% of patients with the chronic form (Restrepo *et al.*, 2012, 2015).

### Eco-epidemiology

As described above, PCM is restricted to certain countries of Latin America. This mycosis has an estimated incidence of 1–3 cases/100 000 inhabitants (Bocca *et al.*, 2013; Bellissimo-Rodrigues *et al.*, 2011) and a mortality rate of 1.4/million, the highest value for any systemic mycosis (Coutinho *et al.*, 2002). One of its most peculiar characteristics is that it is more often diagnosed in males than females (ratio 13:1), and occurs in individuals engaged in agricultural activities (Restrepo *et al.*, 2012, 2015; Colombo *et al.*, 2011; Bellissimo-Rodrigues *et al.*, 2013). It is noteworthy that no outbreaks of this mycosis have ever been reported.

Regarding its phylogenetic classification, *P. brasiliensis* S1 is widely distributed throughout South America, PS2 has been identified only in Brazil and Venezuela, PS3 is apparently restricted to regions of Colombia and PS4 has to date been

reported only from Venezuela (Matute *et al.*, 2006; Salgado-Salazar *et al.*, 2010; Theodoro *et al.*, 2012). The other species, *P. lutzii*, is found mainly in central, south-western and northern Brazil, as well as Ecuador (Teixeira *et al.*, 2009, 2013; Theodoro *et al.*, 2012).

Several environmental and ecological factors have been associated with the diagnosis of PCM, including the presence of tropical and subtropical forests, moderate-to-high precipitation rates (2000–2999 mm/year), watercourses, mild temperatures (<27°C), fertile soils and elevation < 800 m (Calle *et al.*, 2001; Bagagli *et al.*, 2003; Restrepo *et al.*, 2012, 2015). In addition, the El Niño southern oscillation (ENSO) climatic anomaly has been linked with a cluster of 10 cases of acute PCM (Barrozo *et al.*, 2010).

This fungal pathogen has been isolated in culture, as well as being detected by molecular and serological methods, from several wild and domestic animals; these include armadillos (Bagagli *et al.*, 2003; Corredor *et al.*, 2005), raccoons, Brazilian guinea pigs, spiny tree porcupines, ferrets and tayras (Richini-Pereira *et al.*, 2008; Albano *et al.*, 2014), dogs (Ricci *et al.*, 2004; de Farias *et al.*, 2011; Fontana *et al.*, 2010), pigs (Belitardo *et al.*, 2014a), rabbits (Belitardo *et al.*, 2014b), cats (Oliveira *et al.*, 2013), dairy goats (Ferreira *et al.*, 2013), sheep (Oliveira *et al.*, 2012) and chickens (Oliveira *et al.*, 2011). *Paracoccidioides* has also been detected in soil samples from around or inside armadillo burrows (Theodoro *et al.*, 2005).

## Pathogenesis

Different *in vivo* and *in vitro* models have been devised to understand the pathogenesis of PCM. *In vivo*, it has been observed that once conidia or yeast cells of *P. brasiliensis* reach the lungs, these propagules interact initially with the extracellular matrix (ECM) proteins, epithelial cells, alveolar macrophages and pulmonary dendritic cells. Apparently, these interactions are mediated by adhesin-type molecules present on the fungal surface that instead recognize ECM proteins, also present on the epithelial cell surface (Caro *et al.*, 2008; González *et al.*, 2005a, 2008a). In addition, pulmonary cells (mainly macrophages) are activated and an inflammatory process occurs involving the

production of pro-inflammatory cytokines and chemokines; this process is characterized by expression of adhesion molecules in the host and subsequent recruitment of neutrophils and macrophages to the infection site (González *et al.*, 2003, 2005b). After interaction of the fungal propagules with neutrophils and macrophages, these phagocytic cells become activated and express molecules, such as nitric oxide, lysozyme and reactive oxygen intermediates, that may exert a fungicidal effect against *P. brasiliensis*; however, if the fungus is able to overcome these mechanisms, dissemination to other organs and systems will occur (Gonzalez *et al.*, 2000, 2008b; Moreira *et al.*, 2008a, 2008b).

The initial interaction of the phagocytic cells with *P. brasiliensis* has been described as being partly mediated by several molecules including complement receptor-3 (CR3), mannose receptor, dectin-1, Toll-like receptors (TLR)-2 and TLR-4, among others (Jiménez *et al.*, 2006; Loures *et al.*, 2009, 2010; Bonfim *et al.*, 2009).

Moreover, in PCM patients B lymphocytes are activated at polyclonal level with specific antibodies (mainly IgA, IgG and IgE) produced against the fungus (Chequer-Bou-Habib *et al.*, 1989; Baida *et al.*, 1999).

Regarding the immune response developed in PCM patients, it has been found that those with the severe form show increased production of Th2 cytokines (mainly IL-10, IL-4 and IL-5) whereas the Th1 cytokine levels (mainly IFN $\gamma$  and IL-2) are low (Kashino *et al.*, 2000; Benard, 2008). In addition, the different T cell subsets accomplish different functions in this mycosis, as evaluated in animal models; thus, T CD8<sup>+</sup> cells participate in the control of fungal burden, whereas T CD4<sup>+</sup> cells are involved in the delayed-type hypersensitivity response and production of specific protective antibodies (Chiarella *et al.*, 2007). T $\gamma$  $\delta$  lymphocytes appear to participate in the polyclonal activation of B cells, while T-reg (CD4<sup>+</sup> CD25<sup>+</sup> FoxP3<sup>+</sup>) cells are involved in the development of a local and systemic immune response in PCM (Munk *et al.*, 1995; Moreira *et al.*, 2008a, 2008b).

## Life cycle

An important aspect of the life cycle of *Paracoccidioides* is its thermo-dimorphism, in

which the mycelial morphotype occurs at ambient temperature, whereas the yeast morphotype grows at 37°C, equivalent to that of the mammalian host (Queiroz-Telles, 1994).

This is an important issue, since during the life cycle the fungus has to adapt to different conditions in the outside environment and within the host cell; in addition, cell cycle control is vital to maintaining correct cellular function and the transfer of genetic information to new daughter cells. Computational studies searching for genes involved in the cell cycle of *P. brasiliensis* have permitted the discovery of genes involved in cell cycle control, cytoskeleton structure and chromosome segregation, similar to those reported in *Saccharomyces cerevisiae* (Reis *et al.*, 2005). Few studies on this topic have been carried out for *Paracoccidioides*, however. Analysis of *P. brasiliensis* nuclear DNA by flow cytometry enabled four different cellular subpopulations (R1, R2, R4 and R5) to be identified, indicating a 1–4-fold increase in genomic DNA content, characteristic of multiple budding and/or polynucleated cells, as expected in *Paracoccidioides* (Almeida *et al.*, 2006). Approximately 90% of the overall population consisted of two main subpopulations, having 1n and 2n DNA content, R1 and R2 respectively; of note, the R2 subpopulation consisted of mononucleate and binucleate cells that may correspond to cells at different phases of the cycle. Alternatively, these two nuclei may correspond to multinucleate cells; treatment with antifungal drugs induced arrest in the cell cycle profile, suggesting an alteration in nuclear division of *P. brasiliensis* yeast (Rodrigues *et al.*, 2003; Almeida *et al.*, 2006).

On the other hand, several pieces of evidence suggest that this fungal pathogen has a sexual stage. These evidences include the capacity of recombination and the presence of two mating type idiomorphs; thus, each *Paracoccidioides* spp. isolate contains the **a**-box (MAT1-1) or HMG (MAT-12) mating type idiomorph (Matute *et al.*, 2006; Torres *et al.*, 2009).

## Genome and genetics

The first studies in this area were conducted by San-Blas (1986), who demonstrated that both

mycelia and yeast morphotypes contain multiple nuclei, whereas conidia present only a single nucleus. Pulsed-field gel electrophoresis (PFGE) assays were later used to obtain two clinical isolates of *P. brasiliensis*, for which four megabase-sized (2.0–10.0 Mb) bands were identified. A genome size of 45.7–60.9 Mb was estimated for these isolates, based on ranging microfluorometric analysis (M. I. Cano *et al.*, 1998). The same group subsequently employed the contour-clamped homogeneous electric field gel electrophoresis (CHEF) technique and reported a molecular size in the range 3.2–10 Mb for *P. brasiliensis*, with a genome size of approximately 29.7 Mb (Montoya *et al.*, 1999). The differences in genome size suggest that the nuclei of *P. brasiliensis* yeast cells could be diploid, without discarding the possibility of the isolates being haploid or aneuploid. In a later, well-conducted study, Almeida *et al.* (2007a) used flow cytometry (FCM) protocols to evaluate the nucleic content of 10 *P. brasiliensis* isolates from four different PCM-endemic areas (Brazil, Colombia, Uruguay and Venezuela; these isolates represented three different identified cryptic species (S1, PS2 and PS3) (Matute *et al.*, 2006). Results of this study indicated that *P. brasiliensis* were mononucleate yeast cells with a genome size in the range 23–31 Mb, similar to that seen in the previous study (Montoya *et al.*, 1999). Furthermore, the ploidy ratio of 1.0:1.1 observed in the isolates evaluated indicates that this fungus has a haploid DNA content (Almeida *et al.*, 2007a).

In a collaborative study between the Broad Institute and *Paracoccidioides* research community, three referenced isolates (Pb18, Pb03 and Pb01) were selected, representing the S1, PS2 and *P. lutzii* lineages, respectively, for sequencing and genomic analysis. The assembly size for these strains varied in the range 29.1–32.9 Mb and 7875–9132 genes were identified (Desjardins *et al.*, 2011). Even with the methods available at that time, the results produced high-quality draft assemblies, although these included many gaps and nucleotides of uncertain or low quality in the final consensus sequences. More recently, this limitation was overcome by employing high sequencing depth to improve the original assemblies, generated from Sanger sequence reads of the same isolates, allowing more complete and accurate reference assemblies to be obtained (Muñoz *et al.*, 2014). *Paracoccidioides* spp. genome information is



available at: [http://www.broad.mit.edu/annotation/genome/paracoccidioides\\_brasiliensis](http://www.broad.mit.edu/annotation/genome/paracoccidioides_brasiliensis)


These results could be used in future for diverse molecular and biochemical projects as well as in functional studies, not only of *Paracoccidioides* but also in other fields of medical mycology.

## Metabolic pathways

Several studies have been carried out with a view to understanding the metabolic pathways in *Paracoccidioides* spp. that are involved both in cell growth and adaptation to host conditions. These have demonstrated that fungal pathogens have sufficient metabolic plasticity to allow them to exploit different carbon sources (Brock, 2009; Fleck *et al.*, 2011). Arraes *et al.* (2005) found that *P. brasiliensis* is able to develop the most important pathways of central metabolism, involving carbohydrates, lipids, amino acids and nucleotides (Arraes *et al.*, 2005). The major metabolic pathways described so far for *P. brasiliensis* are shown in Figure 2.

## Carbohydrates

With regard to metabolism of carbohydrates, *Paracoccidioides* spp. has both glycolytic and gluconeogenic pathways, the former to obtain pyruvate and the latter to synthesize glucose from non-carbohydrate carbon substrates. The enzymes required for these processes are present in both the mycelium and yeast morphotypes, suggesting that this fungus has sufficient metabolic plasticity to employ different carbon sources and produce glucose, as has been described for other fungi (Arraes *et al.*, 2005; Brock, 2009; Fleck *et al.*, 2011; Ene *et al.*, 2012; Price *et al.*, 2011). Tavares *et al.* (2015) demonstrated that *Paracoccidioides* spp. employ a fermentation process to obtain energy; this strategy could thus facilitate fungal adaptation to glucose-poor microenvironments, such as the phagosomes inside leukocytes. In addition, this pathogen could also produce ATP under low oxygen conditions, in turn reducing the levels of reactive oxygen species produced by the host (Tavares *et al.*, 2015). This was confirmed by Felipe *et al.* (2005), who reported that the fermentation process was mainly developed in *Paracoccidioides* yeast cells; meanwhile aerobic-dependent ATP production



**Figure 2.** A scheme representing the major metabolic pathways described in *Paracoccidioides*. Inside the host tissues, this fungal pathogen reduces the levels of phosphofruktokinase, the enzyme which catalyses phosphorylation of fructose-6 phosphate to fructose-1,6-bisphosphate. In the same way, reactive oxygen species (ROS) or reactive nitrogen species (RNS) decrease the levels of enzymes involved in aerobic respiration. Alterations to any of these pathways generate decreases in the glycolytic pathway. To compensate for the decrease in ATP obtained by carbohydrate metabolism, the fungus possibly increases lipolytic metabolism, this being supported by increases in the methyl citrate and lipid  $\beta$ -oxidation cycles

occurs in the mycelium morphotype. Tavares *et al.* (2007) demonstrated that *Paracoccidioides* spp. co-cultured with murine macrophages showed reduced expression of the mRNA levels of the gene coding for phosphofructokinase, an enzyme that catalyses the phosphorylation of fructose 6-phosphate to fructose 1,6-bisphosphate in the glycolytic pathway, which would provide an alternative way to obtain energy. To confirm this, proteomic studies were performed using *Paracoccidioides* yeast cells after exposition to reactive nitrogen species (RNS) by adding of *S*-nitrosoglutathione (GSNO; a stress generator). In these studies, reduced levels of the enzymes involved in aerobic respiration were observed, such as cytochromes, succinate dehydrogenase and ATP synthases (Parente *et al.*, 2015). These results reinforce the hypothesis that this fungal pathogen uses a fermentation process to obtain energy inside host cells.

On the other hand, a better understanding of the metabolism of the fungus during its adaptation to the host tissues is important to elucidate the alternative metabolic pathway it might adopt under conditions of carbon starvation. Lima *et al.* (2014) reported that the major changes in the transcriptional profile of *Paracoccidioides* during carbon starvation were related to gluconeogenesis and ethanol production. These findings were supported by modulation of the glyoxylate and tricarboxylic cycles; degradation of amino acids and fatty acids was also observed.

## Lipids

Regarding the metabolism of lipids, 16 enzymes have been identified in *Paracoccidioides* as responsible for ergosterol biosynthesis, and many others are involved in different cellular processes. An important characteristic of this fungus is its ability to disseminate from the lungs to other organ systems (Mendes-Giannini *et al.*, 2008). During this process, the fungus must adapt to lipolytic metabolism; Bailão *et al.* (2006) demonstrated that this process is characterized by the expression of 2-methylcitrate dehydratase and acyl CoA dehydrogenase genes which are involved in the methyl citrate and lipid  $\beta$ -oxidation cycles, respectively. In addition, high concentrations of proteins involved in lipid synthesis (fatty acid synthase subunit  $\beta$ -dehydratase, fatty acid synthase subunit  $\alpha$ -reductase and trans-2-enoyl CoA reductase) have been observed during the

interaction of *Paracoccidioides* and RNS (Parente *et al.*, 2015).

## Amino acids and nucleotides

*Paracoccidioides* can synthesize all amino acids except asparagine (Asp). The complete pathway for DOPA–melanin biosynthesis has also been found in this fungus (Arraes *et al.*, 2005). It has been demonstrated that during nitrosative stress in *Paracoccidioides*, four enzymes involved in amino acid metabolism (branched chain amino acid aminotransferase, acetolactate synthase, isovaleryl-CoA dehydrogenase and a subunit of methylcrotonoyl-CoA carboxylase) showed increased activity (Parente *et al.*, 2015); these results are in agreement with those of Missall *et al.* (2006), who observed that amino acid metabolism is affected by nitrosative stress in *C. neoformans*. Further studies are required to characterize the role of RNS in both the synthesis and catabolism of these molecules in *Paracoccidioides*.

To date, seven enzymes involved in *de novo* purine synthesis and four in pyrimidine biosynthesis have been described in *Paracoccidioides*. This fungus probably also produces GMP and AMP *de novo*; however, this hypothesis needs to be tested further, since these previous findings were based on transcriptome analysis (Arraes *et al.*, 2005).

'Moonlighting' proteins are exceptional multifunctional molecules that can accomplish several functions pivotal to cellular metabolism (Huberts and van der Klei, 2010). Several moonlighting proteins are expressed in *Paracoccidioides*, including aconitase, aldolase, glyceraldehyde 3-phosphate dehydrogenase, isocitrate lyase, malate synthase, triose phosphate isomerase, fumarase and enolase. In addition to their metabolic functions, these proteins are involved in cell wall biosynthesis/remodelling, adaptation to different environmental conditions and pathogenesis (Marcos *et al.*, 2014). However, more studies are needed in order to understand their specific and global functions in *Paracoccidioides*.

## Metals (iron, zinc, copper) and haemoglobin

Various metals, such as iron, zinc and copper are essential to the metabolism of different microorganisms, including fungal pathogens such as *Paracoccidioides*. These metals participate in

different metabolic pathways as cofactors and are associated with proteins to form metalloproteins, which play an important role in cell membranes and are involved in cellular respiration (Nevitt, 2011).

Various studies regarding the role of iron in *Paracoccidioides* have demonstrated that this metal is required for conidia-to-yeast transition (an essential event for the development of the disease), as well as yeast replication inside macrophages and monocytes (Cano *et al.*, 1994; Gonzalez *et al.*, 2007). Parente *et al.* (2011) subsequently demonstrated that, under iron-depleted conditions, *P. brasiliensis* undergoes a glycolytic process instead of oxidative pathways, allowing it to survive inside the host cell. Other studies have demonstrated iron to be an important micronutrient required for the growth of several pathogens, including *Paracoccidioides*. Expression of genes involved in the production of siderophores (important iron chelators) under iron-limited conditions has been demonstrated; these siderophores are present at both the extracellular and intracellular levels as coprogens, ferrichromes and fusarinines, and their expression is associated with survival of *Paracoccidioides* after co-culture with IFN $\gamma$ -activated macrophages (Silva-Bailão *et al.*, 2014). Moreover, siderophore production and iron uptake have been described as important factors in cellular metabolism, as well as acting as virulence factors for fungal pathogens (Schrettl *et al.*, 2004; Hwang *et al.*, 2008).

In another related study, it was also demonstrated that *Paracoccidioides* may use haemoglobin as an iron source, most probably through receptor-mediated pathways, since iron-related genes that encode haemoglobin receptors were identified in the genome of this fungal pathogen. In this study, the authors observed that expression of these related genes, including those encoding for proteins involved in amino acid assembly, as well as nitrogen, sulphur and iron–sulphur metabolism, were induced under low-inorganic iron conditions or in the presence of haemoglobin (Bailão *et al.*, 2014).

Although copper and zinc homeostasis have not been extensively studied in *Paracoccidioides*, increased copper transportation has been observed in the tissues of mice infected with *P. brasiliensis* (Bailão *et al.*, 2006). Increased levels of mitochondrial copper transporter and Cu/Zn superoxide dismutase were subsequently observed in *P.*

*brasiliensis* yeast cells infecting macrophages (Tavares *et al.*, 2007). Metalloproteins are known to be involved in many cellular processes, including general metabolism and virulence. Several metalloproteins, such as Cu-, Fe- and Zn-binding proteins, have been identified using bioinformatic tools, representing 7% of the total proteins encoded by the genome of *Paracoccidioides*. Zinc proteins are the most abundant, representing 5.7% of the fungus proteome, with copper and iron proteins accounting for 0.3% and 1.2%, respectively (Tristão *et al.*, 2015).

All the above studies clearly demonstrate the importance of metals (particularly iron) in the cellular metabolism of *Paracoccidioides* as well as its interaction with host cells. However, the exact mechanism by which these metals promote or participate in the metabolism of this fungus is not yet fully understood. Further studies should be carried out on this important topic.

### Oxidative and nitrosative stress response

It has recently been demonstrated that the alternative oxidase (PbAOX) plays an important role in intracellular redox balancing in *P. brasiliensis*; this enzyme is also involved with other components in the mitochondrial respiratory process (Ruiz *et al.*, 2011; Martins *et al.*, 2011). Moreover, additional studies have shown that PbAOX is important not only for maintaining cellular homeostasis, by assisting redox balancing during cell growth and morphological changes, but also in fungal defence against the oxidative stress imposed by immune cells to combat *P. brasiliensis*. This demonstrates that the oxidative stress response is essential to both fungal survival and adaptation to host conditions (Hernández *et al.*, 2015; Ruiz *et al.*, 2011).

Hydrogen peroxide is another molecule that participates in oxidative stress. After the addition of H<sub>2</sub>O<sub>2</sub> to *Paracoccidioides* cultures, it was observed that this pathogen expressed anti-oxidant enzymes, such as catalase, superoxide dismutase, cytochrome *c* peroxidase and thioredoxin, with substantial changes in its metabolism, as observed by activation of pathways to provide NAD(P)H (de Arruda Grossklaus *et al.*, 2013). These studies also showed that an appropriate response to oxidative stress of *Paracoccidioides* is an essential event for its dimorphism process and adaptation to host conditions, both requiring metabolic adaptation by the fungus.

On the other hand, RNS has been reported to reduce the activity of the mitochondrial transport chain in *Paracoccidioides*, possibly through inhibition of aconitase, cytochrome oxidase and other mitochondrial proteins involved in this process (Parente *et al.*, 2015; Mason *et al.*, 2006). The inhibition exerted by RNS compounds is apparently a non-selective but irreversible process, whereas nitric oxide induces a rapid, selective but reversible inhibition of cytochrome oxidase (Brown, 1999). These results clearly indicate that these reactive species not only alter the mitochondrial transport chain in *Paracoccidioides* but also affect important metabolic processes related to ATP production, dimorphism and host adaptation, among others.

### Molecular and research tools

Among the molecular tools that have been employed for studying *Paracoccidioides* spp., amplification of nucleic acids (mainly DNA) has been useful for diagnosing PCM. Development of PCR protocols using primers or probes and targeting the *GP43* gene or the ITS1 ribosomal DNA regions have been the most common assays reported. These have proved to be highly sensitive and specific, able to detect as few as 10 copies or 1.1 pg/ml DNA (Semighini *et al.*, 2002; Dias *et al.*, 2012). Other protocols using conventional, nested or semi-nested PCR are useful for detecting fungal DNA in clinical samples, with the exception of those from serum, which do not appear to be effective (Pitz Ade *et al.*, 2013; Dias *et al.*, 2012; Teles and Martins, 2011; Motoyama *et al.*, 2000).


On the other hand, taking into account that homologous recombination is unusual in *Paracoccidioides*, the development of knockout isolates is not currently feasible (Sturme *et al.*, 2011). Thus, one of the most important advances for functional studies of *Paracoccidioides* has been the implementation of antisense technology. Initially, Leal *et al.* (2004) reported transformation of *P. brasiliensis* yeast cells using *Agrobacterium tumefaciens* GV3101 and the vector pAD1625, although transformation efficiency was not satisfactory (Leal *et al.*, 2004). Afterwards, an efficient strategy based in *Agrobacterium tumefaciens*-mediated transformation (ATMT) was used to obtain *P. brasiliensis* isolates, with downregulation of specific genes; in this study the authors

presented evidence for the occurrence of random single-gene copy integration per haploid nucleus and the generation of homokaryon progeny, a finding that was key to subsequent studies of the biology of this fungus (Almeida *et al.*, 2007b). This approach quickly began to be used for functional studies in *P. brasiliensis*. The first target was CDC42; decreased expression of this molecule in *P. brasiliensis* yeast was associated with reduced cell size and more homogeneous cell growth, alteration in the typical polymorphism of wild-type cells. Increased phagocytosis and decreased virulence in the mutant cells in a mouse model of infection were also observed (Almeida *et al.*, 2009). Several subsequent studies using this technology have allowed the role of various molecules during host–fungus interactions to be defined; these molecules include adhesins (PbHAD32 and GP43), stress proteins (PbAOX), heat shock proteins (HSP90) and other important antigens, such as Pb27 (Hernández *et al.*, 2010, 2012; Ruiz *et al.*, 2011; Tamayo *et al.*, 2013; Torres *et al.*, 2013, 2014; Torres-Gómez *et al.*, 2013; Bailão *et al.*, 2014). It is noteworthy that using these mutants (silenced isolates) in both *in vitro* and *in vivo* (animal) models allowed the involvement of these molecules in several processes, including conidia-to-yeast transition, adhesion and cell homeostasis to be confirmed. The mutant strains showed decreased virulence in all cases. One of the most important results was that the targeted genes remained silenced for long periods, even up to a year after several cultures, or after recovery from animal tissues several months after infection. The expression levels of the majority of genes knocked down using the antisense technology are shown in Figure 3.

An additional strategy to silence genes has been implemented for *Paracoccidioides*; use of RNAi technology to decrease the expression levels of PbGP43 and PbP27 genes indicated reduced expression of these genes in mutant strains, albeit only during the first 20 days after selection, indicating that gene silencing by this methodology was not stable over time (Torres *et al.*, 2013). In a later attempt to improve RNAi technology in *Paracoccidioides*, successful knockdown of GP43 was achieved through the expression of intron-containing hairpin RNA (ihpRNA). Employing this strategy reduced Gp43 transcript levels by 73.1% (Goes *et al.*, 2014).

The implementation of molecular tools for both diagnosis and research has allowed significant





**Figure 3.** Generation of knocked-down strains of *P. brasiliensis*. Gene expression levels of various in wild-type (PbWT), WT transformed with the empty vector (PbEV) and *P. brasiliensis* yeast transformed with the antisense RNA (Pb-X-aRNA) after subculture for 120 days. Gene expression levels obtained by RT-PCR were normalized to the internal reference, Tubulin-2 (TUB2); \* $p < 0.05$  compared with PbWT and PbEV

advances to be made in both fields. However, new tools will need to be developed to evaluate several genes at the same time at functional levels, since host–parasite interactions generate complex responses. Further studies are needed to elucidate the role of genes involved in various metabolic processes, as well as those involved in the response to the host attacks and others encoding for virulence factors.

### Virulence factors

As with other dimorphic fungal pathogens, *Paracoccidioides* spp. exhibits several virulence mechanisms that allow it to combat and overcome host defences (Rappleye and Goldman, 2006). Two main virulence mechanisms are critical for the establishment of disease, the first being the ability of the fungus to adhere to both extracellular matrix (ECM) proteins (mainly laminin, fibronectin, fibrinogen, collagen) and epithelial pulmonary cells. This adherence mechanism is mediated by molecules, named adhesins, present on the fungal surface that allow it not only to adhere but also to invade the host tissues, contributing to development of the mycosis. Thus, several adhesins have been reported, including the 14–3–3 protein enolase, triose phosphate isomerase (TPI), glyceraldehyde 3-phosphate dehydrogenase (GAPDH), the glycoprotein Gp43 (the main immunodominant antigen) and a hydrolase (HAD32) (de Oliveira *et al.*,

2015; Pereira *et al.*, 2007; Barbosa *et al.*, 2006; Torres *et al.*, 2013; Mendes-Giannini *et al.*, 2006; Hernández *et al.*, 2010). Moreover, the 14–3–3 protein and Gp43 induce apoptosis in human pneumocytes (Silva *et al.*, 2015). The second virulence mechanism is the conidia/mycelium-to-yeast transition, a morphological process due not only to the temperature shift to 37 °C but also to the availability of organic sulphur compounds (Menino *et al.*, 2013b). A negative regulator of the inorganic sulphur assimilation pathway has also been described. This molecule (SconCp) is considered to be a novel virulence determinant in *P. brasiliensis* (Menino *et al.*, 2013b).

Other virulence mechanisms have also been described. Thus,  $\alpha$ 1,3-glucan, present in the yeast cell wall, has been suggested to form a 'protective shield' against host defence mechanisms (San-Blas *et al.*, 1977; Rappleye and Goldman, 2006). Melanin, a pigment produced by different pathways, is present in *P. brasiliensis* and is considered to be an important virulence factor that reduces susceptibility to host defence mechanisms and antifungal drugs (Taborda *et al.*, 2008).

Biofilm formation is generally associated with antimicrobial resistance and avoidance of host defences. More recently, it has been reported that *P. brasiliensis* yeasts are able to form biofilms and that their presence is associated with overexpression of adhesins and enzymes (Sardi *et al.*, 2015).

Other virulence determinants include metalloproteinases, phospholipases, an alternative oxidase (AOX; an enzyme involved in intracellular redox

balancing), p27 and heat-shock proteins (Tristão *et al.*, 2015; Soares *et al.*, 2011; Ruiz *et al.*, 2011; Tamayo *et al.*, 2013; Torres *et al.*, 2014). In addition, *P. brasiliensis* is capable of producing an exocellular serine–thiol proteinase which can degrade the basement membrane, allowing the fungus to invade and disseminate to other organ systems (Puccia *et al.*, 1998).

## Animal models

While taking into account that the exact moment when the host acquires the infection remains unknown, animal models have provided important tools to allow host–fungus interactions to be studied, from the beginning of infection to the development of chronic stages that mimic development in humans. Several animal models have been devised and various fungal strains of different morphotypes (conidia or yeast) and virulence have also been employed. Thus, animal models for this mycosis include mice (Calich *et al.*, 1985; Cano *et al.*, 2000; Defaveri *et al.*, 1982; McEwen *et al.*, 1987; Moscardi and Franco, 1980), hamsters (Essayag *et al.*, 2002; Iabuki and Montenegro, 1979; Peraçoli *et al.*, 1982), guinea pigs (Fava-Netto *et al.*, 1961) and rats (Iovannitti *et al.*, 1999). The mouse model is the one that has been the most used. Different inbred strains of mice inoculated intraperitoneally with yeast cells of *P. brasiliensis* showed significantly varying patterns of susceptibility. The A/SN strain was found to be the most resistant, while BIOD2/nSn, BIO.A and BIOD2/oSn were the most susceptible strains. These susceptibility differences were independent of the size of challenge inocula and histocompatibility complexes. However, gender is a factor, since male BALB/c and BIOD2/nSn mouse strains were more susceptible to infection than those of females (Calich *et al.*, 1985).

The same group subsequently established a genetically controlled murine model of PCM using isogenic mice, which allowed several parameters of the host–parasite interactions to be investigated. In this model, resistant mice inoculated intraperitoneally or intratracheally with *P. brasiliensis* yeast showed efficient macrophage activation, Delayed-type Hypersensitivity (DTH) response, low levels of specific antibodies and a tendency towards

resolution of the infection process, suggesting the development of a Th1 immune response. By contrast, susceptible mice developed a predominantly Th2 immune response with inefficient macrophage activation, depressed DTH reactions, high levels of antibodies, greater dissemination of the fungus and the development of chronic progressive disease (Singer-Vermes *et al.*, 1993a, 1993b; Cano *et al.*, 1995). Furthermore, athymic or nude (nu/nu) and euthymic (nu/+) mice have been evaluated in order to evaluate the involvement of T cells. Similar participation of inflammatory cells (neutrophils and macrophages) as well as levels of ECM protein production were observed in the two mouse types. The only difference was that the euthymic animals showed lesions with a more pronounced tendency to encapsulate, increased ECM proteins and higher titres of specific antibodies, indicating that T cells could participate in the containment and control of infection (Lenzi *et al.*, 1994; Burger *et al.*, 1996a, 1996b). In addition to humoral and cellular studies, other investigations have been carried out to evaluate the innate immune response in PCM, in order to understand the initial host–fungus interactions. These studies have allowed the study of mechanisms such as complement protein activation, the microbicidal activity of natural killer cells and phagocytes, and production of inflammatory eicosanoids, cytokines and chemokines (Calich *et al.*, 2008).

More recently, knockout mice have allowed the role of several molecules that participate in host–*Paracoccidioides* interactions to be elucidated. These include interleukin (IL)-10, IL-12, IL-18 (Costa *et al.*, 2013; Livonesi *et al.*, 2008; Ketelut-Carneiro *et al.*, 2015), nitric oxide (Bernardino *et al.*, 2013), dectin-1 (Loures *et al.*, 2014); MyD88, Toll-like receptor (TLR)-2, TLR-4, TLR-9 (González *et al.*, 2008c; Loures *et al.*, 2009, 2015; Menino *et al.*, 2013a), CCR-5 (Moreira *et al.*, 2008a, 2008b) and caspase-1 (Ketelut-Carneiro *et al.*, 2015).

The observation that PCM is more frequent in males than females was attributed to the protection conferred by the female hormone 17- $\beta$ -oestradiol, an elegant animal study being performed to confirm this hypothesis. The course of infection in male and female mice in relation to their hormonal status indicated that, after infection with *P. brasiliensis* conidia, normal males showed progressive infection, whereas normal females restricted

proliferation and progressive disease. Castrated animals exhibited a lesser capacity to restrict disease progression. In addition, castrated male mice reconstituted with 17- $\beta$ -oestradiol initially restricted proliferation, whereas castrated female mice reconstituted with testosterone were unable to restrict disease. These results confirmed that 17- $\beta$ -oestradiol confers protection during initial infection in females (Aristizábal *et al.*, 2002).

### Host defence mechanisms against *Paracoccidioides* spp.

Most knowledge concerning the host defence mechanism against *Paracoccidioides* spp. comes from the infected host (particularly animal models) and from *in vitro* studies using both primary cell or cell line cultures. Different cell types are involved in the defence mechanism against this fungal pathogen, including macrophages, neutrophils, dendritic cells (DCs), T lymphocytes and natural killer (NK) cells. Macrophages, neutrophils and DCs are classified as 'professional phagocytes' and participate in the innate and acquired phases of immunity. Their activation is fundamental to the control of pathogen growth. Both normal macrophages and neutrophils are permissive to *P. brasiliensis* growth, while phagocytes activated by cytokines are able to restrain fungal multiplication (Pina *et al.*, 2008; González *et al.*, 2003, 2000; Rodrigues *et al.*, 2007; Tavian *et al.*, 2008; Kurita *et al.*, 2005). The fungicidal activity of these phagocytes is mediated by nitric oxide or reactive oxygen species (ROS) production, these being molecules that deplete fundamental metabolites for fungal cell cycling and subsequent growth (Pina *et al.*, 2008; González *et al.*, 2003, 2000; Rodrigues *et al.*, 2007; Tavian *et al.*, 2008). In addition to NO and ROS production, the indoleamine 2,3-dioxygenase (IDO; a enzyme that catalyses tryptophan metabolism and is produced mainly by macrophages and DCs) participates in the control of fungal infection (Araújo *et al.*, 2014).

The development and activation of dendritic cells are associated with secretion of IFN $\gamma$ , IL-4 and IL-17 and increased development of T-reg cells, with all of them exerting a positive effect in controlling fungal infection (Pina *et al.*, 2013).

Besides the phagocytosis mechanism, neutrophils release structures called extracellular traps (NETs), which are composed of nuclear (decondensed DNA and histones) and granular material such as elastase; more recently, we have demonstrated that although *P. brasiliensis* is able to induce NETs, this mechanism is ineffective in killing the fungus (Mejía *et al.*, 2015).

T cells comprise different subpopulations that include mainly CD4<sup>+</sup>, CD8<sup>+</sup> and T-reg (CD4<sup>+</sup>CD25<sup>+</sup>PhoxP3<sup>+</sup>) cells. Both CD4<sup>+</sup> and CD8<sup>+</sup> T cells are necessary for controlling paracoccidioidal infection; in this case, CD8<sup>+</sup> T cells control fungal load, while CD4<sup>+</sup> T cells regulate antibody production and DTH reactions (Cano *et al.*, 2000; Chiarella *et al.*, 2007). Depletion of T-reg cells led to a less severe infection in mice, indicating that these cells have a suppressive effect on the immune response, preventing tissue pathology by limiting the inflammatory reaction (Felonato *et al.*, 2012). Nonetheless, the host susceptibility pattern appears to account for functionality of these cells; thus, CD4<sup>+</sup> T cells play a protective role in the resistant and intermediate mouse strains, whereas in susceptible mice they are deleted or anergic. These facts indicate that genetic resistance to PCM is associated with concomitant CD4<sup>+</sup> and CD8<sup>+</sup> T cell immunity, secreting type 1 and type 2 cytokines (Cano *et al.*, 2000; Chiarella *et al.*, 2007). Resistant mice develop higher numbers of more potent T-reg cells than susceptible individuals (Felonato *et al.*, 2012).

Moreover, NK cells accomplish two fundamental functions: (a) they are able to kill *P. brasiliensis* cells directly or recognize and kill infected cells through a cytotoxic mechanism; (b) they produce pro-inflammatory cytokines (IFN $\gamma$  and TNF $\alpha$ ) that in turn influence the acquired immune response against this fungus (Longhi *et al.*, 2012).

Interactions between immune host cells (mainly phagocytic cells) and *Paracoccidioides* spp. are mediated by recognition of conserved fungal structures, known as pathogen-associated molecular patterns (PAMPs), by means of germline-encoded pattern recognition receptors (PRRs) present on the host cell surface (Leibundgut-Landmann *et al.*, 2012; Brown, 2011). PRRs comprise several molecules, including Toll-like receptors (TLR), the C-type lectin receptor (CRL) dectin-1, mannose receptors (MR) and complement receptors (CR), among others. This interaction is critical for

priming an appropriated immune response with synthesis of pro-inflammatory cytokines and chemokines, ROS production and phagocytosis activation; nevertheless, PPR signalling could be beneficial or detrimental to the host, depending on its background. Thus, dectin-1, MR, TLR-2 and TLR-4 control lymphocyte proliferation in *P. brasiliensis* infection, or could exert negative effects by inducing a more severe infection, leading to tissue damage for an uncontrolled or enhanced pro-inflammatory immune response (Loures *et al.*, 2009, 2010, 2015).

As described above, protective immunity to *Paracoccidioides* infection was shown to be mediated by IFN $\gamma$ , IL-12 and TNF $\alpha$ , using gene knockout or cytokine-depleted mice (L. E. Cano *et al.*, 1998; Arruda *et al.*, 2002; Bernardino *et al.*, 2013); while IL-4 and IL-10 apparently both show detrimental effects (Pina *et al.*, 2004; Costa *et al.*, 2013). In addition to cytokines, lipid mediators such as leukotrienes participate in host defence against *Paracoccidioides*; thus, mice deficient in 5-lipoxygenase, a key enzyme that catabolizes the arachidonic acid present in the cell membrane, show increased fungal burdens and mortality (Santos *et al.*, 2013).

## Concluding remarks and future directions

New advances in *Paracoccidioides* biology have been described. These include the description of *P. lutzii* as a new species and annotation of the genome which is already available. Thus, the implementation of novel genetic and molecular tools and the availability of the entire genomic sequence could lead to remarkable progress in the study of virulence determinants and the search for targets in the design of new drugs and diagnostic assays, with a view to improving management of this mycosis.

Nevertheless, several questions remain to be answered regarding the pathogenesis of this dimorphic fungus, its adaptation mechanisms to the human host and how the host defence mechanisms work, as well as its diverse metabolic pathways.

## References

Albano AP, Klafke GB, Brandolt TM, *et al.* 2014. Wild animals as sentinels of *Paracoccidioides brasiliensis* in the state of Rio Grande do Sul, Brazil. *Mycopathologia* **177**: 207–215.

- Almeida AJ, Carmona JA, Cunha C, *et al.* 2007a. Towards a molecular genetic system for the pathogenic fungus *Paracoccidioides brasiliensis*. *Fungal Genet Biol* **44**: 1387–1398.
- Almeida AJ, Cunha C, Carmona JA, *et al.* 2009. Cdc42p controls yeast-cell shape and virulence of *Paracoccidioides brasiliensis*. *Fungal Genet Biol* **46**: 919–926.
- Almeida AJ, Martins M, Carmona JA, *et al.* 2006. New insights into the cell cycle profile of *Paracoccidioides brasiliensis*. *Fungal Genet Biol* **43**: 401–409.
- Almeida AJ, Matute DR, Carmona JA, *et al.* 2007b. Genome size and ploidy of *Paracoccidioides brasiliensis* reveals a haploid DNA content: flow cytometry and GP43 sequence analysis. *Fungal Genet Biol* **44**: 25–31.
- Araújo EF, Loures FV, Bazan SB, *et al.* 2014. Indoleamine 2,3-dioxygenase controls fungal loads and immunity in paracoccidioidomycosis but is more important to susceptible than resistant hosts. *PLoS Negl Trop Dis* **8**: e3330.
- Aristizábal BH, Clemons KV, Cock AM, *et al.* 2002. Experimental *Paracoccidioides brasiliensis* infection in mice: influence of the hormonal status of the host on tissue responses. *Med Mycol* **40**: 169–178.
- Arraes FB, Benoliel B, Burtet RT, *et al.* 2005. General metabolism of the dimorphic and pathogenic fungus *Paracoccidioides brasiliensis*. *Genet Mol Res* **30**: 290–308.
- Arruda C, Franco MF, Kashino SS, *et al.* 2002. Interleukin-12 protects mice against disseminated infection caused by *Paracoccidioides brasiliensis* but enhances pulmonary inflammation. *Clin Immunol* **103**: 185–195.
- Bagagli E, Franco M, Bosco SM, *et al.* 2003. High frequency of *Paracoccidioides brasiliensis* infection in armadillos (*Dasypus novemcinctus*): an ecological study. *Med Mycol* **41**: 217–223.
- Bagagli E, Theodoro RC, Bosco SM, McEwen JG. 2008. *Paracoccidioides brasiliensis*: phylogenetic and ecological aspects. *Mycopathologia* **165**: 197–207.
- Baida H, Biselli PJ, Juvenale M, *et al.* 1999. Differential antibody isotype expression to the major *Paracoccidioides brasiliensis* antigen in juvenile and adult form paracoccidioidomycosis. *Microbes Infect* **1**: 273–278.
- Bailão AM, Schrank A, Borges CL, *et al.* 2006. Differential gene expression by *Paracoccidioides brasiliensis* in host interaction conditions: representational difference analysis identifies candidate genes associated with fungal pathogenesis. *Microbes Infect* **8**: 2686–2697.
- Bailão EF, Parente JA, Pigosso LL, *et al.* 2014. Hemoglobin uptake by *Paracoccidioides* spp. is receptor-mediated. *PLoS Negl Trop Dis* **15**: e2856.
- Barbosa MS, Bão SN, Andreotti PF, *et al.* 2006. Glyceraldehyde-3-phosphate dehydrogenase of *Paracoccidioides brasiliensis* is a cell surface protein involved in fungal adhesion to extracellular matrix proteins and interaction with cells. *Infect Immun* **74**: 382–389.
- Barrozo LV, Benard G, Silva ME, *et al.* 2010. First description of a cluster of acute/subacute paracoccidioidomycosis cases and its association with a climatic anomaly. *PLoS Negl Trop Dis* **4**: e643.
- Belitardo DR, Calefi AS, Borges IK, *et al.* 2014a. Detection of antibodies against *Paracoccidioides brasiliensis* in free-range domestic pigs (*Sus scrofa*). *Mycopathologia* **177**: 91–95.
- Belitardo DR, Calefi AS, Sbeghen MR, *et al.* 2014b. *Paracoccidioides brasiliensis* infection in domestic rabbits (*Oryctolagus cuniculus*). *Mycoses* **57**: 222–227.
- Bellissimo-Rodrigues F, Bollela VR, Da Fonseca BA, Martinez R. 2013. Endemic paracoccidioidomycosis: relationship between



- clinical presentation and patients' demographic features. *Med Mycol* **51**: 313–318.
- Bellissimo-Rodrigues F, Machado AA, Martinez R. 2011. Paracoccidioidomycosis epidemiological features of a 1000-cases series from a hyperendemic area in the south-east of Brazil. *Am J Trop Med Hyg* **85**: 546–550.
- Benard G. 2008. An overview of the immunopathology of human paracoccidioidomycosis. *Mycopathologia* **165**: 209–221.
- Bernardino S, Pina A, Felonato M, et al. 2013. TNF $\alpha$  and CD8<sup>+</sup> T cells mediate the beneficial effects of nitric oxide synthase-2 deficiency in pulmonary paracoccidioidomycosis. *PLoS Negl Trop Dis* **7**: e2325.
- Bocca AL, Amaral AC, Teixeira MM, et al. 2013. Paracoccidioidomycosis: eco-epidemiology, taxonomy and clinical and therapeutic issues. *Future Microbiol* **8**: 1177–1191.
- Bonfim CV, Mamoni RL, Souza MH, Blotta L. 2009. TLR-2, TLR-4 and dectin-1 expression in human monocytes and neutrophils stimulated by *Paracoccidioides brasiliensis*. *Med Mycol* **47**: 722–733.
- Brock M. 2009. Fungal metabolism in host niches. *Curr Opin Microbiol* **12**: 371–376.
- Brown GC. 1999. Nitric oxide and mitochondrial respiration. *Biochim Biophys Acta* **1411**: 351–369.
- Brown GD. 2011. Innate antifungal immunity: the key role of phagocytes. *Annu Rev Immunol* **29**: 1–21.
- Brummer E, Castaneda E, Restrepo A. 1993. Paracoccidioidomycosis: an update. *Clin Microbiol Rev* **6**: 89–117.
- Burger E, Miyaji M, Sano A, et al. 1996a. Histopathology of paracoccidioidomycotic infection in athymic and euthymic mice: a sequential study. *Am J Trop Med Hyg* **55**: 235–242.
- Burger E, Vaz CC, Sano A, et al. 1996b. *Paracoccidioides brasiliensis* infection in nude mice: studies with isolates differing in virulence and definition of their T cell-dependent and T cell-independent components. *Am J Trop Med Hyg* **55**: 391–398.
- Calich VL, da Costa TA, Felonato M, et al. 2008. Innate immunity to *Paracoccidioides brasiliensis* infection. *Mycopathologia* **165**: 223–236.
- Calich VL, Singer-Vermees LM, Siqueira AM, Burger E. 1985. Susceptibility and resistance of inbred mice to *Paracoccidioides brasiliensis*. *Br J Exp Pathol* **66**: 585–594.
- Calle D, Rosero S, Orozco LC, et al. 2001. Paracoccidioidomycosis in Colombia: an ecological study. *Epidemiol Infect* **126**: 309–315.
- Cano LE, Gomez B, Brummer E, et al. 1994. Inhibitory effect of dexamethasone or macrophage activation on transformation of *Paracoccidioides brasiliensis* conidia ingested by macrophages: reversal by holotransferrin. *Infect Immun* **62**: 1494–1496.
- Cano LE, Kashino SS, Arruda C, et al. 1998. Protective role of  $\gamma$ -interferon in experimental pulmonary paracoccidioidomycosis. *Infect Immun* **66**: 800–806.
- Cano LE, Singer-Vermees LM, Costa TA, et al. 2000. Depletion of CD8<sup>+</sup> T cells *in vivo* impairs host defense of mice resistant and susceptible to pulmonary paracoccidioidomycosis. *Infect Immun* **68**: 352–359.
- Cano LE, Singer-Vermees LM, Vaz CA, et al. 1995. Pulmonary paracoccidioidomycosis in resistant and susceptible mice: relationship among progression of infection, bronchoalveolar cell activation, cellular immune response, and specific isotype patterns. *Infect Immun* **63**: 1777–1783.
- Cano MI, Cisalpino PS, Galindo I, et al. 1998. Electrophoretic karyotypes and genome sizing of the pathogenic fungus *Paracoccidioides brasiliensis*. *J Clin Microbiol* **36**: 742–747.
- Caro E, González A, Muñoz C, et al. 2008. Recognition of laminin by *Paracoccidioides brasiliensis* conidia: a possible mechanism of adherence to human type II alveolar cells. *Med Mycol* **46**: 795–804.
- Chequer-Bou-Habib D, Daniel-Ribeiro C, Banic DM, et al. 1989. Polyclonal B cell activation in paracoccidioidomycosis. Polyclonal activation in paracoccidioidomycosis. *Mycopathologia* **108**: 89–93.
- Chiarella AP, Arruda C, Pina A, et al. 2007. The relative importance of CD4<sup>+</sup> and CD8<sup>+</sup> T cells in immunity to pulmonary paracoccidioidomycosis. *Microbes Infect* **9**: 1078–1088.
- Colombo AL, Tobon A, Restrepo A, et al. 2011. Epidemiology of endemic systemic fungal infections in Latin America. *Med Mycol* **49**: 785–798.
- Corredor GG, Peralta LA, Castanão JH, et al. 2005. The naked-tailed armadillo *Cabassous centralis* (Miller 1899): a new host to *Paracoccidioides brasiliensis*. Molecular identification of the isolate. *Med Mycol* **43**: 275–280.
- Costa TA, Bazan SB, Feriotti C, et al. 2013. In pulmonary paracoccidioidomycosis IL-10 deficiency leads to increased immunity and regressive infection without enhancing tissue pathology. *PLoS Negl Trop Dis* **7**: e2512.
- Coutinho ZF, Silva D, Lazera M, et al. 2002. Paracoccidioidomycosis mortality in Brazil (1980–1995). *Cad Saude Publica* **18**: 1441–1454.
- de Arruda Grossklaus D, Bailão AM, Vieira Rezende TC, et al. 2013. Response to oxidative stress in *Paracoccidioides* yeast cells as determined by proteomic analysis. *Microbes Infect* **15**: 347–364.
- de Farias MR, Condas LA, Ribeiro MG, et al. 2011. Paracoccidioidomycosis in a dog: case report of generalized lymphadenomegaly. *Mycopathologia* **172**: 147–152.
- de Oliveira HC, da Silva JF, Scorzoni L, et al. 2015. Importance of adhesins in virulence of *Paracoccidioides* spp. *Front Microbiol* **6**: 303.
- Defaveri J, REXKALLAH-IWASSO MT, FRANCO MF. 1982. Experimental pulmonary paracoccidioidomycosis in mice: morphology and correlation of lesions with humoral and cellular immune response. *Mycopathologia* **77**: 3–11.
- Desjardins CA, Champion MD, Holder JW, et al. 2011. Comparative genomic analysis of human fungal pathogens causing paracoccidioidomycosis. *PLoS Genet* **7**: e1002345.
- Dias L, de Carvalho LF, Romano CC. 2012. Application of PCR in serum samples for diagnosis of paracoccidioidomycosis in southern Bahia-Brazil. *PLoS Negl Trop Dis* **6**: e1909.
- Ene IV, Heilmann CJ, Sorgo AG, et al. 2012. Carbon source-induced reprogramming of the cell wall proteome and secretome modulates the adherence and drug resistance of the fungal pathogen *Candida albicans*. *Proteomics* **12**: 3164–3179.
- Essayag SM, Landaeta ME, Hartung C, et al. 2002. Histopathologic and histochemical characterization of calcified structures in hamsters inoculated with *Paracoccidioides brasiliensis*. *Mycoses* **45**: 351–357.
- Fava-Netto C, De Brito T, Lacaz CS. 1961. Experimental South American blastomycosis of the guinea pig. *Pathol Microbiol* **24**: 192–206.
- Felipe MS, Andrade RV, Arraes FB, et al. 2005. Transcriptional profiles of the human pathogenic fungus *Paracoccidioides brasiliensis* in mycelium and yeast cells. *J Biol Chem* **280**: 24706–24714.
- Felonato M, Pina A, de Araujo EF, et al. 2012. Anti-CD25 treatment depletes T<sub>reg</sub> cells and decreases disease severity in susceptible and

- resistant mice infected with *Paracoccidioides brasiliensis*. *PLoS One* **7**: e51071.
- Ferreira JB, Navarro IT, Freire RL, *et al.* 2013. Evaluation of *Paracoccidioides brasiliensis* Infection in dairy goats. *Mycopathologia* **176**: 95–99.
- Fleck CB, Schöbel F, Brock M. 2011. Nutrient acquisition by pathogenic fungi: nutrient availability, pathway regulation, and differences in substrate utilization. *Int J Med Microbiol* **301**: 400–407.
- Fontana FF, Dos Santos CT, Esteves FM, *et al.* 2010. Seroepidemiological survey of paracoccidioidomycosis infection among urban and rural dogs from Uberaba, Minas Gerais, Brazil. *Mycopathologia* **169**: 159–165.
- Goes T, Bailão EF, Correa CR, *et al.* 2014. New developments of RNAi in *Paracoccidioides brasiliensis*: prospects for high-throughput, genome-wide, functional genomics. *PLoS Negl Trop Dis* **8**: e3173.
- González A, Caro E, Muñoz C, *et al.* 2008a. *Paracoccidioides brasiliensis* conidia recognize fibronectin and fibrinogen which subsequently participate in adherence to human type II alveolar cells: involvement of a specific adhesin. *Microb Pathog* **44**: 389–401.
- Gonzalez A, de Gregori W, Velez D, *et al.* 2000. Nitric oxide participation in the fungicidal mechanism of  $\gamma$ -interferon-activated murine macrophages against *Paracoccidioides brasiliensis* conidia. *Infect Immun* **68**: 2546–2552.
- González A, Gómez BL, Díez S, *et al.* 2005a. Purification and partial characterization of a *Paracoccidioides brasiliensis* protein with capacity to bind to extracellular matrix proteins. *Infect Immun* **73**: 2486–2495.
- González A, Lenzi HL, Motta EM, *et al.* 2005b. Expression of adhesion molecules in lungs of mice infected with *Paracoccidioides brasiliensis* conidia. *Microbes Infect* **7**: 666–673.
- Gonzalez A, Restrepo A, Cano LE. 2007. Role of iron in the nitric oxide-mediated fungicidal mechanism of IFN $\gamma$ -activated murine macrophages against *Paracoccidioides brasiliensis* conidia. *Rev Inst Med Trop Sao Paulo* **49**: 11–16.
- González A, Restrepo A, Cano LE. 2008b. Pulmonary immune responses induced in BALB/c mice by *Paracoccidioides brasiliensis* conidia. *Mycopathologia* **165**: 313–330.
- González A, Sahaza JH, Ortiz BL, *et al.* 2003. Production of pro-inflammatory cytokines during the early stages of experimental *Paracoccidioides brasiliensis* infection. *Med Mycol* **41**: 391–399.
- González A, Yáñez A, Gozalbo D, Gil ML. 2008c. MyD88 is dispensable for resistance to *Paracoccidioides brasiliensis* in a murine model of blood-borne disseminated infection. *FEMS Immunol Med Microbiol* **54**: 365–374.
- Hernández O, Almeida AJ, Gonzalez A, *et al.* 2010. A 32-kDa hydrolase plays an important role in *Paracoccidioides brasiliensis* adherence to host cells and influences pathogenicity. *Infect Immun* **78**: 5280–5286.
- Hernández O, Almeida AJ, Tamayo D, *et al.* 2012. The hydrolase PbHAD32 participates in the adherence of *Paracoccidioides brasiliensis* conidia to epithelial lung cells. *Med Mycol* **50**: 533–537.
- Hernández O, Araque P, Tamayo D, *et al.* 2015. Alternative oxidase plays an important role in *Paracoccidioides brasiliensis* cellular homeostasis and morphological transition. *Med Mycol* **53**: 205–214.
- Huberts DH, van der Kleij IJ. 2010. Moonlighting proteins: an intriguing mode of multitasking. *Biochim Biophys Acta* **1803**: 520–525.
- Hwang LH, Mayfield JA, Rine J, Sil A. 2008. *Histoplasma* requires SID1, a member of an iron-regulated siderophore gene cluster, for host colonization. *PLoS Pathog* **4**: e1000044.
- Iabuki K, Montenegro MR. 1979. Experimental paracoccidioidomycosis in the Syrian hamster: morphology, ultrastructure and correlation of lesions with presence of specific antigens and serum levels of antibodies. *Mycopathologia* **67**: 131–141.
- Iovannitti CA, Finkelievich JL, Negroni R, Elías-Costa MR. 1999. Histopathological evolution of experimental paracoccidioidomycosis in Wistar rats. *Zentralbl Bakteriol* **289**: 211–216.
- Jiménez MP, Restrepo A, Radzioch D, *et al.* 2006. Importance of complement 3 and mannose receptors in phagocytosis of *Paracoccidioides brasiliensis* conidia by Nramp1 congenic macrophages lines. *FEMS Immunol Med Microbiol* **47**: 56–66.
- Kashino SS, Fazioli RA, Cafalli-Favati C, *et al.* 2000. Resistance to *Paracoccidioides brasiliensis* infection is linked to a preferential Th1 immune response, whereas susceptibility is associated with absence of IFN $\gamma$  production. *J Interferon Cytokine Res* **20**: 89–97.
- Ketelut-Carneiro N, Silva GK, Rocha FA, *et al.* 2015. IL-18 triggered by the Nlrp3 inflammasome induces host innate resistance in a pulmonary model of fungal infection. *J Immunol* **194**: 4507–4517.
- Kurita N, Oarada M, Brummer E. 2005. Fungicidal activity of human peripheral blood leukocytes against *Paracoccidioides brasiliensis* yeast cells. *Med Mycol* **43**: 417–422.
- Leal CV, Montes BA, Mesa AC, *et al.* 2004. *Agrobacterium tumefaciens*-mediated transformation of *Paracoccidioides brasiliensis*. *Med Mycol* **42**: 391–395.
- Leibundgut-Landmann S, Wuthrich M, Hohl TM. 2012. Immunity to fungi. *Curr Opin Immunol* **24**: 449–458.
- Lenzi HL, Calich VL, Miyaji M, *et al.* 1994. Fibrosis patterns of lesions developed by athymic and euthymic mice infected with *Paracoccidioides brasiliensis*. *Braz J Med Biol Res* **27**: 2301–2308.
- Lima PS, Casaletti L, Bailão AM, *et al.* 2014. Transcriptional and proteomic responses to carbon starvation in *Paracoccidioides*. *PLoS Negl Trop Dis* **8**: e2855.
- Livonesi MC, Souto JT, Campanelli AP, *et al.* 2008. Deficiency of IL-12 p40 subunit determines severe paracoccidioidomycosis in mice. *Med Mycol* **46**: 637–646.
- Longhi LN, da Silva RM, Fornazim MC, *et al.* 2012. Phenotypic and functional characterization of NK cells in human immune response against the dimorphic fungus *Paracoccidioides brasiliensis*. *J Immunol* **189**: 935–945.
- Loures FV, Araújo EF, Feriotti C, *et al.* 2015. TLR-4 cooperates with Dectin-1 and mannose receptor to expand Th17 and Tc17 cells induced by *Paracoccidioides brasiliensis* stimulated dendritic cells. *Front Microbiol* **6**: 261.
- Loures FV, Araújo EF, Feriotti C, *et al.* 2014. Dectin-1 induces M1 macrophages and prominent expansion of CD8<sup>+</sup> IL-17<sup>+</sup> cells in pulmonary paracoccidioidomycosis. *J Infect Dis* **210**: 762–773.
- Loures FV, Pina A, Felonato M, Calich VL. 2009. TLR2 is a negative regulator of Th17 cells and tissue pathology in a pulmonary model of fungal infection. *J Immunol* **183**: 1279–1290.
- Loures FV, Pina A, Felonato M, *et al.* 2010. Toll-like receptor 4 signalling leads to severe fungal infection associated with enhanced proinflammatory immunity and impaired expansion of regulatory T cells. *Infect Immun* **78**: 1078–1088.
- Marcos CM, de Oliveira HC, da Silva JF, *et al.* 2014. The multifaceted roles of metabolic enzymes in the *Paracoccidioides* species complex. *Front Microbiol* **5**: 719.

- Marques SA. 2013. Paracoccidioidomycosis: epidemiological, clinical, diagnostic and treatment up-dating. *An Bras Dermatol* **88**: 700–711.
- Martins VP, Dinamarco TM, Soriani FM, *et al.* 2011. Involvement of an alternative oxidase in oxidative stress and mycelium-to-yeast differentiation in *Paracoccidioides brasiliensis*. *Eukariot Cell* **10**: 237–248.
- Mason MG, Nicholls P, Wilson MT, Cooper CE. 2006. Nitric oxide inhibition of respiration involves both competitive (heme) and noncompetitive (copper) binding to cytochrome *c* oxidase. *Proc Natl Acad Sci* **103**: 708e13.
- Matute DR, McEwen JG, Puccia R, *et al.* 2006. Cryptic speciation and recombination in the fungus *Paracoccidioides brasiliensis* as revealed by gene genealogies. *Mol Biol Evol* **23**: 65–73.
- McEwen JG, Bedoya V, Patiño MM, *et al.* 1987. Experimental murine paracoccidioidomycosis induced by the inhalation of conidia. *J Med Vet Mycol* **25**: 165–175.
- Mejía SP, Cano LE, López JA, *et al.* 2015. Human neutrophils produce extracellular traps against *Paracoccidioides brasiliensis*. *Microbiology* **161**: 1008–1017.
- Mendes-Giannini MJ, Andreotti PF, Vincenzi LR, *et al.* 2006. Binding of extracellular matrix proteins to *Paracoccidioides brasiliensis*. *Microbes Infect* **8**: 1550–1559.
- Mendes-Giannini MJ, Monteiro da Silva JL, de Fátima da Silva J, *et al.* 2008. Interactions of *Paracoccidioides brasiliensis* with host cells: recent advances. *Mycopathologia* **165**: 237–248.
- Menino JF, Almeida AJ, Rodrigues F. 2012. Gene knockdown in *Paracoccidioides brasiliensis* using antisense RNA. *Methods Mol Biol* **845**: 187–198.
- Menino JF, Saraiva M, Gomes-Alves AG, *et al.* 2013a. TLR9 activation dampens the early inflammatory response to *Paracoccidioides brasiliensis*, impacting host survival. *PLoS Negl Trop Dis* **7**: e2317.
- Menino JF, Saraiva M, Gomes-Rezende J, *et al.* 2013b. *P. brasiliensis* virulence is affected by SconC, the negative regulator of inorganic sulfur assimilation. *PLoS One* **8**: e74725.
- Missall TA, Pusateri ME, Donlin MJ, *et al.* 2006. Posttranslational, translational, and transcriptional responses to nitric oxide stress in *Cryptococcus neoformans*: implications for virulence. *Eukaryot Cell* **5**: 518–529.
- Montoya AE, Alvarez AL, Moreno MN, *et al.* 1999. Electrophoretic karyotype of environmental isolates of *Paracoccidioides brasiliensis*. *Med Mycol* **37**: 219–222.
- Moreira AP, Cavassani KA, Massafra-Tristão FS, *et al.* 2008a. CCR5-dependent regulatory T cell migration mediates fungal survival and severe immunosuppression. *J Immunol* **180**: 3049–3056.
- Moreira AP, Dias-Melicio LA, Peracoli MT, *et al.* 2008b. Killing of *Paracoccidioides brasiliensis* yeast cells by IFN $\gamma$  and TNF $\alpha$  activated murine peritoneal macrophages: evidence of H $_2$ O $_2$  and NO effector mechanisms. *Mycopathologia* **166**: 17–23.
- Moscardi M, Franco MF. 1980. Experimental paracoccidioidomycosis in mice. Immunopathological aspects of intraperitoneal infection. *Rev Inst Med Trop Sao Paulo* **22**: 286–293.
- Motoyama AB, Venancio EJ, Brandão GO, *et al.* 2000. Molecular identification of *Paracoccidioides brasiliensis* by PCR amplification of ribosomal DNA. *J Clin Microbiol* **38**: 3106–3109.
- Munk ME, Fazioli RA, Calich VL, Kaufmann SH. 1995. *Paracoccidioides brasiliensis*-stimulated human  $\gamma\delta$  T cells support antibody production by B cells. *Infect Immun* **63**: 1608–1610.
- Muñoz JF, Gallo JE, Misas E, *et al.* 2014. Genome update of the dimorphic human pathogenic fungi causing paracoccidioidomycosis. *PLoS Negl Trop Dis* **8**: e3348.
- Nevitt T. 2011. War-Fe-re: iron at the core of fungal virulence and host immunity. *Biomet. Int J Role Met Ions Biol Biochem Med* **24**: 547–558.
- Oliveira GG, Belitardo DR, Balarin MR, *et al.* 2013. Serological survey of paracoccidioidomycosis in cats. *Mycopathologia* **176**: 299–302.
- Oliveira GG, Navarro IT, Freire RL, *et al.* 2012. Serological survey of paracoccidioidomycosis in sheep. *Mycopathologia* **173**: 63–68.
- Oliveira GG, Silveira LH, Itano EN, *et al.* 2011. Serological evidence of *Paracoccidioides brasiliensis* infection in chickens from Paraná and Mato Grosso do Sul States, Brazil. *Mycopathologia* **171**: 197–202.
- Parente AF, Bailão AM, Borges CL, *et al.* 2011. Proteomic analysis reveals that iron availability alters the metabolic status of the pathogenic fungus *Paracoccidioides brasiliensis*. *PLoS One* **6**: e22810.
- Parente AF, Naves PE, Pigosso LL, *et al.* 2015. The response of *Paracoccidioides* spp. to nitrosative stress. *Microbes Infect* **17**: 575–85.
- Peraçoli MTS, Mota NGS, Montenegro MR. 1982. Experimental paracoccidioidomycosis in the Syrian hamster. Morphology and correlation of lesions with humoral and cell mediated immunity. *Mycopathologia* **79**: 7–17.
- Pereira LA, Bão SN, Barbosa MS, *et al.* 2007. Analysis of the *Paracoccidioides brasiliensis* triosephosphate isomerase suggests the potential for adhesin function. *FEMS Yeast Res* **7**: 1381–1388.
- Pina A, Bernardino S, Calich VL. 2008. Alveolar macrophages from susceptible mice are more competent than those of resistant mice to control initial *Paracoccidioides brasiliensis* infection. *J Leukoc Biol* **83**: 1088–1099.
- Pina A, de Araujo EF, Felonato M, *et al.* 2013. Myeloid dendritic cells (DCs) of mice susceptible to paracoccidioidomycosis suppress T cell responses whereas myeloid and plasmacytoid DCs from resistant mice induce effector and regulatory T cells. *Infect Immun* **81**: 1064–1077.
- Pina A, Valente-Ferreira RC, Molinari-Madlum EE, *et al.* 2004. Absence of interleukin-4 determines less severe pulmonary paracoccidioidomycosis associated with impaired Th2 response. *Infect Immun* **72**: 2369–2378.
- Pitz Ade F, Koishi AC, Tavares ER, *et al.* 2013. An optimized one-tube, semi-nested PCR assay for *Paracoccidioides brasiliensis* detection. *Rev Soc Bras Med Trop* **46**: 783–785.
- Price MS, Betancourt-Quiroz M, Price JL, *et al.* 2011. *Cryptococcus neoformans* requires a functional glycolytic pathway for disease but not persistence in the host. *MBio* **2**: e00103.
- Puccia R, Carmona AK, Gesztesi JL, Juliano L, Travassos LR. 1998. Exocellular proteolytic activity of *Paracoccidioides brasiliensis*: cleavage of components associated with the basement membrane. *MBio* **36**: 345–348.
- Queiroz-Telles F. 1994. *Paracoccidioides brasiliensis* ultrastructural findings. In *Paracoccidioidomycosis*, Franco M, Lacaz CS, Restrepo-Moreno A, *et al.* (eds). CRC Press: London; 27–44.
- Rappleye CA, Goldman WE. 2006. Defining virulence genes in the dimorphic fungi. *Annu Rev Microbiol* **60**: 281–303.
- Reis VC, Torres FA, Poças-Fonseca MJ, *et al.* 2005. Cell cycle, DNA replication, repair, and recombination in the dimorphic



- human pathogenic fungus *Paracoccidioides brasiliensis*. *Genet Mol Res* **4**: 232–250.
- Restrepo A, Gómez BL, Tobón A. 2012. Paracoccidioidomycosis: Latin America's own fungal disorder. *Curr Fungal Infect Rep* **6**: 303–311.
- Restrepo A, Gonzalez A, Agudelo C. 2011. Paracoccidioidomycosis. In *Essentials of Clinical Mycology*, Kauffman CA, Sobel JD, Pappas PG, *et al.* (eds), 2nd edn. Springer: New York; 367–386.
- Restrepo A, Tobón A, Cano L. 2015. Paracoccidioides brasiliensis. In *Mandell, Douglas and Bennett's Principles and Practice of Infectious Diseases*, Bennett JE, Dolin R, Blaser MJ (eds), 8th edn. Elsevier: Philadelphia, PA; 2995–3002.
- Ricci G, Mota FT, Wakamatsu A, *et al.* 2004. Canine Paracoccidioidomycosis. *Med Mycol* **42**: 379–383.
- Richini-Pereira VB, Bosco Sde M, Griese J, *et al.* 2008. Molecular detection of *Paracoccidioides brasiliensis* in road-killed wild-animals. *Med Mycol* **46**: 35–40.
- Rodrigues DR, Dias-Melicio LA, Calvi SA, *et al.* 2007. *Paracoccidioides brasiliensis* killing by IFN $\gamma$ , TNF $\alpha$  and GM-CSF activated human neutrophils: role for oxygen metabolites. *Med Mycol* **45**: 27–33.
- Rodrigues F, Ludovico P, Sousa MJ, *et al.* 2003. The spoilage yeast *Zygosaccharomyces bailii* forms mitotic spores: a screening method for haploidization. *Appl Environ Microbiol* **69**: 649–653.
- Ruiz OH, Gonzalez A, Almeida AJ, *et al.* 2011. Alternative oxidase mediates pathogen resistance in *Paracoccidioides brasiliensis* infection. *PLoS Negl Trop Dis* **5**: e1353.
- Salgado-Salazar C, Jones LR, Restrepo A, McEwen JG. 2010. The human fungal pathogen *Paracoccidioides brasiliensis* (Onygenales: Ajellomycetaceae) is a complex of two species: phylogenetic evidence from five mitochondrial markers. *Cladistics* **26**: 613–224.
- San-Blas F. 1986. Ultrastructure of spore formation in *Paracoccidioides brasiliensis*. *J Med Vet Mycol* **24**: 203–210.
- San-Blas G, San-Blas F, Serrano LE. 1977. Host–parasite relationships in the yeast-like form of *Paracoccidioides brasiliensis* strain IVIC Pb9. *Infect Immun* **15**: 343–346.
- Santos PC, Santos DA, Ribeiro LS, *et al.* 2013. The pivotal role of 5-lipoxygenase-derived LTB $_4$  in controlling pulmonary paracoccidioidomycosis. *PLoS Negl Trop Dis* **7**: e2390.
- Sardi JC, Pitangui NS, Voltan AR, *et al.* 2015. *In vitro* *Paracoccidioides brasiliensis* biofilm and gene expression of adhesins and hydrolytic enzymes. *Virulence* **6**: 642–651.
- Schrettl M, Bignell E, Kragl C, *et al.* 2004. Siderophore biosynthesis but not reductive iron assimilation is essential for *Aspergillus fumigatus* virulence. *J Exp Med* **200**: 1213–1219.
- Semighini CP, de Camargo ZP, Puccia R, *et al.* 2002. Molecular identification of *Paracoccidioides brasiliensis* by 5' nuclease assay. *Diagn Microbiol Infect Dis* **44**: 383–386.
- Silva JF, Vicentim JI, Oliveira HC *et al.* 2015. Influence of the *Paracoccidioides brasiliensis* 14–3–3 and gp43 proteins on the induction of apoptosis in A549 epithelial cells. *Mem Inst Oswaldo Cruz* **110**: 476–484.
- Silva-Bailão MG, Bailão EF, Lechner BE, *et al.* 2014. Hydroxamate production as a high affinity iron acquisition mechanism in *Paracoccidioides* spp. *PLoS One* **9**: e105805.
- Singer-Vermes LM, Burger E, Russo M, *et al.* 1993a. Advances in experimental paracoccidioidomycosis using an isogenic murine model. *Arch Med Res* **24**: 239–245.
- Singer-Vermes LM, Caldeira CB, Burger E, Calich LG. 1993b. Experimental murine paracoccidioidomycosis: relationship among the dissemination of the infection, humoral and cellular immune responses. *Clin Exp Immunol* **94**: 75–79.
- Soares DA1, Oliveira MB, Evangelista AF, *et al.* 2011. Phospholipase gene expression during *Paracoccidioides brasiliensis* morphological transition and infection. *Mem Inst Oswaldo Cruz* **108**: 808–811.
- Sturme MH, Puccia R, Goldman GH, Rodrigues F. 2011. Molecular biology of the dimorphic fungi *Paracoccidioides* spp. *Fungal Biol Rev* **25**: 89–97.
- Taborda CP, da Silva MB, Nosanchuk JD, Travassos LR. 2008. Melanin as a virulence factor of *Paracoccidioides brasiliensis* and other dimorphic pathogenic fungi: a mini-review. *Mycopathologia* **165**: 331–339.
- Tamayo D, Muñoz JF, Torres I, *et al.* 2013. Involvement of the 90 kDa heat shock protein during adaptation of *Paracoccidioides brasiliensis* to different environmental conditions. *Fungal Genet Biol* **51**: 34–41.
- Tavares AH, Fernandes L, Bocca AL, *et al.* 2015. Transcriptomic reprogramming of genus *Paracoccidioides* in dimorphism and host niches. *Fungal Genet Biol* **81**: 98–109.
- Tavares AH, Silva SS, Dantas A, *et al.* 2007. Early transcriptional response of *Paracoccidioides brasiliensis* upon internalization by murine macrophages. *Microbes Infect* **9**: 583–590.
- Tavian EG, Dias-Melicio LA, Acorsi MJ, *et al.* 2008. Interleukin-15 increases *Paracoccidioides brasiliensis* killing by human neutrophils. *Cytokine* **41**: 48–53.
- Teixeira MD, Theodoro RC, Oliveira FF, *et al.* 2013. *Paracoccidioides lutzii* sp. nov.: biological and clinical implications. *Med Mycol* **52**: 19–28.
- Teixeira MM, Theodoro RC, de Carvalho MJ, *et al.* 2009. Phylogenetic analysis reveals a high level of speciation in the *Paracoccidioides* genus. *Mol Phylogenet Evol* **52**: 273–283.
- Teles FR, Martins ML. 2011. Laboratorial diagnosis of paracoccidioidomycosis and new insights for the future of fungal diagnosis. *Talanta* **85**: 2254–2264.
- Theodoro RC, Candeias JM, Araújo JP, Jr, *et al.* 2005. Molecular detection of *Paracoccidioides brasiliensis* in soil. *Med Mycol* **43**: 725–729.
- Theodoro RC, Teixeira MM, Felipe MS, *et al.* 2012. Genus *Paracoccidioides*: species recognition and biogeographic aspects. *PLoS One* **7**: e37694.
- Torres I, Garcia AM, Hernandez O, *et al.* 2009. Presence and expression of the mating type locus in *Paracoccidioides brasiliensis* isolates. *Fungal Genet Biol* **47**: 373–380.
- Torres I, Hernandez O, Tamayo D, *et al.* 2013. Inhibition of PbGP43 expression may suggest that gp43 is a virulence factor in *Paracoccidioides brasiliensis*. *PLoS One* **8**: e68434.
- Torres I, Hernandez O, Tamayo D, *et al.* 2014. *Paracoccidioides brasiliensis* PbP27 gene: knockdown procedures and functional characterization. *FEMS Yeast Res* **14**: 270–280.
- Torres-Gómez I, Hernandez-Ruiz O, Muñoz JF, *et al.* 2013. RNAi technology targeting PbGP43 and PbP27 in *Paracoccidioides brasiliensis*. *Open J Genet* **3**: 1–8.
- Tristão GB, Assunção L do P, Dos Santos LP *et al.* 2015. Predicting copper-, iron-, and zinc-binding proteins in pathogenic species of the *Paracoccidioides* genus. *Front Microbiol* **5**: 761.